ISSN 0021-9673

CUMULATIVE AUTHOR AND SUBJECT INDEXES VOLUMES 301–325

JOURNAL OF

CHROMATOGRAPHY

INTERNATIONAL JOURNAL ON CHROMATOGRAPHY, ELECTROPHORESIS AND RELATED METHODS

EDITOR, Michael Lederer (Switzerland)
ASSOCIATE EDITOR, K. Macek (Prague)
EDITOR, SYMPOSIUM VOLUMES, E. Heftmann (Orinda, CA)
EDITORIAL BOARD

W. A. Aue (Halifax)
V. G. Berezkin (Moscow)
V. Betina (Bratislava)
A. Bevenue (Honolulu, HI)
P. Boček (Brno)
P. Boulanger (Lille)
A. A. Boulton (Saskatoon)
G. P. Cartoni (Rome)
S. Dilli (Kensington, N.S.W.)
L. Fishbein (Jefferson, AR)
R. W. Frei (Amsterdam)
A. Frigerio (Milan)
C. W. Gehrke (Columbia, MO)
E. Gil-Av (Rehovot)
G. Guiochon (Palaiseau)
I. M. Hais (Hradec Králové)
J. K. Haken (Kensington, N.S.W.)
S. Hjertén (Uppsala)
E. C. Horning (Houston, TX)
Cs. Horváth (New Haven, CT)
J. F. K. Huber (Vienna)
A. T. James (Sharnbrook)
J. Janák (Brno)
E. sz. Kováts (Lausanne)
K. A. Kraus (Oak Ridge, TN)
E. Lederer (Gif-sur-Yvette)
A. Liberti (Rome)
H. M. McNair (Blacksburg, VA)
Y. Marcus (Jerusalem)
G. B. Marini-Bettòlo (Rome)
A. J. P. Martin (Lausanne)
Č. Michalec (Prague)
R. Neher (Basel)
G. Nickless (Bristol)
J. Novák (Brno)
N. A. Parris (Wilmington, DE)
R. L. Patience (London)
P. G. Righetti (Milan)
O. Samuelson (Göteborg)
R. Schwarzenbach (Dübendorf)
G. Semenza (Zürich)
L. R. Snyder (Yorktown Heights, NY)
A. Zlatkis (Houston, TX)

EDITORS, BIBLIOGRAPHY SECTION
Z. Deyl (Prague), J. Janák (Brno), K. Macek (Prague)

UNIVERSITY OF HAWAII LIBRARIES
LIBRARY USE ONLY
PERIODICAL
QD 271 .J6
U.H. LIBRARY

ELSEVIER

JOURNAL OF CHROMATOGRAPHY

Scope. The *Journal of Chromatography* publishes papers on all aspects of chromatography, electrophoresis and related methods. Contributions consist mainly of research papers dealing with chromatographic theory, instrumental development and their applications. The section *Biomedical Applications,* which is under separate editorship, deals with the following aspects: developments in and applications of chromatographic and electrophoretic techniques related to clinical diagnosis (including the publication of normal values); screening and profiling procedures with special reference to metabolic disorders; results from basic medical research with direct consequences in clinical practice; combinations of chromatographic and electrophoretic methods with other physicochemical techniques such as mass spectrometry. In *Chromatographic Reviews,* reviews on all aspects of chromatography, electrophoresis and related methods are published.

Submission of Papers. Papers in English, French and German may be submitted, in three copies. Manuscripts should be submitted to: The Editor of *Journal of Chromatography,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands, or to: The Editor of *Journal of Chromatography, Biomedical Applications,* P.O. Box 681, 1000 AR Amsterdam, The Netherlands. Review articles are invited or proposed by letter to the Editors and will appear in *Chromatographic Reviews* or *Biomedical Applications.* An outline of the proposed review should first be forwarded to the Editors for preliminary discussion prior to preparation. Submission of an article is understood to imply that the article is original and unpublished and is not being considered for publication elsewhere. For copyright regulations, see below.

Subscription Orders. Subscription orders should be sent to: Elsevier Science Publishers B.V., P.O. Box 211, 1000 AE Amsterdam, The Netherlands. The *Journal of Chromatography* and the *Biomedical Applications* section can be subscribed to separately.

Publication. The *Journal of Chromatography* (incl. *Biomedical Applications, Chromatographic Reviews* and *Cumulative Author and Subject Indexes, Vols. 301–325*) has 34 volumes in 1985. The subscription prices for 1985 are:

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes, Vols. 301–325*) + *Biomed. Appl.* (Vols. 312–345):
Dfl. 5440.00 plus Dfl. 748.00 (postage) (total *ca.* US$ 2291.75)

J. Chromatogr. (incl. *Chromatogr. Rev.* and *Cum. Indexes, Vols. 301–325*) only (Vols. 312–335):
Dfl. 4320.00 plus Dfl. 582.00 (postage) (total *ca.* US$ 1795.50)

Biomed. Appl. only (Vols. 336–345):
Dfl. 1750.00 plus Dfl. 220.00 (postage) (total *ca.* US$ 729.75).

Journals are automatically sent by airmail at no extra costs to Australia, Brasil, Canada, China, Hong Kong, India, Israel, Japan, Malaysia, New Zealand, Pakistan, Singapore, South Africa, South Korea, Taiwan and the U.S.A. Back volumes of the *Journal of Chromatography* (Vols. 1 through 311) are available at Dfl. 204.00 (plus postage). Claims for issues not received should be made within three months of publication of the issue. If not, they cannot be honoured free of charge. Customers in the U.S.A. and Canada wishing information on this and other Elsevier journals, please contact Journal Information Center, Elsevier Science Publishing Co. Inc., 52 Vanderbilt Avenue, New York, NY 10017. Tel. (212) 916-1250.

Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Referativnyi Zhurnal and Science Citation Index.

See page 3 of cover for Publication Schedule, Information for Authors and information on Advertisements.

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1985

0021-9673/85/$03.30

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Printed in The Netherlands

JOURNAL
of
CHROMATOGRAPHY

INTERNATIONAL JOURNAL OF CHROMATOGRAPHY,
ELECTROPHORESIS AND RELATED METHODS

Cumulative Author and Subject Indexes Volumes 301–325 (1984–1985)

ELSEVIER
AMSTERDAM — OXFORD — NEW YORK — TOKYO
1985

Electronic data processing:
Büro für Satztechnik W. Meyer KG
Weissensberg, F.R.G.

© ELSEVIER SCIENCE PUBLISHERS B.V. — 1985 0021-9673/85/$03.30

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior written permission of the publisher, Elsevier Science Publishers B.V., P.O. Box 330, 1000 AH Amsterdam, The Netherlands.

Upon acceptance of an article by the journal, the author(s) will be asked to transfer copyright of the article to the publisher. The transfer will ensure the widest possible dissemination of information.

Submission of an article for publication implies the transfer of the copyright from the author(s) to the publisher and entails the authors' irrevocable and exclusive authorization of the publisher to collect any sums or considerations for copying or reproduction payable by third parties (as mentioned in article 17 paragraph 2 of the Dutch Copyright Act of 1912 and in the Royal Decree of June 20, 1974 (S. 351) pursuant to article 16 b of the Dutch Copyright Act of 1912) and/or to act in or out of Court in connection therewith.

Special regulations for readers in the U.S.A. This journal has been registered with the Copyright Clearance Center, Inc. Consent is given for copying of articles for personal or internal use, or for the personal use of specific clients. This consent is given on the condition that the copier pays through the Center the per-copy fee stated in the code on the first page of each article for copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law. The appropriate fee should be forwarded with a copy of the first page of the article to the Copyright Clearance Center, Inc., 27 Congress Street, Salem, MA 01970, U.S.A. If no code appears in an article, the author has not given broad consent to copy and permission to copy must be obtained directly from the author. All articles published prior to 1980 may be copied for a per-copy fee of US$ 2.25, also payable through the Center. This consent does not extend to other kinds of copying, such as for general distribution, resale, advertising and promotion purposes, or for creating new collective works. Special written permission must be obtained from the publisher for such copying.

Printed in The Netherlands

AUTHOR INDEX

VOLUMES 301—325

A

Abad, P., Poupet, A., Ponchet, M., Venard, P. and Bettachini, B.
Separation and quantitative assay of three pathogenesis-related (b)proteins from tobacco mosaic virus hypersensitive*Nicotiana tabacum* by reversed-phase HPLC 318(1985)417

Abadie, F., see Lecaillon, J.-B. 305(1984)411

Abbott, F.S., see Kapil, R.P. 307(1984)305

Abbott, R.W., see Gill, R. 301(1984)155

Abdel-Baky, S., see Nazareth, A. 314(1984)201

Abdel-Moety, E.M.
GLC determination of azintamide (Oragallin) in pharmaceutical formulations 324(1985)475

Abegaz, B. and Gerba, S.
Use of retention time plots as a means of identifying isomeric methyl ketononanoates 324(1985)440

Aben, G.V.A., see De Bruijn, C.H.M.M. 320(1985)205

Aben, G.V.A., see Everaerts, F.M. 320(1985)263

Aben, G.V.A., see Reijenga, J.C. 320(1985)245

Abernethy, D.R., Greenblatt, D.J., Ochs, H.R., Willis, C.R., Miller, D.D. and Shader, R.I.
Haloperidol determination in serum and cerebrospinal fluid using GLC with nitrogen-phosphorus detection: application to pharmacokinetic studies 307(1984)194

Abernethy, D.R., see Willis, C.R. 307(1984)200

Abidi, S.L.
High-efficiency resolution of isomeric rotenone compounds by HPLC 317(1984)383

Abidi, S.L.
Retention behavior of long chain quaternary ammonium homologues and related nitroso-alkylmethylamines 324(1985)209

Abramson, R.K., see Kiel, J.S. 320(1985)313

Ackman, R.G.
Silica gel particle size and coating thicknesses of Chromarods-S and -SII 323(1985)410

Ackman, R.G., see Banerjee, A.K. 319(1985)215

Adam, A.M. and Rogers, H.J.
GLC assay of aminoglutethimide and a HPLC assay for its acetyl metabolite in biological fluids 307(1984)129

Adam, M.J., see Grierson, J.R. 325(1985)103

Adamovics, J.
Rapid determination of metronidazole in human serum and urine using a normal-phase HPLC column with aqueous solvents 309(1984)436

Adams, A.K., Van Engelen, D.L. and Thomas, L.C.
Detection of GC eluates by simultaneous absorbance and fluorescence measurements 303(1984)341

Adams, M.A., see O'Kruk, R.J. 310(1984)343

Adams, N.W., see Lee, M.L. 302(1984)303

Agarwal, S.P. and Nwaiwu, J.
Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

Ager, R.P. and Oliver, R.W.A.
Separation of oestrogen conjugates in urine and synthetic mixtures by HPLC methods (Review) 309(1984)1

Agnew, M.P., see Lauren, D.R. 303(1984)206

Aguilar, M.I., see Hart, S.J. 306(1984)215

Aherne, G.W., see Burton, N.K. 309(1984)409

Airaudo, C.B., Gayte-Sorbier, A., Laurent, P. and Creusevau, R.
TLC of age resisters for elastomers (phenols and organic phosphites) (in French) 314(1984)349

Airoldi, L., see Carlucci, G. 311(1984)141

Aizawa, T., see Amagaya, S. 320(1985)430

Åkerblom, M.
Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Åkerlund, H.-E., see Johansson, G. 311(1984)277

Åkerlund, H.-E., see Svensson, P. 323(1985)363

Al-Saigh, Z.Y., see Card, T.W. 301(1984)261

Alasandro, M., Wiesler, D. and Novotny, M.
Effects of starvation and refeeding on the excretion of urinary steroid metabolites in

mice with different genetic background 308(1984)1
Albert, K., see Bayer, E. 312(1984)91
Albert, L.S.R., Mitchell, S.D. and Gray, D.O. Recovery of plant amines with carboxylic ion-exchange substrates and their subsequent analysis by TLC 312(1984)357
Albertsson, P.-Å., see Svensson, P. 323(1985)363
Alborn, H. and Stenhagen, G. Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47
Albro, P.W., Schroeder, J.S., Harvan, D.J. and Corbett, B.J. Characteristics of an extraction and purification procedure for chlorinated dibenzo-*p*-dioxins and dibenzofurans in soil and liver 312(1984)165
Alén, R., Niemelä, K. and Sjöström, E. GLC separation of hydroxy monocarboxylic acids and dicarboxylic acids on a fused-silica capillary column 301(1984)273
Alexander, G., see Varhegyi, G. 318(1985)247
Alexander, J.N. IV, McNally, M.E. and Rogers, L.B. LC fractionation of oligomers of nonylphenyl-oligo(ethylene glycol) 318(1985)289
Alexander, P.W., see Haddad, P.R. 315(1984)261, 321(1985)363, 324(1985)319
Alexander, R., see Rowland, S.J. 312(1984)395
Alexander, R., see Wajon, J.E. 319(1985)187
Ali, H.M., Bennett, J.L., Sulaiman, S.M. and Gaillot, J. Method for the determination of oltipraz, a new antischistosomal agent, in blood 305(1984)465
Ali, L.Z., see Sloan, D.L. 316(1984)43
Ali, S.G.A.H., Purnell, J.H. and Williams, P.S. Window analysis optimization of GC separations using mixed Porapaks 302(1984)119
Alken, R.G., see Grove, J. 306(1984)383
Allen, A.C., see Lurie, I.S. 317(1984)427
Allen, K.G., MacGee, J., Fellows, M.E., Tornheim, P.A. and Wagner, K.R. New procedure to analyze free fatty acids. Application to 20-mg brain tissue samples 309(1984)33
Allenmark, S., Bomgren, B. and Borén, H. Direct LC separation of enantiomers on immobilized protein stationary phases. IV. Molecular interaction forces and retention behaviour in chromatography on bovine serum albumin as a stationary phase 316(1984)617
Alley, C.C., see Brooks, J.B. 309(1984)269
Allgire, J.F., Juenge, E.C., Damo, C.P., Sullivan, G.M. and Kirchhoefer, R.D. HPLC determination of *d*-/*l*-epinephrine enantiomer ratio in lidocaine-epinephrine local anesthetics 325(1985)249
Alric, R., see Cociglio, M. 307(1984)351
Alten, R. von, see Heizmann, P. 310(1984)119
Alton, K.B., Leitz, F., Bariletto, S., Jaworsky, L., Desrivieres, D. and Patrick, J. HPLC assay for labetalol in human plasma using a PRP-1 column and fluorometric detection 311(1984)319
Alvinerie, M. and Toutain, P.L. Determination of methylprednisolone and methylprednisolone acetate in synovial fluid using HPLC 309(1984)385
Amagaya, S., Sugishita, E., Ogihara, Y., Ogawa, S., Okada, K. and Aizawa, T. Separation and quantitative analysis of 18α-glycyrrhetinic acid and 18β-glycyrrhetinic acid in Glycyrrhizae Radix by GLC 320(1985)430
Amagaya, S., see Shimizu, K. 307(1984)488
Amato, A., see Gagliardi, L. 315(1984)465, 325(1985)353
Amédée-Manesme, O., see Furr, H.C. 309(1984)299
Ames, M.M., see Miller, K.J. 307(1984)335
Amione, C., see Leone, L. 321(1985)407
Amoah, A.G.B., Gould, B.J. and Parke, D.V. Single-dose pharmacokinetics of perhexiline administered orally to humans 305(1984)401
Amourache, L. and Vijayalakshmi, M.A. Affinity chromatography of kid chymosin on histidyl-Sepharose 303(1984)285
Amsterdam, P.H. van, see Ruijten, H.M. 314(1984)183
Andary, C., Roussel, J.L., Rascol, J.P. and Privat, G. Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312
Anderegg, R.J., see LaBrosse, J.L. 314(1984)83, 314(1984)93
Anders, J.C., Theoharides, A.D., Thomas, L.M., Smyth, M.H. and Chung, H. HPLC method for the analysis of a candidate 8-aminoquinoline antileishmanial drug using oxidative electrochemical detection 311(1984)117
Andersen, Ø.M. and Francis, G.W. Simultaneous analysis of anthocyanins and anthocyanidins on cellulose thin layers 318(1985)450
Andersson, K., Hallgren, C., Levin, J.-O. and Nilsson, C.-A. Determination of aliphatic amines in air by reversed-phase HPLC using 1-naphthyl isocyanate derivatives 312(1984)482
Andersson, L. Fractionation of human serum proteins by immobilized metal affinity chromatography 315(1984)167
Andersson, L. Experimental evaluation of the errors in high-performance GPC measurements 325(1985)37
Andersson, M., see Johansson, G. 303(1984)39
Andrasko, J. Anion chromatography on a conventional

reversed-phase column using two UV detectors 314(1984)429
Andrawes, F., Holzer, G., Roedder, E., Gibson, E.K. Jr. and Oro, J.
GC analysis of volatiles in fluid and gas inclusions 302(1984)181
Andreani, A., see Guerra, M.C. 320(1985)281
Andreolini, F., Borra, C., Di Corcia, A. and Samperi, R.
Direct determination of valproate in minute whole blood samples 310(1984)208
Andrews, R.S., see Powles, P. 311(1984)434
Andronikashvili, T.T., see Hradil, J. 324(1985)277
Änggård, E., see Jones, A.W. 309(1984)73
Angi, M.R., see Bettero, A. 310(1984)390
Anjaneyulu, B., see Nene, S. 308(1984)334
Annapoorani, K.S., Damodaran, C. and Chandra Sekharan, P.
Solid-state fluorodensitometric quantitation of arylnaphthalene lignan lactones of *Cleistanthus collinus* 303(1984)296
Annesley, T., Giacherio, D. and Wilkerson, K.
Analysis of retinoids by HPLC using programmed gradient separation 305(1984)199
Antle, P.E., Goldberg, A.P. and Snyder, L.R.
Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1
Antle, P.E., see Goldberg, A.P. 316(1984)241
Antonsson, A.-M., Gyllenhaal, O., Kylberg-Hanssen, K., Johansson, L. and Vessman, J.
Monitoring of *S*- and *R*-tocainide in human plasma after heptafluorobutyrylation, separation on Chirasil-Val and electron-capture detection 308(1984)181
Aoki, N., see Hirokawa, T. 312(1984)11
Aoyama, H., see Obuchi, A. 312(1984)247
Aoyama, M., see Kodama, K. 311(1984)369
Aoyama, T., see Nakano, Y. 311(1984)390
Aoyama, Y., see Yamamoto, S. 303(1984)99
Apffel, J.A., Brinkman, U.A.T. and Frei, R.W.
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153
Arata, J., see Mikasa, H. 305(1984)204, 310(1984)401
Aravind, M.K., Miceli, J.N. and Kauffman, R.E.
Determination of ibuprofen by HPLC 308(1984)350
Aravind, M.K., see Bowman, D.B. 309(1984)209
Archer, A.W.
Determination of 3-amino-1,2,4-triazole (amitrole) in urine by ion-pair HPLC 303(1984)267
Ardalan, B., see Chandrasekaran, B. 309(1984)403
Ardenne, R.A.M. van, see De Raat, W.K. 310(1984)41
Argoudelis, C.J.
Isocratic LC method for the simultaneous determination of aspartame and other additives in soft drinks 303(1984)256
Aries, R.E., Gutteridge, C.S. and Macrae, R.
Pyrolysis-MS investigations of reversed-phase HPLC stationary phases 319(1985)285
Arima, M., see Konomi, H. 311(1984)375
Arita, T., see Miyazaki, K. 310(1984)219
Armstrong, D.J., see Kim, S.-G. 321(1985)175
Arnold, W., see Dell, D. 317(1984)483
Arpino, P., see Ignatiadis, I. 324(1985)87
Arpino, P.J.
Ten years of LC-MS 323(1985)3
Arranz Peña, M.I.
Rapid GC assay for serum thiopental 305(1984)210
Arrendale, R.F., see Snook, M.E. 324(1985)141
Arsov, A., Kostowa, W. and Mesrob, B.
TLC of acid dyes on polyamide layers (in German) 322(1985)267
Artemi, F., see Felici, M. 320(1985)435
Artigas, F., see Tusell, J.M. 306(1984)338
Arvidsson, T., Wahlund, K.-G. and Daoud, N.
Procedures for direct injections of untreated blood plasma into LC columns with emphasis on a pre-column venting technique 317(1984)213
Asanoma, M., see Terada, H. 318(1985)299
Ashoor, S.H. and Knox, M.J.
Identification of fish species by HPLC 324(1985)199
Ashton, R.W. and Sloan, D.L.
Enzymatic coupled assay procedures that employ HPLC: the synthesis of orotidylate from ribose 317(1984)245
Ashworth, A.J. and Price, G.J.
Mixed solvents in GLC. Activity coefficients for chloroform, dichloromethane and ethyl acetate in squalane-dinonyl phthalate mixtures at 303°K 324(1985)231
Asmus, P.A. and Landis, J.B.
Analysis of steroids in bulk pharmaceuticals by LC with light-scattering detection 316(1984)461
Assmann, G., see Schmitz, G. 307(1984)65, 307(1984)81, 320(1985)253
Audunsson, G. and Mathiasson, L.
Direct analysis of free amines in salt solutions at sub-ppm levels by GLC 315(1984)299
Autio, K., see Kanko, S. 324(1985)395
Avram, M.J. and Shanks, C.A.
Determination of D-tubocurarine chloride or metocurine iodide in human plasma by HPLC with UV detection 306(1984)398
Axelson, J.E., see Kapil, R.P. 307(1984)305
Axelson, M., see Tandon, R. 302(1984)1
Ayafor, J.F., see Kubo, I. 321(1985)246
Aybar-Batista, D., see Sloan, D.L. 316(1984)43
Azerad, R., see Maurs, M. 325(1985)444
Azhitsky, G.Y., see Troitsky, G.V. 324(1985)285
Azoulay, M., Desbene, P.-L., Frappier, F. and Georges, Y.
Use of LC-MS for the quantitation of dethiobiotin and biotin in biological samples 303(1984)272

Azumaya, H., see Lam, S. 302(1984)21

B

Baba, S., Shinohara, Y., Sano, H., Inoue, T., Masuda, S. and Kurono, M.
Application of HPLC with synchronized accumulating radioisotope detector to analysis of glyceryl trinitrate and its metabolites in rat plasma 305(1984)119
Baba, S., Utoh, M., Horie, M. and Mori, Y.
Revised method for the quantitative determination of 5-hydroxytryptamine in human plasma by GC-MS-selected ion monitoring 307(1984)1
Baba, Y., Yoza, N. and Ohashi, S.
Simultaneous determination of phosphate and phosphonate by flow injection analysis and HPLC with a series detection system 318(1985)319
Baba, Y., see Yoza, N. 325(1985)385
Babitch, J.A., see Sharma, S.K. 312(1984)435
Bächmann, K., see Reineke, F.J. 323(1985)323
Bachur, N.R., see Nakazawa, H. 307(1984)323
Bacic, A., Harris, P.J., Hak, E.W. and Clarke, A.E.
Capillary GC of partially methylated alditol acetates on a high-polarity bonded-phase vitreous-silica column 315(1984)373
Backhaus, J., Dingler, E. and Weyhenmeyer, R.
Assay of the antiarrhythmic compound stirocainide in plasma by fused-silica GLC and nitrogen-selective detection 307(1984)190
Bacopoulos, N.G., see Harris, P.Q. 309(1984)379
Badaisha, K.K., see Kohli, J.C. 320(1985)455
Badcock, N.R.
Micro-determination of ketoconazole in plasma or serum by HPLC 306(1984)436
Bailey, E., Farmer, P.B., Hoskins, J.A., Lamb, J.H. and Peal, J.A.
Determination of plasma phenytoin by capillary GC with nitrogen-phosphorus detection and with selective ion monitoring 310(1984)199
Bailey, J.E. Jr.
Purification of D&C Red No. 33 by preparative HPLC 314(1984)379
Bailey, J.E. Jr.
Determination of 1,3-diphenyltriazene and azobenzene in D&C Red No. 33 by solvent extraction and reversed-phase HPLC 321(1985)185
Bain, R., see Gordon, S.M. 322(1985)246
Baiocchi, C., see Mentasti, E. 322(1985)177
Baker, B.W., see Yeung, P.K.F. 303(1984)412
Baker, L.H., see Evans, L.J. Jr. 305(1984)163
Baldessarini, R.J., see Stoll, A.L. 307(1984)457
Baldwin, R.P., see Halbert, M.K. 306(1984)269
Balen, H.C.J.G. van, see Van Hogezand, R.A. 305(1984)470
Baltar, A., see Purser, C. 311(1984)135
Baltassat, P., see Boulieu, R. 307(1984)469
Baltensperger, U. and Hertz, J.
Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153
Banerjee, A.K., Ratnayake, W.M.N. and Ackman, R.G.
Enhanced response of more highly unsaturated lipids on TLC-Iatroscan flame ionization detection after exposing the developed Chromarods to iodine vapour 319(1985)215
Bank, B., see Pirkle, W.H. 316(1984)585
Baranowski, J.D.
HPLC separation of pungency components of ginger 319(1985)471
Barbaro, A.M., see Guerra, M.C. 320(1985)281
Barber, W.E. and Carr, P.W.
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography; factors that control retention and selectivity 301(1984)25
Barber, W.E. and Carr, P.W.
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography: factors that control sensitivity and detection 316(1984)211
Barberán, F.A.T., Tomás, F. and Ferreres, F.
TLC behaviour and chemical structure of 6- and 8-methoxy-5-hydroxyflavones 315(1984)101
Barchowsky, A., see Luzzi, F.A. 311(1984)291
Bardalaye, P.C., Wheeler, W.B. and Templeton, J.L.
Derivatization of Oryzalin for characterization by GC-MS 314(1984)450
Barends, D.M., Blauw, J.S., Mijnsbergen, C.W., Govers, C.J.L.R. and Hulshoff, A.
Pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene 322(1985)321
Bargnoux, H., see Lartigue-Mattei, C. 310(1984)407
Bariletto, S., see Alton, K.B. 311(1984)319
Barkan, S., see Yang, Z.-Y. 324(1985)444
Barkley, R.M., see Hutte, R.S. 302(1984)173
Barnes, R.M., see Slatkavitz, K.J. 302(1984)277
Barone, S., Stitzel, R.E. and Head, R.J.
Non-radiochemical procedure for the measurement of O-methylation of the stereoisomers of isoprenaline 310(1984)283
Barrett, G.M., see Brooks, C.J.W. 315(1984)119
Barron, R.E. and Fritz, J.S.
Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201
Barry, E.F., see Hubball, J.A. 302(1984)341
Bartha, Á., Vigh, G., Billiet, H.A.H. and De Galan, L.

Studies in reversed-phase ion-pair chromatography. IV. The role of the chain length of the pairing ion 303(1984)29
Bascom, C.C., see Sharifi, B.G. 324(1985)173
Basili, A., see Gagliardi, L. 315(1984)465, 325(1985)353
Basta, M.T., see Brooks, J.B. 309(1984)269
Batley, M., see McNicholas, P.A. 315(1984)451
Battistoni, R., see Grossi, G. 309(1984)214
Bauer, C., see Josić, D. 317(1984)35
Bauer, M., Degude, C. and Mailhe, L.
Simultaneous determination of chlorhexidine, tetracaine and their degradation products by ion-pair LC 315(1984)457
Bauer, S., Neupert, M. and Spiteller, G.
Characterization of artefacts produced by treatment of organic acids with diazomethane 309(1984)243
Baumann, R.A., see Gooijer, C. 312(1984)337
Baumann, R.A., see Meering, P.G. 310(1984)159
Bauza, M.T., Smith, R.V., Knutson, D.E. and Witter, F.R.
GC determination of pentoxifylline and its major metabolites in human breast milk 310(1984)61
Bayer, C., see Hermann, K. 312(1984)273
Bayer, E. and Albert, K.
Continuous-flow carbon-13 nuclear magnetic resonance spectroscopy 312(1984)91
Bayer, E., Küsters, E., Nicholson, G.J. and Frank, H.
GC separation of enantiomeric sulphur compounds on Chirasil-Val 320(1985)393
Bayne, W.F., see Gravallese, D.A. 310(1984)71
Bazyl'chick, V.V., Fedorov, P.I. and Polyakova, N.P.
GLC study of monoterpene hydrocarbons of conifer balsams and ring opening products of pinane and carane structure 321(1985)240
Beach, C.A., see Mays, D.C. 311(1984)301
Bear, G.R., Lawley, C.W. and Riddle, R.M.
Separation of sulfonate and carboxylate mixtures by ion-exchange HPLC 302(1984)65
Beattie, K.L., see Revich, G.G. 317(1984)283
Bechtold, H., Klein, F., Trenk, D. and Jähnchen, E.
Improved method for quantitative analysis of vitamin K_1 and vitamin K_1 2,3-epoxide in human plasma by electron-capture GLC 306(1984)333
Beck, O. and Pévet, P.
Analysis of melatonin, 5-methoxytryptophol and 5-methoxyindoleacetic acid in the pineal gland and retina of hamster by capillary column GC-MS 311(1984)1
Beck, R.H.F., see Praznik, W. 303(1984)417
Beckers, J.L.
Determination of absolute mobilities and p*K* values by isotachophoresis 320(1985)147
Bedford, G.K. and Chiong, M.A.
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 305(1984)183
Beem, E.P., Goormachtig, P.F.J., Hooghwinkel, G.J.M., Lisman, J.J.W. and Overdijk, B.
Preparative isolation of the polypeptide chains of human liver β-N-acetylhexosaminidase A by fast protein LC-ion-exchange chromatography 323(1985)439
Beer, J.O. de, see De Beer, J.O. 307(1984)475
Begutter, H., Klus, H. and Ultsch, I.
Capillary GC determination of volatile and tobacco-specific N-nitrosamines with the thermo-energy-analyzer (in German) 321(1985)475
Beland, F.A., Fullerton, N.F. and Heflich, R.H.
Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 308(1984)121
Belew, M., see Maisano, F. 321(1985)305
Bellon, G., Lundy, A.M., Malgras, A. and Borel, J.P.
Fluorometric evaluation of sarcosine in urine and serum 311(1984)405
Bellon, P., Canal, P., Bernadou, J. and Soula, G.
Use of electrochemical detection in the HPLC determination of hydroxylated ellipticine derivatives 309(1984)170
Ben-Avram, C.M., see Reeve, J.R. Jr. 321(1985)421
Benassi, C.A., see Bettero, A. 310(1984)390
Benbouzid, A., Guinet, R. and Cloppet, H.
Purification of soluble specific antigens of systemic candidiasis by antibody affinity chromatography 306(1984)117
Bender, A.O., see Tameesh, A.H.H. 321(1985)59
Benedek, K., Dong, S. and Karger, B.L.
Kinetics of unfolding of proteins on hydrophobic surfaces in reversed-phase LC 317(1984)227
Benedict, C.R. and Risk, M.
Determination of urinary and plasma dihydroxyphenylalanine by coupled-column HPLC with C_8 and C_{18} stationary phases 317(1984)27
Benet, L.Z., see Langendijk, P.N.J. 307(1984)371
Benet, L.Z., see Smith, P.C. 306(1984)315
Benjamin, D.M., see Willis, C.R. 307(1984)200
Bennett, J.C., see Bhown, A.S. 314(1984)467
Bennett, J.L., see Ali, H.M. 305(1984)465
Benoiton, N.L., see Steinauer, R. 325(1985)111
Benowitz, N.L., see Jacob, P. III 306(1984)173
Berens, M.E., Salmon, S.E. and Davis, T.P.
Quantitative analysis of prostaglandins in cell culture medium by high-resolution GC with electron-capture detection 307(1984)251
Beretta, E. and Vanazzi, G.
Determination of nanogram amounts of dicyclomine with GC and nitrogen-selective detection 308(1984)341
Berg, J.H.M. van den, see Glöckner, G. 317(1984)615
Bergamini, C., see Chiavari, G. 318(1985)427

Bergenhem, N., Carlsson, U. and Klasson, K.
Rapid ion-exchange chromatography for preparative separation of proteins. Application to porcine and bovine carbonic anhydrases 319(1985)59
Berger, J.A., see Lartigue-Mattei, C. 310(1984)407
Bergqvist, Y. and Eckerbom, S.
Improved GC method for the simultaneous determination of chloroquine and two metabolites using capillary columns 306(1984)147
Bermejo, J. and Guillen, M.D.
Study of Kováts retention indices of aliphatic saturated esters and their relation to the polarity of the stationary phase 318(1985)187
Bernadou, J., see Bellon, P. 309(1984)170
Bernard, D., Maurizis, J.C., Chollet, P., Chassagne, J. and Plagne, R.
Isolation and purification of HLA-DR antigens from tumour cells by affinity chromatography and chromatofocusing 308(1984)322
Bernareggi, A., Ratti, B. and Toselli, A.
Quantitative determination of premazepam in human plasma by HPLC 309(1984)415
Bernauer, K., see Su, H. 301(1984)461
Berridge, J.C. and Morrissey, E.G.
Automated optimisation of reversed-phase HPLC separations. An improved method using the sequential simplex procedure 316(1984)69
Berry, B.W., see Jamali, F. 310(1984)327
Berry, V.V.
Universal LC methods. V. "Pulsing" with weak eluent to resolve peaks 321(1985)33
Bertaux, O., see Valencia, R. 325(1985)207
Bertrand, C., see Felix, G. 319(1985)432
Bettachini, B., see Abad, P. 318(1985)417
Bettendorff, L., see Bontemps, J. 307(1984)283
Bettero, A., Angi, M.R., Moro, F. and Benassi, C.A.
Histamine assay in tears by fluorescamine derivatization and HPLC 310(1984)390
Beyerlein, F.H., see Sura, D.J. 314(1984)471
Bhamra, R.K., Flanagan, R.J. and Holt, D.W.
HPLC method for the measurement of mexiletine and flecainide in blood plasma or serum 307(1984)439
Bhatia, S.C. and Shanbhag, V.D.
Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325
Bhatnagar, R., see Nakazawa, H. 307(1984)323
Bhatnager, P.K., see Kabra, P.M. 307(1984)224
Bhattacharya, A., see Tripathi, D.N. 315(1984)417
Bhown, A.S. and Bennett, J.C.
Identification of phenylthiohydantoin derivatives of substituted amino acids encountered during amino acid sequence analysis by HPLC 314(1984)467
Biagi, G.L., see Guerra, M.C. 320(1985)281
Bianchetti, G., see Caqueret, H. 311(1984)199
Bianchetti, G., see Padovani, P. 308(1984)229
Bianchi Bosisio, A., see Cossu, G. 307(1984)103
Bianchine, J.R., see Mays, D.C. 311(1984)301
Bidleman, T.F., Simon, C.G., Burdick, N.F. and You, F.
Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448
Bier, M., see Mosher, R.A. 320(1985)23
Biffar, S., Greely, V. and Tibbetts, D.
Determination of penicillamine in encapsulated formulations by HPLC 318(1985)404
Biggs, W.R., see Fetzer, J.C. 322(1985)275
Bikle, D.D., see Halloran, B.P. 303(1984)229
Billiet, H.A.H., Drouen, A.C.J.H. and De Galan, L.
Rapid optimization of the concentration of the ion-pairing reagent in ion-pairing reversed-phase LC 316(1984)231
Billiet, H.A.H., see Bartha, A. 303(1984)29
Binder, H., Weis, L. and Janz, P.
Reaction gas chromatographic studies of 3-oxocarboxylic acid esters with a tautomeric enol form. I. A semi-quantitative study (in German) 303(1984)375
Binkhuyzen, D., see Eichholtz, P.C.N. 305(1984)438
Binsard, A.M., see Paphassarang, S. 319(1985)412
Biondi, P.A., Simonic, T., Secchi, C. and Ronchi, S.
Sensitive assay for diamine oxidase activity using HPLC 309(1984)151
Birks, J.W., see Sigvardson, K.W. 316(1984)507
Bischoff, R. and McLaughlin, L.W.
Mixed-mode chromatographic matrices for the resolution of transfer ribonucleic acids 317(1984)251
Björkman, S. and Idvall, J.
HPLC method for methohexital and thiopental in plasma or whole blood 307(1984)481
Blaas, W., see Tiebach, R. 318(1985)103, 323(1985)121
Blake, J.D., Clarke, M.L. and Richards, G.N.
Determination of D-gluconic, 5-keto-D-gluconic, 2-keto-D-gluconic and 2,5-diketo-D-gluconic acids by HPLC 312(1984)211
Blanchard, F., see Olek, M. 325(1985)239
Blanche, F. and Couder, M.
Use of isotachophoresis to monitor the synthesis and purification of L-α-amino-ε-ketopimelate 323(1985)451
Blaschke, T.F., see Smith, G.A. 307(1984)410
Blasius, E., Müller, K. and Ziegler, K.
Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161
Blatchford, C. and Malcolme-Lawes, D.J.
Electrochemiluminescence as a detection

technique for reversed-phase HPLC. I. Preliminary experiments 321(1985)227
Blauw, J.S., see Barends, D.M. 322(1985)321
Blitterswijk, H. van, see Jansen, E.H.J.M. 319(1985)436
Blom, Y., see Jones, A.W. 309(1984)73
Blomberg, L., see Buijten, J. 301(1984)265
Blomberg, L., see Markides, K. 302(1984)319
Blomberg, L.G., see Hoffmann, S. 302(1984)95
Blomberg, S., see Roeraade, J. 301(1984)454, 322(1985)55
Bobik, A., see Jackman, G.P. 308(1984)301
Bobik, A., see Oddie, C.J. 308(1984)370
Boček, P. and Foret, F.
Isotachophoresis in inorganic chemistry (Review) 313(1984)189
Boček, P., see Deml, M. 320(1985)159
Boček, P., see Dolník, V. 320(1985)89
Boček, P., see Everaerts, F.M. 320(1985)263
Boček, P., see Gebauer, P. 320(1985)49
Boček, P., see Pospíchal, J. 320(1985)139
Bock, J.L., Lam, S. and Karmen, A.
Therapeutic drug monitoring using high-speed LC and rapid sample preparation: an assay for serum theophylline 308(1984)354
Bockbrader, H.N. and Reuning, R.H.
Digoxin and metabolites in urine: a derivatization-HPLC method capable of quantitating individual epimers of dihydrodigoxin 310(1984)85
Boehm, T.M., see Young, P.M. 311(1984)79
Boiron, F., see Heape, A.M. 322(1985)391
Bologna, M., see Forchetti, C. 309(1984)177
Bomgren, B., see Allenmark, S. 316(1984)617
Bond, A.M., Heritage, I.D. and Briggs, M.H.
Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313
Bondesson, U., see Jones, A.W. 309(1984)73
Bonn, G.
HPLC elution behaviour of oligosaccharides, monosaccharides and sugar degradation products on series-connected ion-exchange resin columns using water as the mobile phase 322(1985)411
Bonnaud, B., Calmel, F., Patoiseau, J.-F., N'Guyen, N.-T. and Cousse, H.
Separation of diastereomeric amides by preparative HPLC and analysis of enantiomers on a chiral support (in French) 318(1985)398
Bontemps, J., Laschet, J., Dandrifosse, G., Van Cutsem, J.-L. and Forget, P.-P.
Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59
Bontemps, J., Philippe, P., Bettendorff, L., Lombet, J., Dandrifosse, G., Schoffeniels, E. and Crommen, J.
Determination of thiamine and thiamine phosphates in excitable tissues as thiochrome derivatives by reversed-phase HPLC on octadecyl silica 307(1984)283
Boo, A.T. and Krohn, J.
Combination of automated pre-column trace enrichment and on-line evaporation in HPLC 301(1984)335
Boobis, A.R., see Gulaid, A.A. 318(1985)393
Booth, C., see Styring, M.G. 319(1985)115
Boppana, V.K., Heineman, F.C., Lynn, R.K., Randolph, W.C. and Ziemniak, J.A.
Determination of fenoldopam (SK&F 82526) and its metabolites in human plasma and urine by HPLC with electrochemical detection 317(1984)463
Borea, P.A., see Guerra, M.C. 320(1985)281
Borel, J.P., see Bellon, G. 311(1984)405
Borén, H., see Allenmark, S. 316(1984)617
Borgmann, U., see Schmitz, G. 320(1985)253
Bories, G.F., see Peleran, J.C. 324(1985)469
Borra, C., Cosmi, G., Di Corcia, A. and Samperi, R.
Reliable measurement of non-esterified long-chain fatty acid pattern in blood plasma 311(1984)9
Borra, C., see Andreolini, F. 310(1984)208
Bory, C., see Boulieu, R. 307(1984)469
Bossche, W. van den, see De Croo, F. 325(1985)395
Bossche, W. van den, see De Spiegeleer, B. 315(1984)481
Bosserhoff, A., see Kamp, R.M. 317(1984)181
Both-Miedema, R., see Jansen, E.H.J.M. 319(1985)436
Botha, E., see Labadarios, D. 309(1984)375
Bottiglieri, T., Lim, C.K. and Peters, T.J.
Isocratic analysis of 3-methoxy-4-hydroxyphenyl glycol, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in cerebrospinal fluid by HPLC with amperometric detection 311(1984)354
Boukerche, H., see Clezardin, P. 319(1985)67
Boulieu, R., Bory, C., Baltassat, P. and Gonnet, C.
Simultaneous determination of allopurinol, oxipurinol, hypoxanthine and xanthine in biological fluids by HPLC 307(1984)469
Bouquet, S., see Fourtillan, J.-B. 309(1984)391
Bourne, D.W.A., see Schneider, J.J. 308(1984)359
Boutron, H., see Diquet, B. 311(1984)430
Bovalini, L., see Martelli, P. 301(1984)297
Bovenkamp, J.W., Lacroix, B.V. and Henshaw, P.F.
Retention times of oximes in reversed-phase HPLC 301(1984)492
Bowman, D.B., Aravind, M.K., Miceli, J.N. and Kauffman, R.E.
Reversed-phase HPLC method to determine ceftriaxone in biological fluids 309(1984)209
Bowyer, D.E., see Schmitz, G. 307(1984)81
Bradberry, C., see McKay, L. 311(1984)167
Bradshaw, J.S., see Lee, M.L. 302(1984)303
Brady, R.L., see Lacey, E. 315(1984)233

Branch, R.A., see Wedlund, P.J. 307(1984)121
Brandts, P.M., Gelsema, W.J. and De Ligny, C.L.
Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. I. Sorption isotherms of *n*-alcohols on octyl-agarose 322(1985)399
Brauw, M.C. ten Noever de, see Van der Greef, J. 323(1985)81
Bray, C.M., see Standard, S.A. 318(1985)433
Brazell, R.S. and Todd, R.A.
New design for helium ionization detection 302(1984)257
Breckenridge, A.M., see Coleman, M.D. 308(1984)363
Breckenridge, A.M., see Ward, S.A. 305(1984)239
Bree, H. de, see Ruijten, H.M. 314(1984)183
Breimer, D.D., see Kleinbloesem, C.H. 308(1984)209
Breyer-Pfaff, U., Wiatr, R. and Nill, K.
Measurement of maprotiline and oxaprotiline in plasma by HPLC of fluorescent derivatives 309(1984)107
Briggs, M.H., see Bond, A.M. 315(1984)313
Brimer, L. and Dalgaard, L.
Cyanogenic glycosides and cyanohydrins in plant tissues. Qualitative and quantitative determination by enzymatic post-column cleavage and electrochemical detection, after separation by HPLC 303(1984)77
Brimer, L., see Dalgaard, L. 303(1984)67
Brinkman, U.A.T., see Apffel, J.A. 312(1984)153
Brinkman, U.A.T., see Goewie, C.E. 301(1984)325
Brinkman, U.A.T., see Jansen, H. 325(1985)255
Brinkman, U.A.T., see Kok, W.T. 315(1984)271
Brinkman, U.A.T., see Maris, F.A. 323(1985)113
Brinkman, U.A.T., see Nielen, M.W.F. 317(1984)557
Brodie, R.R., Chasseaud, L.F. and Rooney, L.
Determination of bromopride in human plasma and urine by HPLC 310(1984)353
Brodie, R.R., see Walmsley, L.M. 311(1984)227
Bromet, N., see Nicot, G. 306(1984)279
Brondz, I. and Olsen, I.
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19
Brondz, I. and Olsen, I.
Determination of bound cellular fatty acids in *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* by GC and GC-MS 308(1984)282
Brondz, I. and Olsen, I.
Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261
Brondz, I. and Olsen, I.
Carbohydrates of whole defatted cells as a basis for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)31
Brondz, I. and Olsen, I.
Whole-cell methanolysis as a rapid method for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)347
Brooks, C.J.W., Cole, W.J. and Barrett, G.M.
Studies of cyclic di-*tert.* -butylsilylene derivatives of diols and hydroxy acids by GC-MS 315(1984)119
Brooks, J.B., Basta, M.T., Alley, C.C., Holler, J.S. and El Kholy, A.M.
Identification of diethylene glycol in sera from egyptian children by frequency-pulsed electron-capture GLC 309(1984)269
Brooks, M.A., see Lee, T.L. 306(1984)429
Brown, G.M., see Rosenfeld, J.M. 325(1985)309
Brown, H., see Risk, M. 317(1984)367
Brown, J.E., see Young, P.M. 311(1984)79
Brown, M.J., see Causon, R.C. 306(1984)257, 310(1984)11, 317(1984)319
Brown, N.D., Gray, R.R., Stermer-Cox, M.G., Doctor, B.P. and Hagedorn, I.
Stability study of HI-6 dichloride in various anticholinergic formulations 315(1984)389
Brown, N.D., Stermer-Cox, M.G., Poon, B.T. and Chulay, J.D.
Separation and identification of a plasma and urinary mono-acetylated conjugate of chloroquine in man by ion-pair HPLC 309(1984)426
Brown, N.D., see Nelson, L.D. 324(1985)203
Brown, P.R., see Perrone, P.A. 307(1984)53, 317(1984)301
Brown, R., see Timmons, L. 314(1984)476
Brown, S.A., see Thompson, H.J. 314(1984)323
Brown, S.J., see Harris, P.Q. 309(1984)379
Brubaker, W.F. Jr. and Ogliaruso, M.A.
GLC of substituted benzils and benzophenones 324(1985)450
Brubaker, W.F. Jr. and Prusoff, W.H.
Preparative and analytical HPLC methods in the synthesis and analysis of decomposition of nitrosourea nucleosides 322(1985)455
Bruijn, C.H.M.M. de, see De Bruijn, C.H.M.M. 320(1985)205
Bruijn, C.H.M.M. de, see Reijenga, J.C. 320(1985)245
Bruins, A.P.
Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99
Brummelen, P. van, see Kleinbloesem, C.H. 308(1984)209
Bruner, F., Furlani, G. and Mangani, F.
Sample enrichment for GC-MS analysis of polynuclear aromatic hydrocarbons in water and in organic mixtures 302(1984)167
Brunner, C., see Yang, Z.-Y. 324(1985)444
Buchanan, J.F., see Pollack, G.M. 311(1984)101
Buchman, O., Cao, G.-Y. and Peng, C.T.
Structure assignment by retention index in gas-liquid radiochromatography of substituted cyclohexenes 312(1984)75
Buchta, R.C., Wyles, H.F., Hensler, C.J., Van Lenten, F.J., Westerberg, R.B. and

Williams, L.A.
Determination of polychlorinated biphenyls in copper phthalocyanine pigments 325(1985)456
Buck, R.H. and Krummen, K.
HPLC with automated pre-column derivatization for amino acids 303(1984)238
Buck, R.H. and Krummen, K.
Resolution of amino acid enantiomers by HPLC using automated pre-column derivatisation with a chiral reagent 315(1984)279
Buckingham, D.A.
Reversed-phase high-performance ion-pair chromatography of cobalt(III) coordination complexes (Review) 313(1984)93
Bui-Nguyên, M.H.
Direct determination of D-panthenol in pharmaceutical preparations by ion-pair chromatography 303(1984)291
Buijten, J., Blomberg, L., Hoffmann, S., Markides, K. and Wännman, T.
Use of poly(silarylene-methylphenylsiloxane) block copolymer as a thermostable stationary phase in capillary column GC 301(1984)265
Buijten, J., see Hoffmann, S. 302(1984)95
Buijten, J., see Markides, K. 302(1984)319
Bullita, R., see Cossu, G. 307(1984)103
Burch, J.E., Roberts, S.G. and Raddats, M.A.
Amitriptyline and its basic metabolites determined in plasma by GC 308(1984)165
Burdick, N.F., see Bidleman, T.F. 301(1984)448
Burford, R.P., Haken, J.K. and Vimalasiri, P.A.D.T.
Chromatographic analysis of polyester-based polyurethane elastomers after alkali fusion 321(1985)295
Burford, R.P., see Vimalasiri, P.A.D.T. 319(1985)121
Burri, R., see Honegger, C.G. 309(1984)53
Bursey, J.T., see Voyksner, R.D. 312(1984)221, 323(1985)383
Bursill, D.B., see Nicholson, B.C. 325(1985)221
Burton, G., Veleiro, A.S. and Gros, E.G.
Reversed-phase chromatographic separation of withanolides from *Acnistus breviflorus* 315(1984)435
Burton, N.K., Aherne, G.W. and Marks, V.
Novel method for the quantitation of 6-mercaptopurine in human plasma using HPLC with fluorescence detection 309(1984)409
Buser, C., see Heizmann, P. 310(1984)119
Büser, W. and Erbersdobler, H.F.
Determination of lysinoalanine as the heptafluorobutyryl isobutyl ester derivative by GLC 303(1984)234
Bushway, R.J.
HPLC analysis of rotenone and rotenonone in water by direct injection 303(1984)263
Butler, H.T., see Poole, C.F. 302(1984)149
Butler, L.G., see Putman, L.J. 318(1985)85

C

Caboni, M.F., Lercker, G. and Ghe, A.M.
Analysis of phospholipids in cow's milk by high-temperature injection GC and HPLC 315(1984)223
Caddy, B., see Chow, W.M.L. 318(1985)255
Calder, I.C., see Hart, S.J. 306(1984)215
Calderone, R., see Cantwell, A.M. 316(1984)133
Calman, K.C., see Cummings, J. 311(1984)125
Calmel, F., see Bonnaud, B. 318(1985)398
Calvey, T.N., see Murray, G.R. 308(1984)143
Canal, P., see Bellon, P. 309(1984)170
Cani, D., see Marcucci, F. 309(1984)132
Cannon, M., see Timmons, L. 314(1984)476
Cano, J.-P., see Santoni, Y. 306(1984)165
Canselier, J.-P., see Castro, V. 325(1985)43
Cantelli Forti, G., see Guerra, M.C. 320(1985)281
Cantwell, A.M., Calderone, R. and Sienko, M.
Process scale-up of a β-lactam antibiotic purification by HPLC 316(1984)133
Cao, G.-Y., see Buchman, O. 312(1984)75
Caqueret, H. and Bianchetti, G.
Simple method for routine determination of betaxolol in blood and urine by automated HPLC with fluorimetric detection 311(1984)199
Caradonna, P., see Castagnola, M. 307(1984)91
Carcía-Domínguez, J.A., see Fernández-Sánchez, E. 312(1984)69
Card, T.W., Al-Saigh, Z.Y. and Munk, P.
Diffusion in the bubble flow meter in inverse GC experiments 301(1984)261
Carignan, G., Lodge, B.A. and Skakum, W.
HPLC analysis of estradiol valerate-testosterone enanthate in oily formulations 301(1984)292
Carignan, G., Lodge, B.A. and Skakum, W.
General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470
Carisano, A.
Rapid and sensitive method for the determination of proline by reversed-phase HPLC with automated pre-column fluorescence derivatization 318(1985)132
Carlebom, P., see Lagerström, P.-O. 307(1984)230
Carlsson, U., see Bergenhem, N. 319(1985)59
Carlucci, G., Airoldi, L., Fanelli, R. and Laguzzi, B.
Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141
Carlucci, G., see Forchetti, C. 309(1984)177
Carr, P.W., see Barber, W.E. 301(1984)25, 316(1984)211
Carson, P.E., see Nora, M.V. 307(1984)451
Carson, S.W. and Jusko, W.J.
Simultaneous analysis of cortexolone and

cortisol by HPLC for use in the metyrapone test 306(1984)345
Casoli, A., Mangia, A., Predieri, G. and Sappa, E.
HPLC of homo- and hetero-tetrametallic iron, ruthenium, osmium and nickel clusters 303(1984)404
Cassagne, C., see Heape, A.M. 322(1985)391
Cassiano, L., see Castagnola, M. 307(1984)91
Castagnetta, L., see D'Agostino, G. 305(1984)13
Castagnola, M., Caradonna, P., Cassiano, L., Degen, C., Lorenzin, F., Rossetti, D. and Salvi, M.L.
Analysis of the globins from fast human haemoglobins by isoelectrofocusing on polyacrylamide gel rods 307(1984)91
Castello, G.
GC identification of alkyl radicals formed in plasma radiofrequency discharges by using iodine as a scavenger 303(1984)61
Castello, G. and D'Amato, G.
GC separation and identification of linear and branched-chain alkyl bromides 324(1985)363
Castro, V. and Canselier, J.-P.
Analysis of α-olefinsulphonates by HPLC 325(1985)43
Catlow, D.A.
Applications of thermospray ionisation LC-MS to compounds of pharmaceutical interest 323(1985)163
Cattan, A., see Enger, A. 305(1984)335
Caude, M., see Souteyrand, C. 316(1984)373
Causon, R.C. and Brown, M.J.
Measurement of tyramine in human plasma, utilising ion-pair extraction and HPLC with amperometric detection 310(1984)11
Causon, R.C. and Brown, M.J.
HPLC with amperometric determination of plasma tyramine 317(1984)319
Causon, R.C., Desjardins, R., Brown, M.J. and Davies, D.S.
Determination of *d*-isoproterenol sulphate by HPLC with amperometric detection 306(1984)257
Cautreels, W., see Fels, J.-P. 308(1984)273
Cavallini, D., see Kodama, H. 311(1984)183
Cavazzutti, G., see Gagliardi, L. 315(1984)465, 325(1985)353
Cavicchio, G., see Forchetti, C. 309(1984)177
Čelap, M.B., see Vučković, G. 324(1985)431
Čellár, P., see Krupčík, J. 312(1984)1
Čellár, P., see Matisová, E. 303(1984)151
Centen, J.D., see Tong, H.Y. 324(1985)373
Chabard, J.L., see Lartigue-Mattei, C. 310(1984)407
Chadwick, V.S., see Selden, C. 305(1984)43
Chaláni, P., see Zelenský, I. 325(1985)161
Chan, K., see Murray, G.R. 308(1984)143
Chan, K.K.H., see Hui, K.-S. 325(1985)467
Chan, K.Y., Ohlweiler, D.F., Lang, J.F. and Okerholm, R.A.
Simultaneous analysis of a new cardiotonic agent, MDL 17,043, and its major sulfoxide metabolite in plasma by HPLC 306(1984)249
Chan, P., see Tyman, J.H.P. 303(1984)137
Chandra Sekharan, P., see Annapoorani, K.S. 303(1984)296
Chandrasekaran, B. and Ardalan, B.
Determination of 3-deazaguanine in mice plasma by HPLC 309(1984)403
Chang, C.A. and Fong, K.L.
Substitution-inert metal complex mobile phases in non-suppressed anion chromatography 312(1984)99
Chang, J.P.
Effect of surfactants on the separation of proteins by reversed-phase HPLC. I. Non-ionic surfactants (Tween) 317(1984)157
Chang, J.-P., El Rassi, Z. and Horváth, C.
Silica-bound polyethyleneglycol as stationary phase for separation of proteins by HPLC 319(1985)396
Chang, J.P., Melander, W.R. and Horváth, C.
Reversed-phase chromatography of proteins with diphenyl-silica column and hydro-organic eluents containing two organic solvents 318(1985)11
Chaplin, R.P., see Wainwright, M.S. 321(1985)287
Chapman, J.R.
LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153
Charikofsky, J.G., see Glajch, J.L. 318(1985)23
Charles, B.G., Ravenscroft, P.J. and Jacobsen, N.W.
Analysis of propantheline bromide in serum by HPLC 306(1984)424
Chase, T.N., see Knight, M. 301(1984)277
Chassagne, J., see Bernard, D. 308(1984)322
Chasseaud, L.F., see Brodie, R.R. 310(1984)353
Chasseaud, L.F., see Walmsley, L.M. 311(1984)227
Chatfield, J.M., see Kim, S.-G. 321(1985)175
Chatterji, D.C., see Narang, P.K. 305(1984)135
Chatterji, D.C., see Yeager, R.L. 305(1984)496
Chaumont, P.R. and Merrill, E.W.
Analytical solution for the area under chromatographic peaks obtained from the deconvolution method of Vaidya and Hester 303(1984)177
Checchini, L., see Desideri, P.G. 322(1985)107
Cheh, A.M., see Sweeney, A.G. 325(1985)95
Chen, B.-K., see Melander, W.R. 318(1985)1
Chen, E.C.M., see Hernandez-Gil, N. 312(1984)31
Chen, F.M.F., see Steinauer, R. 325(1985)111
Chen, M.-L., see Prueksaritanont, T. 306(1984)89
Chen, P., see Fogel, J. 317(1984)507
Chen, S.S., see Hsu, D.-S. 311(1984)396
Chen, S.S.-H. and Kou, A.Y.
HPLC of phospholipids using deuterated solvents for IR detection 307(1984)261
Chen, T.-M. and Contario, J.J.
HPLC resolution of enantiomers of γ-vinyl-γ-aminobutyric acid 314(1984)495

Cheng, L.K., see Krebs, H.A. 310(1984)412
Cheng, W. and McCown, M.
Effect of alkyl chain length on surface silanization of silica 318(1985)173
Chiavari, G. and Bergamini, C.
HPLC of carbonyl compounds as 2,4-dinitrophenylhydrazones with electrochemical detection 318(1985)427
Chiong, M.A., see Bedford, G.K. 305(1984)183
Chiou, W.L., see Prueksaritanont, T. 306(1984)89
Chism, G.W. III, Long, A.R. and Rolle, R.
Assay of enzymes involved in cytokinin metabolism by means of reversed-phase HPLC 317(1984)263
Choi, K.S., see Kim, K.R. 325(1985)127
Choi, R.L., see Rosenberg, M. 308(1984)382
Choi, S.-G., see Miyazaki, M. 323(1985)443
Chollet, P., see Bernard, D. 308(1984)322
Chooi, W.Y., see Stark, B.C. 319(1985)367
Chorev, M., see Laufer, R. 301(1984)415
Chortyk, O.T., see Snook, M.E. 324(1985)141
Chou, C.-K., see Lai, S.-T. 324(1985)436
Chou, R.C., see Wiegand, U.-W. 311(1984)218
Chovan, J.P., see Klett, R.P. 310(1984)361
Chow, C.K., see Collins, C.A. 317(1984)349
Chow, W.M.L. and Caddy, B.
Separation of underivatised barbiturates by capillary column GC. Preparation of medium polar polymethylsiloxane columns to optimize selectivity 318(1985)255
Chrambach, A.
Unified view of moving boundary electrophoresis: practical implications (plenary lecture) 320(1985)1
Chrétien, J.R., Szymoniak, K., Lion, C. and Haken, J.K.
GC of homologous esters. XXX. Unexpected predominancy of polar effects over steric effects in crowded aliphatic esters: a GLC appraisal of specific interactions 324(1985)355
Chrin, L., see Kelley, J. 311(1984)400
Christensen, R.G. and White, E. V
Determination of dibenzothiophene in oils by LC-tandem MS 323(1985)33
Christie, W.W. and Hunter, M.L.
Separation of molecular species of phosphatidylcholine by HPLC on a PLRP-S column 325(1985)473
Christopherson, R.J., see Early, R.J. 310(1984)1
Chubb, C. and Nolan, C.
Quantitative metabolic profiling of testicular steroid secretions with bonded-phase capillary GC 308(1984)11
Chulay, J.D., see Brown, N.D. 309(1984)426
Chung, H., see Anders, J.C. 311(1984)117
Churáček, J., see Horna, A. 325(1985)367
Chvalovský, V., see Nondek, L. 312(1984)303
Cillard, J., see Torel, J. 323(1985)447
Cillard, P., see Torel, J. 323(1985)447
Čižmárová, E., see Madajová, V. 320(1985)131
Claeys, M., see Vandewoude, M. 311(1984)176
Clark, B.J., see Fell, A.F. 316(1984)423
Clark, R.A., see Nieman, R.H. 317(1984)271
Clark, R.J., see Denham, E.L. 301(1984)253
Clarke, A.E., see Bacic, A. 315(1984)373
Clarke, A.F., see Lagerström, P.-O. 307(1984)230
Clarke, M.L., see Blake, J.D. 312(1984)211
Claus-Walker, J., see Rodriguez, G.P. 308(1984)65
Clezardin, P., McGregor, J.L., Manach, M., Boukerche, H. and Dechavanne, M.
One-step procedure for the rapid isolation of mouse monoclonal antibodies and their antigen binding fragments by fast protein LC on a Mono Q anion-exchange column 319(1985)67
Clift, R.G., see Morreal, C.E. 319(1985)153
Cline, R.E., Yert, L.W. and Needham, L.L.
Determination of germicidal phenols in blood by capillary column GC 307(1984)420
Cloppet, H., see Benbouzid, A. 306(1984)117
Coas, W., see Lepri, L. 322(1985)363
Coates, P.E., see Yip, M.S. 307(1984)343
Cobelli, L., see Marcucci, F. 309(1984)132
Cochetière-Collinet, M.-F. de la, see De la Cochetière-Collinet, M.-F. 305(1984)178
Cociglio, M., Sauvaire, D. and Alric, R.
LC assay of heptaminol in serum and its oral pharmacokinetics in the dog 307(1984)351
Cockrell, C.S. and Ellis, E.F.
Simple single-step HPLC method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 308(1984)316
Coddens, M.E., see Dhanesar, S.C. 324(1985)415
Coddens, M.E., see Poole, C.F. 302(1984)149
Cody, W.L., Wilkes, B.C. and Hruby, V.J.
Reversed-phase HPLC studies of α-MSH fragments 314(1984)313
Cohen, A.S. and Grushka, E.
Structure-retention relationships: chromatographic behaviour and properties of nucleotide-cation complexes 318(1985)221
Cohen, B.M., see Stoll, A.L. 307(1984)457
Cohen, J.D.
Convenient apparatus for the generation of small amounts of diazomethane 303(1984)193
Cohen, J.L., see Terry, M.D. 311(1984)213
Cohen, K.A., Dolan, J.W. and Grillo, S.A.
Examination of theoretical principles of gradient elution as applied to reversed-phase HPLC separation of phenylthiohydantoin amino acids 316(1984)359
Cohen, S.A., see Yu, T.J. 301(1984)425
Cohn, L.J., Greely, V.J. and Tibbetts, D.L.
Determination of demecarium bromide and related compounds by HPLC 321(1985)401
Coign, M.J., see Pimprikar, G.D. 317(1984)413
Cole, W.J., see Brooks, C.J.W. 315(1984)119
Coleman, M.D., Edwards, G., Mihaly, G.W., Howells, R.E. and Breckenridge, A.M.
HPLC method for the determination of pyrimethamine and its 3-N-oxide metabolite in biological fluids 308(1984)363

Coletti, O., see Cremonesi, P. 308(1984)310
Collins, A.J., see Gordon, S.M. 322(1985)246
Collins, C.A. and Chow, C.K.
Determination of vitamin A and vitamin A acetate by HPLC with fluorescence detection 317(1984)349
Collins, J.M., see Klecker, R.W. Jr. 307(1984)361
Collins, M.A., see Origitano, T.C. 311(1984)17
Colombo, L., see Marcucci, F. 309(1984)132
Colombo, L., see Mussini, E. 305(1984)450
Combaut, G., see Serve, L. 322(1985)257
Congote, L.F.
HPLC separation of serum erythrotropin and erythropoietin 310(1984)396
Coninx, P., see Enger, A. 305(1984)335
Conron, D.W., see Driscoll, J.N. 302(1984)43, 302(1984)269
Contario, J.J., see Chen, T.-M. 314(1984)495
Conway, W.D., see Ito, Y. 301(1984)405
Cook, S.E., Pinkerton, T.C., Sadler, G. and Nelson, P.E.
LC of tomato paste serum with sample preparation by ultrafiltration 325(1985)450
Cooke, M., see Gaffney, M.H. 306(1984)303
Cookman, J.R., see Hartley, R. 306(1984)191
Cookson, D.J., Rix, C.J., Shaw, I.M. and Smith, B.E.
HPLC procedure for direct determination of the compound class composition of diesel and kerosine fuels using refractive index detection 312(1984)237
Cooper, H.A. and Hurtubise, R.J.
Mobile phase optimization for the reversed-phase LC of complex hydroxyl aromatic mixtures 324(1985)1
Cooper, W.T. and Hayes, J.M.
Characterization of adsorption at kaolinite surfaces by heterogeneous GSC 314(1984)111
Cooper,C.S., see Toren, P.C. 305(1984)171
Cooperman, B.S., see Hasan, T. 321(1985)462
Cooperman, B.S., see Kerlavage, A.R. 317(1984)201
Corbett, B.J., see Albro, P.W. 312(1984)165
Corcia, A. di, see Andreolini, F. 310(1984)208
Corcia, A. di, see Borra, C. 311(1984)9
Corradini, D. and Horváth, C.
Chromatography of Astra Blue dyes with eluents containing potassium thiocyanate in organic solvents 319(1985)400
Corradini, D., see Melander, W.R. 317(1984)67
Cosmi, G., see Borra, C. 311(1984)9
Cossu, G., Manca, M., Pirastru, M.G., Bullita, R., Bianchi Bosisio, A. and Righetti, P.G.
Determination of glycosylated haemoglobin by isoelectric focusing in non-linear pH gradients 307(1984)103
Costanzo, S.J.
Selection of mixed ion-pair modifiers for HPLC mobile phases 314(1984)402
Cotts, P.M., see Mumby, S.J. 319(1985)241
Couche, D.J., see Nicholson, B.C. 325(1985)221
Couder, M., see Blanche, F. 323(1985)451
Čoupek, J., see Mikeš, O. 301(1984)93
Cousse, H., see Bonnaud, B. 318(1985)398
Covey, T.R., see Skrabalak, D.S. 315(1984)359
Covi, G., see Peuchant, E. 310(1984)297
Cowie, C.E., see Haddad, P.R. 303(1984)321
Cox, J., see Ray, G. 317(1984)455
Cox, J.W. and Pullen, R.H.
Isomodal column switching HPLC technique for the analysis of ciglitazone and its metabolites in human serum 307(1984)155
Crabbe, M.J.C., see Petchey, M. 307(1984)180
Cremonesi, P., Galimberti, G., Coletti, O. and Sportoletti, G.
HPLC determination of imidazole in biological fluids 308(1984)310
Crétin, C., Vidal, J., Suzuki, A. and Gadal, P.
Isolation of plant phosphoenolpyruvate carboxylase by high-performance size-exclusion chromatography 315(1984)430
Creusevau, R., see Airaudo, C.B. 314(1984)349
Crews, C., see Gilbert, J. 319(1985)376
Crolla, T., Santini, F., Visconti, M. and Pifferi, G.
HPLC separation of cadralazine from its potential metabolites and degradation products. Quantitation of the drug in human plasma and urine 310(1984)139
Crommen, J., see Bontemps, J. 307(1984)283
Croo, F. de, see De Croo, F. 325(1985)395
Crook, M., see Ray, G. 317(1984)455
Crooks, P.A., see Cundy, K.C. 306(1984)291
Crosby, S.D. and Dale, G.L.
Simplified procedure for the quantitation of radioactive phosphoinositides by TLC 323(1985)462
Crouch, R.L., Hawley-Fedder, R.A., Parsons, M.L. and Karasek, F.W.
Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53
Crouzat-Reynes, G., see Maingault, P. 314(1984)445
Crowe, J.H., see Womersley, C. 318(1985)112
Cummings, J., Stuart, J.F.B. and Calman, K.C.
Determination of adriamycin, adriamycinol and their 7-deoxyaglycones in human serum by HPLC 311(1984)125
Cundy, K.C. and Crooks, P.A.
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291
Cunningham, F.M., see Mallet, A.I. 309(1984)160
Curran, N.M., see Lovering, E.G. 319(1985)235
Cutsem, J.-L. van, see Bontemps, J. 311(1984)59

D

Dąbrowski, R., see Witkiewicz, Z. 315(1984)145
Dadgar, D., see Hojabri, H. 311(1984)189
Daftsios, A.C., Johnson, E.L., Keeley, F.J., Gryczko, C. and Rawski, V.
HPLC analysis of isoxicam in human plasma and urine 305(1984)145
D'Agostino, G., Mitchell, F., Castagnetta, L. and O'Hare, M.J.
Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13
Dale, G.L., see Crosby, S.D. 323(1985)462
Dalene, M., Lundh, T. and Mathiasson, L.
N-Permethylation of polyamines at trace levels for GC analysis 322(1985)169
Dalene, M., see Skarping, G. 303(1984)89
Dalgaard, L. and Brimer, L.
Electrochemical detection of cyanogenic glycosides after enzymatic post-column cleavage 303(1984)67
Dalgaard, L., see Brimer, L. 303(1984)77
Dalgaard, L., see Køppen, B. 321(1985)385
Dalgaard, L., see Nordholm, L. 305(1984)391
D'Amato, G., see Castello, G. 324(1985)363
Damo, C.P., see Allgire, J.F. 325(1985)249
Damodaran, C., see Annapoorani, K.S. 303(1984)296
Dandrifosse, G., see Bontemps, J. 307(1984)283, 311(1984)59
Daniel, R., see Mallet, A.I. 309(1984)160
Danse, I.H.R., see Klett, R.P. 310(1984)361
Daoud, N., see Arvidsson, T. 317(1984)213
Darvas, F., see Valkó, K. 301(1984)355
Dätwyler, P., see Hirter, P. 323(1985)89
Davies, D.S., see Causon, R.C. 306(1984)257
Davies, I.A.I., see Shamoon, J.M. 314(1984)499
Davies, N.W.
Optimizing flow-rates in capillary GC-MS 325(1985)23
Davies, N.W., Veronese, M.E. and McLean, S.
MS determination of N-hydroxyphenacetin in urine using multiple metastable peak monitoring following TLC 310(1984)179
Davies, T.J. and Hayward, N.J.
Volatile products from acetylcholine as markers in the rapid urine test using head-space GLC 307(1984)11
Davis, A.T., Ingalls, S.T. and Hoppel, C.L.
Determination of free trimethyllysine in plasma and tissue specimens by HPLC 306(1984)79
Davis, H.M., see Wisneski, H.H. 317(1984)421
Davis, T.P., see Berens, M.E. 307(1984)251
Day, B.P.F. and Gregory, J.F. III
Determination of folylpoly-γ-glutamate carboxypeptidase (folacin conjugase) activity using reversed-phase HPLC 318(1985)387
De Beer, J.O., Jacobs, G.A. and Martens, M.A.
Simple reversed-phase HPLC determination of antipyrine in rabbit plasma for pharmacokinetic studies 307(1984)475
De Brauw, M.C. ten Noever, see Tas, A.C. 310(1984)243
De Bree, H., see Ruijten, H.M. 314(1984)183
De Bruijn, C.H.M.M., Reijenga, J.C., Aben, G.V.A., Verheggen, T.P.E.M. and Everaerts, F.M.
Isotachophoresis of allergenic extracts 320(1985)205
De Bruijn, C.H.M.M., see Reijenga, J.C. 320(1985)245
De Croo, F., Van den Bossche, W. and De Moerloose, P.
HPLC behaviour of some pharmaceutically important thiazide, loop and potassium-sparing diuretics 325(1985)395
De Galan, L., see Bartha, Á. 303(1984)29
De Galan, L., see Billiet, H.A.H. 316(1984)231
De Jong, J., Schouten, J.P., Muusze, R.G. and Tjaden, U.R.
Analysis of β-carbolines by reversed-phase ion-pair partition chromatography with fluorometric detection 319(1985)23
De Jong, J., Van Valkenburg, C.F.M. and Tjaden, U.R.
Selective retention of catecholamines and their derivatives in reversed-phase ion-pair partition chromatography 322(1985)43
De la Cochetière-Collinet, M.-F. and Larsson, L.
Studies on volatile metabolites of some potentially pathogenic *Bacillus* species, using automated head-space GC 305(1984)178
De la Teja, E., see Dulitzky, M. 317(1984)403
De Leeuw, I., see Vandewoude, M. 311(1984)176
De Ligny, C.L., Spanjer, M.C., Van Houwelingen, J.C and Weesie, H.M.
Three-mode factor analysis of data on retention in normal-phase HPLC 301(1984)311
De Ligny, C.L., see Brandts, P.M. 322(1985)399
De Moerloose, P., see De Croo, F. 325(1985)395
De Moerloose, P., see De Spiegeleer, B. 315(1984)481
De Moerloose, P., see Van den Eeckhout, E. 318(1985)343
De Ponte, G., see Marcucci, F. 309(1984)132
De Ponte, G., see Mussini, E. 305(1984)450
De Pourcq, P., Hoebus, J., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Quantitative determination of amoxicillin and its decomposition products by HPLC 321(1985)441
De Pourcq, P., see Roets, E. 303(1984)117
De Raat, W.K. and Van Ardenne, R.A.M.
Sorption of organic compounds from urine in mutagenicity testing: choice of sorbent 310(1984)41
De Spiegeleer, B., Slegers, G., Van den Bossche, W. and De Moerloose, P.
Quantitative analysis of *cis*-

dichlorodiammineplatinum(II) by high-performance TLC 315(1984)481
De Vries, J.X., Staiger, C., Wang, N.S., Schlicht, F. and Koroknay, B.
Simultaneous determination of antipyrine and metabolites in human plasma and urine by HPLC 309(1984)219
Debackere, M., see Delbeke, F.T. 325(1985)304
Dechavanne, M., see Clezardin, P. 319(1985)67
Deelder, R.S., see Jansen, H. 325(1985)255
Degen, C., see Castagnola, M. 307(1984)91
Degude, C., see Bauer, M. 315(1984)457
Dehaas Vermeulen, J. and Thompson, T.A.
Assay for the determination of the tetracyclic antidepressant compound aptazapine in plasma by HPLC 306(1984)412
Deiana, S., Dessi, A., Micera, G., Gessa, C. and Solinas, V.
Determination of vanadate(V) by conductometric anion chromatography 320(1985)450
Del Rio, R.M., see Herranz, A.S. 309(1984)139
Delbeke, F.T. and Debackere, M.
Quantitation of caffeine in doping analysis using an automated extractor/concentrator 325(1985)304
Delcour, J.A., see Vercruysse, S.A.R. 324(1985)495
Dell, D., Joly, R., Meister, W., Arnold, W., Partos, C. and Guldimann, B.
Determination of tenoxicam, and the isolation, identification and determination of Ro 17-6661, its major metabolite, in human urine 317(1984)483
Dellacherie, E., see Mathis, R. 319(1985)43
Deltour, G., see Enger, A. 305(1984)335
Demaster, E.G., Shirota, F.N., Redfern, B., Goon, D.J.W. and Nagasawa, H.T.
Analysis of hepatic reduced glutathione, cysteine and homocysteine by cation-exchange HPLC with electrochemical detection 308(1984)83
Deming, S.N., see Wu, C.-G. 302(1984)79
Deml, M., Foret, F. and Boček, P.
Electric sample splitter for capillary zone electrophoresis 320(1985)159
Deml, M., see Dolník, V. 320(1985)89
Deml, M., see Pospíchal, J. 320(1985)139
Den Hartigh, J., Van Oort, W.J. and Hulshoff, A.
LC determination of mitomycin C in human plasma and urine 306(1984)444
Denham, E.L. and Clark, R.J.
GC studies on group VIB organometallic complexes. The analytical and preparative separation of chromium, molybdenum and tungsten trifluorophosphine carbonyl complexes 301(1984)253
Desai, H.K., Joshi, B.S., Panu, A.M. and Pelletier, S.W.
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223
Desbene, P.-L., see Azoulay, M. 303(1984)272
Desideri, P.G., Lepri, L., Heimler, D., Checchini, L. and Giannessi, S.
Fingerprinting of crude oil spills 322(1985)107
Desideri, P.G., see Lepri, L. 322(1985)363
Desimoni, E., see Palmisano, F. 306(1984)205
Desjardins, R., see Causon, R.C. 306(1984)257
Desrivieres, D., see Alton, K.B. 311(1984)319
Dessi, A., see Deiana, S. 320(1985)450
Deves, C., see Padovani, P. 308(1984)229
DeVilbiss, E.D., see Leonhardt, B.A. 322(1985)484
Devine, P.L. and Milborrow, B.V.
Use of Sep-Pak cartridges for the recovery of compounds from scintillation fluids 325(1985)323
DeZwaan, J., see Plaisted, S.M. 314(1984)369
Dhanesar, S.C., Coddens, M.E. and Poole, C.F.
Influence of phase loading on the performance of whisker-walled open tubular columns coated with organic molten salts 324(1985)415
Di Corcia, A., see Andreolini, F. 310(1984)208
Di Corcia, A., see Borra, C. 311(1984)9
Diamondstone, B.I., Wise, S.A. and Sander, L.C.
Carbon analysis for the study of bonded phases in LC 321(1985)319
Dibbelt, L. and Kuss, E.
Isocratic reversed-phase HPLC of free and sulphoconjugated steroids 322(1985)371
Dickinson, A.G., see Pierotti, J.A. 306(1984)377
Diegelmann, R.F., see Lindblad, W.J. 315(1984)447
DiFeo, T.J., see Sisco, W.R. 322(1985)380
Dihuidi, K., Kucharski, M.J., Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413
Dilks, C.H. Jr., Yau, W.W. and Kirkland, J.J.
Sampling and flow effects in sedimentation field flow fractionation 315(1984)45
Dilli, S., Maitra, A.M. and Patsalides, E.
Volatility and GC of some derivatives of Schiff base chelates 318(1985)350
Dilli, S., Miller, S. and Robards, K.
Improved elution of metal β-diketonates on fused-silica wall-coated open-tubular columns 324(1985)75
Dilli, S. and Robards, K.
Spectroscopic properties and GC behaviour of aryl β-diketones and selected chelates 312(1984)109
Dilli, S. and Robards, K.
Volatility studies of metal chelates. VI. GC behaviour of aryl β-diketones and their metal chelates 322(1985)1
Dilli, S. and Robards, K.
Volatility studies of metal chelates. VII. GC behaviour of chelates derived from the aromatic β-diketones 1-phenylbutane-1,3-dione and 4,4,4-trifluoro-1-(pentafluorophenyl)butane-1,3-dione 322(1985)23
Dimauro, P.R., see Hubball, J.A. 302(1984)341
Dingler, E., see Backhaus, J. 307(1984)190

Diquet, B., Doare, L. and Gaudel, G.
New method for the determination of yohimbine in biological fluids by HPLC with amperometric detection 311(1984)449
Diquet, B., Nguyen-Huu, J.J. and Boutron, H.
Determination of pindolol in human plasma by HPLC with amperometric detection 311(1984)430
Dirks, R.J.M., see Oosterbaan, M.J.M. 306(1984)323
Dixon, J.S., Lowe, J.R. and Galloway, D.B.
Rapid method for the determination of either piroxicam or tenoxicam in plasma using HPLC 310(1984)455
Dixon, L.M., see McCormick, M.J. 322(1985)478
Dizdaroglu, M.
Separation of sequence isomeric dipeptides by high-resolution GC 318(1985)384
Dizdaroglu, M., see Margolis, S.A. 322(1985)117
Dmitriev, L.B., see Zamureenko, V.A. 303(1984)109
Doare, L., see Diquet, B. 311(1984)449
Dobberstein, P., see Levsen, K. 323(1985)135
Dobrinska, M.R., see Musson, D.G. 308(1984)251
Doctor, B.P., see Brown, N.D. 315(1984)389
Doerr, W.W., see Shiao, S.-Y. 303(1984)13
Dohji, T., Fushimi, M., Kawabe, T., Kamiyama, F., Mori, M., Sugita, N. and Tanizawa, O.
Rapid measurement of oestradiol and oestriol by HPLC after automatic pretreatment 311(1984)249
Dolan, J.W., see Cohen, K.A. 316(1984)359
Dolecka, E., see Oszczapowicz, J. 315(1984)95
Dolník, V., Deml, M. and Boček, P.
Large sample volume preseparation for trace analysis in isotachophoresis 320(1985)89
Domingo-García, M., López-Garzón, F.J., López-Garzón, R. and Moreno-Castilla, C.
GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19
Domon, B., Dorsaz, A.-C. and Hostettmann, K.
HPLC of oleanane saponins 315(1984)441
Dondeyne, P., see Vercruysse, S.A.R. 324(1985)495
Dondi, F. and Remelli, M.
Characterization of extracolumn and concentration-dependent distortion of chromatographic peaks by Edgeworth-Cramér series 315(1984)67
Dong, S., see Benedek, K. 317(1984)227
Donn, K.H., see Wainer, I.W. 306(1984)405
Dorey, R.C., see Narasimhachari, N. 311(1984)424
Dorsaz, A.-C., see Domon, B. 315(1984)441
Dorsey, J.G., Khaledi, M.G., Landy, J.S. and Lin, J.-L.
Gradient elution micellar LC 316(1984)183
Douse, J.M.F.
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137
Doyle, T.D., see Wainer, I.W. 306(1984)405
Doyle, T.D., see Yang, Z.-Y. 324(1985)444
Drayer, D.E., see Katz, E.Z. 305(1984)477
Drevin, I., see Johansson, B.-L. 321(1985)335
Drewes, L.R., see Singh, A.K. 324(1985)163
Dreyer, R.N., see Siegel, S.A. 319(1985)351
Drinkwater, L., see Womersley, C. 318(1985)112
Driscoll, J.N., Conron, D.W. and Ferioli, P.
Comparison of a new electrochemical detector for GC analysis with the electrolytic conductivity detector 302(1984)269
Driscoll, J.N., Conron, D.W., Ferioli, P., Krull, I.S. and Xie, K.-H.
Trace analysis of organic compounds by HPLC with photoionization detection 302(1984)43
Driskell, J.A., see Pierotti, J.A. 306(1984)377
Drost, R.H., Van Ooijen, R.D., Ionescu, T. and Maes, R.A.A.
Determination of morphine in serum and cerebrospinal fluid by GC and selected ion monitoring after reversed-phase column extraction 310(1984)193
Drouen, A.C.J.H., see Billiet, H.A.H. 316(1984)231
Drummer, O.H., Jarrott, B. and Louis, W.J.
Combined GC-MS procedure for the measurement of captopril and sulfur-conjugated metabolites of captopril in plasma and urine 305(1984)83
Du, Y. and Oshima, R.
Analysis of long-chain phenols in the sap of the Burmese lac tree, *Melanorrhoea usitate* by capillary GLC 318(1985)378
Dubois, J.P., see Menge, G. 310(1984)431
Ducret, P., see Kolarovic, L. 314(1984)233
Ducrocq, C., see Maurs, M. 325(1985)444
Duez, P., Vanhaelen, M., Vanhaelen-Fastré, R., Hanocq, M. and Molle, L.
Comparison between HPTLC fluorometry and HPLC for the determination of sennosides A and B in Senna (Cassia spp.) pods and leaves 303(1984)391
Dufka, O., see Horna, A. 325(1985)367
Duggan, D.E., Hooke, K.F. and Ramjit, H.G.
Sensitive GLC procedure for urinary Nτ-methylimidazole acetic acid, an index of histamine turnover 306(1984)69
Dulitzky, M., De la Teja, E. and Lewis, H.F.
Determination of caffeine in tea by HPLC and a modified digestion procedure 317(1984)403
Dunn, B.E., see Torres, A.R. 316(1984)125
Dunn, D.L., see Scott, B.S. 319(1985)419
Dupeyron, J.P., see Guyon, F. 311(1984)160
Duval, M., see Godbillon, J. 309(1984)198
Dziedzic, S.Z., see Ireland, P.A. 325(1985)275

E

Early, R.J., Thompson, J.R., McAllister, T., Fenton, T.W. and Christopherson, R.J. Branched-chain α-keto acid analysis in biological fluids: preparative clean-up by anion-exchange and analysis by capillary GC 310(1984)1

Ebling, W.F., Szefler, S.J. and Jusko, W.J. Analysis of cortisol, methylprednisolone, and methylprednisolone hemisuccinate. Absence of effects of troleandomycin on ester hydrolysis 305(1984)271

Eckerbom, S., see Bergqvist, Y. 306(1984)147

Ecknig, W., see Kleinert, T. 315(1984)75, 315(1984)85

Edberg, S.C., see Torres, A.R. 316(1984)125

Edqvist, L.-E., see Hallin, P. 319(1985)195

Edstein, M. Quantification of antimalarial drugs. I. Simultaneous measurement of sulphadoxine, N_4-acetylsulphadoxine and pyrimethamine in human plasma 305(1984)502

Edstein, M. Quantification of antimalarial drugs. II. Simultaneous measurement of dapsone, monoacetyldapsone and pyrimethamine in human plasma 307(1984)426

Edwards, D.J., see Kapil, R.P. 307(1984)305

Edwards, G., see Coleman, M.D. 308(1984)363

Edwards, G., see Ward, S.A. 305(1984)239

Eeckhout, E. van den, see Van den Eeckhout, E. 318(1985)343

Eek, L. and Ferrer, N. Sensitive determination of nitrite and nitrate by ion-exchange chromatography 322(1985)491

Eglinton, G., see Marriott, P.J. 301(1984)107

Egorin, M.J., see Nakazawa, H. 307(1984)323

Ehrlich, M., see Gehrke, C.W. 301(1984)199

Ehrsson, H., see Eksborg, S. 306(1984)446

Eibs, G. and Schöneshöfer, M. Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Eichholtz, P.C.N., Binkhuyzen, D. and Thieme, R.E. Direct determination of 3-methoxy-4-sulphonyloxyphenylglycol (MHPG sulphate) in urine using GLC 305(1984)438

Eksborg, S., Ehrsson, H. and Lindfors, A. LC determination of mitomycin C in plasma and urine 306(1984)446

El Kholy, A.M., see Brooks, J.B. 309(1984)269

El Rassi, Z., see Chang, J.-P. 319(1985)396

El Tayar, N., Van de Waterbeemd, H. and Testa, B. Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. I. Relationship between capacity factors and the methanol concentration in methanol-water eluents 320(1985)293

El Tayar, N., Van de Waterbeemd, H. and Testa, B. Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. II. Procedure for the determination of a lipophilic index measured by reversed-phase HPLC 320(1985)305

El-Merzabani, M.M., see Yoshioka, M. 309(1984)63

Elferink, F., Van der Vijgh, W.J.F. and Pinedo, H.M. Analysis of antitumour [1,1-bis(aminomethyl)cyclohexane]platinum(II) complexes derived from spiroplatin by HPLC with differential pulse amperometric detection 320(1985)379

Elferink, F., see Van der Vijgh, W.J.F. 310(1984)335

Elias-Baker, B.A., see Jacob, P. III 306(1984)173

Ellis, E.F., see Cockrell, C.S. 308(1984)316

Ellren, O., see Peetre, I.-B. 318(1985)41

Ellwood, P.A., see Walker, R.F. 301(1984)485

Eloy, C., Fardel, G. and Flandrois, J.P. Fast protein LC for the isolation of *Clostridium perfringens* type A α-toxin 321(1985)235

Emanuel, R.L., Williams, G.H. and Giese, R.W. Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Engelen, D.L. van, see Adams, A.K. 303(1984)341

Engelhorn, S.C., see Goheen, S.C. 317(1984)55

Enger, A., Deltour, G., Coninx, P., Jezekova, D., Cattan, A. and Fournaise, R. GLC determination of a new anticancer agent, 1,3,3,5,5-pentakis-(azaridino)-λ^6,2,4,6,3λ^5,5λ^5-thiatriazadiphosphorine-1-oxide, for pharmacokinetic investigations (in French) 305(1984)335

Enomoto, N., see Mihara, S. 324(1985)428

Epstein, P., see Fogel, J. 317(1984)507

Er, S.S., see Hsia, J.C. 303(1984)425

Erbersdobler, H.F., see Büser, W. 303(1984)234

Ericsson, L.H., see Sasagawa, T. 307(1984)29

Ernst, K., see König, W.A. 301(1984)129

Ersser, R.S. and Mitchell, J.D. Automated anion-exchange chromatographic method for the quantitation of lactulose in physiological fluids and processed milks 307(1984)393

Essers, L. Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Eva, C., see Meek, J.L. 317(1984)343

Evans, L.J. Jr., Young, J.D., Hutchison, J.S. and Baker, L.H. HPLC determination of nafazatrom in human plasma using fluorescence detection 305(1984)163

Evans, S.V., see Hayman, A.R. 325(1985)462
Everaerts, F.M., Verheggen, T.P.E.M., Reijenga, J.C., Aben, G.V.A., Gebauer, P. and Boček, P.
Determination of heavy metals by isotachophoresis 320(1985)263
Everaerts, F.M., see De Bruijn, C.H.M.M. 320(1985)205
Everaerts, F.M., see Lemmens, A.A.G. 320(1985)193
Everaerts, F.M., see Reijenga, J.C. 320(1985)67, 320(1985)245
Everaerts, F.M., see Verheggen, T.P.E.M. 320(1985)99
Everett, R., see Tamma, R.V. 322(1985)236
Evershed, R.P., see Marriott, P.J. 301(1984)107

F

Falcy, C., see Guyon, F. 311(1984)160
Fanali, S., Ossicini, L. and Prosperi, T.
Paper electrophoretic study of ion pair formation. XV. Resolution of optically active cobalt(III)-cyclohexanediamine complexes 318(1985)440
Fanelli, R., see Carlucci, G. 311(1984)141
Farbotko, M.J., see Paryjczak, T. 301(1984)270
Fardel, G., see Eloy, C. 321(1985)235
Farmer, P.B., see Bailey, E. 310(1984)199
Fatmi, A.A., see Parekh, N.J. 314(1984)65
Fausnaugh, J.L., Kennedy, L.A. and Regnier, F.E.
Comparison of hydrophobic-interaction and reversed-phase chromatography of proteins 317(1984)141
Febvre, N., see Lecaillon, J.-B. 317(1984)493
Fedorov, P.I., see Bazyl'chick, V.V. 321(1985)240
Feffer, M., see Loche, S. 317(1984)377
Feibush, B., see Miller, N.T. 316(1984)519
Fekete, J., see Niegreisz, Z. 316(1984)451
Feldhoff, R.C., Ledden, D.J., Steffen, M.C., Steffen, J.M. and Musacchia, X.J.
Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267
Felici, M., Artemi, F., Luna, M. and Speranza, M.
Determination of laccase activity with various aromatic substrates by HPLC 320(1985)435
Felix, G. and Bertrand, C.
Separation of polyaromatic hydrocarbons on caffeine-bonded silica gel 319(1985)432
Fell, A.F., Clark, B.J. and Scott, H.P.
Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423
Fellows, M.E., see Allen, K.G. 309(1984)33
Fels, J.-P., Lechat, P., Rispe, R. and Cautreels, W.
Determination of glaucine in plasma and urine by HPLC 308(1984)273
Felt, V., see Hušek, P. 305(1984)442
Fennessey, P.V., see Pike, A.W. 306(1984)39
Fenton, T.W., see Early, R.J. 310(1984)1
Ferguson, S.M., see Tam, Y.K. 310(1984)438
Ferioli, P., see Driscoll, J.N. 302(1984)43, 302(1984)269
Fernández-Sánchez, E., Carcía-Domínguez, J.A., Menendez, V. and Pertierra-Rimada, E.
Indirect determination of specific retention volumes in GC. Use of retentions of the McReynolds probes relative to *n*-decane 312(1984)69
Ferrer, N., see Eek, L. 322(1985)491
Ferreres, F., see Barberán, F.A.T. 315(1984)101
Ferri, S., see Martelli, P. 301(1984)297
Fetzer, J.C. and Biggs, W.R.
Solvated structure-retention relationships of peropyrene-type polycyclic aromatic hydrocarbons 322(1985)275
Fidler, V., Vacík, J. and Fidler, Z.
Dynamics of isotachophoretic separation. I. Computer simulation 320(1985)167
Fidler, V., see Fidler, Z. 320(1985)175
Fidler, V., see Hanibalová, D. 320(1985)185
Fidler, Z., Fidler, V. and Vacík, J.
Dynamics of isotachophoretic separation. II. Impurities in the electrolytes and sample injection 320(1985)175
Fidler, Z., see Fidler, V. 320(1985)167
Fiedler, H.-P.
Screening for new microbial products by HPLC using a photodiode array detector 316(1984)487
Figueira, M.A. and Ribeiro, J.A.
Separation of adenosine and its derivatives by TLC on silica gel 325(1985)317
Finklestein, S.P., see Stoll, A.L. 307(1984)457
Fisher, D., see Nazareth, A. 319(1985)382
Flach, S.M., see Weiser, E.L. 303(1984)1
Flammini, D., see Forchetti, C. 309(1984)177
Flanagan, R.J. and Jane, I.
HPLC analysis of basic drugs on silica columns using non-aqueous ionic eluents. I. Factors influencing retention, peak shape and detector response 323(1985)173
Flanagan, R.J., see Bhamra, R.K. 307(1984)439
Flanagan, R.J., see Jane, I. 323(1985)191
Flandrois, J.P., see Eloy, C. 321(1985)235
Flodberg, G., see Roeraade, J. 301(1984)454, 322(1985)55
Flora, K., see Ray, G. 317(1984)455
Fogel, J., Epstein, P. and Chen, P.
Simultaneous HPLC assay of acetylsalicylic acid and salicylic acid in film-coated aspirin tablets 317(1984)507
Fong, K.L., see Chang, C.A. 312(1984)99

Forchetti, C., Flammini, D., Carlucci, G., Cavicchio, G., Vaggi, L. and Bologna, M.
HPLC procedure for the quantitation of norfloxacin in urine, serum and tissues 309(1984)177
Foret, F., see Boček, P. 313(1984)189
Foret, F., see Deml, M. 320(1985)159
Forget, P.-P., see Bontemps, J. 311(1984)59
Forsman, U.
Enantiomeric resolution of an optically active guanine derivative by HPLC with phenylalanine-Cu(II) in the mobile phase 303(1984)217
Först, C., see Liebich, H.M. 309(1984)225
Forti, G. Cantelli, see Guerra, M.C. 320(1985)281
Foucault, A. and Rosset, R.
Ligand-exchange chromatography on copper(II)-modified silica gel. Improvements and use for screening of protein hydrolyzate and quantitation of dipeptides and amino acid fractions 317(1984)41
Fournaise, R., see Enger, A. 305(1984)335
Fourtillan, J.-B., Girault, J., Bouquet, S. and Lefèbvre, M.-A.
Determination of mequitazin in human plasma and urine by capillary column GLC-MS 309(1984)391
Frahn, J.L.
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167
Frahn, J.L.
Detection of volatile fatty and dicarboxylic acids on alkaline pherograms with Mills' reagent 318(1985)446
Franchi, G.G., see Martelli, P. 301(1984)297
Francis, G.W., see Andersen, Ø.M. 318(1985)450
Frank, H., Thiel, D. and Langer, K.
Determination of N-acetyl-L-cysteine in biological fluids 309(1984)261
Frank, H., see Bayer, E. 320(1985)393
Frappier, F., see Azoulay, M. 303(1984)272
Freeman, C.J., see Musial, S.P. 319(1985)467
Freeman, D.H., see Hoering, T.C. 316(1984)333
Freeston, L.K., see Gordon, S.M. 322(1985)246
Frei, R.W., see Apffel, J.A. 312(1984)153
Frei, R.W., see Goewie, C.E. 301(1984)325
Frei, R.W., see Gooijer, C. 312(1984)337
Frei, R.W., see Jansen, H. 325(1985)255
Frei, R.W., see Kok, W.T. 315(1984)271
Frei, R.W., see Maris, F.A. 323(1985)113
Frei, R.W., see Nielen, M.W.F. 317(1984)557
Frenz, J., see Jacobson, J. 316(1984)53
Fričová, V., see Kramlová, M. 303(1984)422
Friedel, R.O., see Narasimhachari, N. 311(1984)424
Friedlander, E.R., see Powell, S.C. 317(1984)87
Fries, R., see Schaber, P.M. 316(1984)25
Frigerio, A., see Marcucci, F. 309(1984)132
Fritig, B., see Geoffroy, P. 315(1984)333
Fritz, J.S., see Barron, R.E. 316(1984)201
Frye, J.S., see Rudzinski, W.E. 323(1985)281
Fu, P.P., see Yang, S.K. 316(1984)569
Fuchs, A., see Wiehle, R.D. 307(1984)39
Fujii, K., see Hirokawa, T. 305(1984)429
Fujita, J., see Nakajima,K. 301(1984)241
Fujita, K., see Ito, S. 311(1984)154
Fujita, M., see Nakajima,K. 301(1984)241
Fukunaga, K., see Shoji, S. 319(1985)359
Fukushige, S., see Kato, Y. 320(1985)440
Fullerton, N.F., see Beland, F.A. 308(1984)121
Funakoshi, T., see Shoji, S. 319(1985)359
Funazo, K., Wu, H.-L., Morita, K., Tanaka, M. and Shono, T.
Butylation of inorganic anions for simultaneous determination by GC 319(1985)143
Fung, H.-L., see Morrison, R.A. 308(1984)153
Furlani, G., see Bruner, F. 302(1984)167
Furr, H.C., Amédée-Manesme, O. and Olson, J.A.
Gradient reversed-phase HPLC separation of naturally occurring retinoids 309(1984)299
Fushimi, M., see Dohji, T. 311(1984)249

G

Gabutti, V., see Leone, L. 321(1985)407
Gadal, P., see Crétin, C. 315(1984)430
Gaffney, M.H., Cooke, M. and Simpson, R.
Improved method for the determination of chlorhexidine in urine 306(1984)303
Gagliardi, L., Amato, A., Basili, A., Cavazzutti, G., Gattavecchia, E. and Tonelli, D.
Determination of preservatives in cosmetic products by reversed-phase HPLC 315(1984)465
Gagliardi, L., Amato, A., Basili, A., Cavazzutti, G., Gattavecchia, E. and Tonelli, D.
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353
Gaillot, J., see Ali, H.M. 305(1984)465
Gal, J.
Resolution of the enantiomers of ephedrine, norephedrine and pseudoephedrine by HPLC 307(1984)220
Gal, J. and Sedman, A.J.
R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275
Gal, J., see Miller, K.J. 307(1984)335
Gal, J., see Sedman, A.J. 306(1984)155
Galan, L. de, see Bartha, Á. 303(1984)29
Galan, L. de, see Billiet, H.A.H. 316(1984)231

Galante, L.J., see Ng, K.-T. 309(1984)125
Galimberti, G., see Cremonesi, P. 308(1984)310
Gallagher, R.T. and Hawkes, A.D.
HPLC with stop-flow UV spectral characterization of lolitrem neurotoxins from perennial ryegrass 322(1985)159
Gallagher, R.T., Hawkes, A.D. and Stewart, J.M.
Rapid determination of the neurotoxin lolitrem B in perennial ryegrass by HPLC with fluorescence detection 321(1985)217
Gallelli, J.F., see Narang, P.K. 305(1984)135
Galloway, D.B., see Dixon, J.S. 310(1984)455
Galzy, G., see Krishnamurti, C.R. 315(1984)321
Gama-Sosa, M.A., see Gehrke, C.W. 301(1984)199
Gamerith, G.
Influence of trifluoroacetic anhydride concentration during derivatization and chromatography of N(O,S)-trifluoroacetyl amino acid *n*-propyl esters 318(1985)65
Games, D.E. and Ramsey, E.D.
HPLC-MS of derivatized and underivatized amino acids 323(1985)67
Ganten, D., see Hermann, K. 312(1984)273
Garaj, J., see Krupčík, J. 312(1984)1
Garaj, J., see Matisová, E. 303(1984)151
García-Raso, A., Saura-Calixto, F. and Raso, M.A.
Study of GC behaviour of alkenes based on molecular orbital calculations 302(1984)107
García-Raso, A., see Saura-Calixto, F. 322(1985)35
García-Raso, J., see Saura-Calixto, F. 322(1985)35
Gardner, J.D., see Knight, M. 301(1984)277
Gardner, J.J.
Determination of sodium cromoglycate in human urine by HPLC on an anion-exchange column 305(1984)228
Gargan, P.E., see Otterburn, M.S. 303(1984)429
Gärtner, W., see Ostrowksi, J. 309(1984)115
Gattavecchia, E., see Gagliardi, L. 315(1984)465, 325(1985)353
Gaudel, G., see Diquet, B. 311(1984)449
Gayte-Sorbier, A., see Airaudo, C.B. 314(1984)349
Gazda, K. and Makuch, B.
Selection of optimum conditions for the preparative-scale separation of a dipeptide diastereomer mixture on the basis of measurements made in an analytical column 320(1985)397
Gazdag, M., Szepesi, G. and Hernyes, M.
Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267
Gazzotti, G., Sonnini, S. and Ghidoni, R.
Separation of ganglioside molecular species, with homogeneous long-chain base composition, by reversed-phase TLC 315(1984)395
Gebauer, P. and Boček, P.
Optimization in isotachophoresis. The concept of selectivity and separation speed 320(1985)49
Gebauer, P., see Everaerts, F.M. 320(1985)263
Geerdink, R.B., see Maris, F.A. 323(1985)113
Gehrke, C.W., McCune, R.A., Gama-Sosa, M.A., Ehrlich, M. and Kuo, K.C.
Quantitative reversed-phase HPLC of major and modified nucleosides in DNA 301(1984)199
Gelpí, E., see Tusell, J.M. 306(1984)338
Gelsema, W.J., see Brandts, P.M. 322(1985)399
Genieser, H.-G., see Lüllmann, C. 323(1985)273
Gennaro, M.C., see Mentasti, E. 322(1985)177
Geoffroy, P., Legrand, M., Hermann, C. and Fritig, B.
HPLC of proteins. Purification of plant enzymes by ion-exchange chromatography 315(1984)333
George, S., see McBurney, A. 308(1984)387
George, W.J., see Rege, A.B. 309(1984)397
Georges, Y., see Azoulay, M. 303(1984)272
Gerba, S., see Abegaz, B. 324(1985)440
Gerber, N., see Mays, D.C. 311(1984)301
Geschke, R., see Heizmann, P. 310(1984)129, 311(1984)411
Gessa, C., see Deiana, S. 320(1985)450
Ghe, A.M., see Caboni, M.F. 315(1984)223
Ghidoni, R., see Gazzotti, G. 315(1984)395
Ghosal, P.K., see Ray, B. 315(1984)401
Ghosh, A.K., see Keller, S. 305(1984)461, 305(1984)461
Giacherio, D., see Annesley, T. 305(1984)199
Giacomini, K.M., see Smith, G.A. 307(1984)410
Giannessi, S., see Desideri, P.G. 322(1985)107
Gibson, E.K. Jr., see Andrawes, F. 302(1984)181
Giese, R.W., see Emanuel, R.L. 312(1984)285
Giese, R.W., see Mohamed, G.B. 314(1984)211
Giese, R.W., see Nazareth, A. 309(1984)357, 314(1984)201, 314(1984)219, 319(1985)382
Giese, R.W., see Warden, B.A. 314(1984)295
Giessmann, U., see Schmelzeisen-Redeker, G. 323(1985)127
Giger, W., see Schaffner, C. 312(1984)413
Gil-Av, E., see Watabe, K. 318(1985)235
Gilbert, H.S., see Stump, D.D. 306(1984)371
Gilbert, J., Shepherd, M.J. and Wallwork, M.A.
Multi-stage chromatographic procedure for the isolation of a single oligomeric species (vinyl chloride tetramer) from poly(vinyl chloride) resin used for food packaging applications 320(1985)361
Gilbert, J., Startin, J.R. and Crews, C.
Optimisation of conditions for the trimethylsilylation of trichothecene mycotoxins 319(1985)376
Gilbert, S.G., see Stavric, B. 310(1984)107
Gill, J.P., see Marriott, P.J. 301(1984)107
Gill, R., Abbott, R.W. and Moffat, A.C.
HPLC systems for the separation of local anaesthetic drugs with applicability to the analysis of illicit cocaine samples 301(1984)155
Gill, R., see Law, B. 301(1984)165, 325(1985)294
Gilon, C., see Laufer, R. 301(1984)415

Gilvydis, D.M., see Walters, S.M. 317(1984)533
Ginneken, C.A.M. van, see Koopmans, P.P. 307(1984)445
Girault, J., see Fourtillan, J.-B. 309(1984)391
Glajch, J.L., Gluckman, J.C., Charikofsky, J.G., Minor, J.M. and Kirkland, J.J.
Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23
Glennon, J.D., see Hojabri, H. 311(1984)189
Glöckner, G., Van den Berg, J.H.M., Meijerink, N.L.J., Scholte, T.G. and Koningsveld, R.
Two-dimensional analysis of copolymers by size-exclusion chromatography and gradient-elution reversed-phase precipitation chromatography 317(1984)615
Głód, B. and Kemula, W.
Electrokinetic detection of lauric, myristic, stearic and arachidic acids separated by reversed-phase HPLC 321(1985)433
Gluckman, J., Shelly, D. and Novotný, M.
Laser fluorimetry for capillary column LC: high-sensitivity detection of derivatized biological compounds 317(1984)443
Gluckman, J.C. and Novotny, M.
New thermionic detector utilizing orthogonal nebulization for capillary column LC 314(1984)103
Gluckman, J.C., see Glajch, J.L. 318(1985)23
Godbillon, J. and Duval, M.
Determination of two metoprolol metabolites in human urine by HPLC 309(1984)198
Godbillon, J., see Lecaillon, J.-B. 305(1984)411
Godeau, R.P., see Paphassarang, S. 319(1985)412
Goebel, H., see Stan, H.-J. 314(1984)413
Goetz, M., see Joshua, H. 303(1984)185
Goewie, C.E., Nielen, M.W.F., Frei, R.W. and Brinkman, U.A.T.
Optimization of precolumn design in LC 301(1984)325
Gogitidze, N.M., see Hradil, J. 324(1985)277
Goheen, S.C. and Engelhorn, S.C.
Hydrophobic interaction HPLC of proteins 317(1984)55
Goldberg, A.P., Nowakowska, E., Antle, P.E. and Snyder, L.R.
Retention-optimization strategy for the HPLC ion-pair separation of samples containing basic compounds 316(1984)241
Goldberg, A.P., see Antle, P.E. 321(1985)1
Goldberg, P.A., see Walker, R.F. 301(1984)485
Goldstein, D.S., Stull, R., Markey, S.P., Marks, E.S. and Keiser, H.R.
Dihydrocaffeic acid: a common contaminant in the LC-electrochemical measurement of plasma catecholamines in man 311(1984)148
Gonnet, C., see Boulieu, R. 307(1984)469
Gonnet, C., see Nicot, G. 306(1984)279
Gooderham, N.J. and Gorrod, J.W.
Application of reversed-phase HPLC to in vitro drug metabolism studies with N-alkylarylamines 309(1984)339
Gooijer, C., Veltkamp, A.C., Baumann, R.A., Velthorst, N.H., Frei, R.W. and Van der Vijgh, W.J.F.
Analysis of platinum complexes by LC with quenched phosphorescence detection 312(1984)337
Goon, D.J.W., see Demaster, E.G. 308(1984)83
Goormachtig, P.F.J., see Beem, E.P. 323(1985)439
Gordon, S.M., Freeston, L.K., Collins, A.J. and Bain, R.
Determination of emepronium bromide in tablets by reversed-phase HPLC 322(1985)246
Gorrod, J.W., see Gooderham, N.J. 309(1984)339
Gosset, G., see Lecaillon, J.-B. 305(1984)411
Goto, M., see Ishii, D. 316(1984)441
Gouda, T.A., see Strang, C. 314(1984)482
Gould, B.J., see Amoah, A.G.B. 305(1984)401
Goulle, J.P., see Lacroix, C. 307(1984)137, 309(1984)183
Govers, C.J.L.R., see Barends, D.M. 322(1985)321
Gower, D.B., see Kwan, T.K. 301(1984)189
Grajek, H., see Witkiewicz, Z. 312(1984)141
Grandberg, I.I., see Zamureenko, V.A. 303(1984)109
Granit, L., see Katz, E.Z. 305(1984)477
Grassini-Strazza, G. and Nicoletti, I.
HPTLC on amino-bonded silica gel: application to barbiturates and steroids 322(1985)149
Gravallese, D.A., Musson, D.G., Pauliukonis, L.T. and Bayne, W.F.
Determination of imipenem (N-formimidoyl thienamycin) in human plasma and urine by HPLC, comparison with microbiological methodology and stability 310(1984)71
Gray, D.O., see Albert, L.S.R. 312(1984)357
Gray, D.O., see Hayman, A.R. 325(1985)462
Gray, R.R., see Brown, N.D. 315(1984)389
Grayeski, M.L., see Koziol, T. 317(1984)355
Grech-Bélanger, O.
GC method for the routine serum monitoring of mexiletine 309(1984)165
Greef, J. van der, see Van der Greef, J. 323(1985)81
Greely, V., see Biffar, S. 318(1985)404
Greely, V.J., see Cohn, L.J. 321(1985)401
Greenbaum, L.M., see Narayanan, T.K. 306(1984)109
Greenblatt, D.J., see Abernethy, D.R. 307(1984)194
Greenblatt, D.J., see Matlis, R. 310(1984)445
Greenblatt, D.J., see Willis, C.R. 307(1984)200
Greenhut, J. and Rudolph, F.B.
Preparative separation of nucleotides by ion exchange HPLC using a volatile buffer system 319(1985)461
Greenland, R.D., see Lasley, S.M. 305(1984)27

Gregg, M.R. and Jack, D.B.
Determination of timolol in plasma and breast milk using HPLC with electrochemical detection 305(1984)244
Grego, B., see Skinner, S.J.M. 308(1984)111
Gregory, J.F. III, see Day, B.P.F. 318(1985)387
Gregory, P.B., see Smith, G.A. 307(1984)410
Grehlinger, M., see Trumbore, C.N. 322(1985)443
Greibrokk, T., see Lundanes, E. 322(1985)347
Grese, R., see Timmons, L. 314(1984)476
Gribnau, F.W.J., see Koopmans, P.P. 307(1984)445
Grierson, J.R. and Adam, M.J.
In-situ preparation of a chemically bonded chiral stationary phase for the separation of aromatic α-amino acid enantiomers 325(1985)103
Griest, W.H., see Maskarinec, M.P. 302(1984)51
Grillo, S.A., see Cohen, K.A. 316(1984)359
Grimsrud. E.P. and Valkenburg, C.A.
New schemes for the electron-capture sensitization of aromatic hydrocarbons 302(1984)243
Grimsrud, E.P.
Effect of trialkylamines on the response of the electron-capture detector to acridine 312(1984)49
Grinberg, N. and Weinstein, S.
Enantiomeric separation of Dns-amino acids by reversed-phase TLC 303(1984)251
Grinberg, N., see Weinstein, S. 318(1985)117
Grob, K. and Habich, A.
Headspace gas analysis: the role and the design of concentration traps specifically suitable for capillary GC 321(1985)45
Grob, K. Jr.
Band broadening in space in splitless injection 324(1985)251
Grob, K. Jr. and Kuhn, S.
Speed of temperature increase when using large retention gaps in capillary GC 301(1984)1
Grob, K. Jr. and Neukom, H.P.
Method for testing retention gaps and data on the GC retention power of differently deactivated capillary tubes 323(1985)237
Grobecker, H., see Kees, F. 305(1984)363
Groenendijk, G., see Kok, W.T. 315(1984)271
Gros, E.G., see Burton, G. 315(1984)435
Grossi, G., Lippi, A., Battistoni, R., Martelli, E. and Sturani, C.
HPLC determination of mepixanthone in serum 309(1984)214
Grove, J., Alken, R.G. and Schechter, P.J.
Assay of γ-vinyl-γ-aminobutyric acid (4-amino-hex-5-enoic acid) in plasma and urine by automatic amino acid analysis. Application to human pharmacokinetics 306(1984)383
Grünes, R. and Sawodny, W.
Polymeric Schiff base complexes as solid phases in GC 322(1985)63
Grushka, E.
Effect of high speed on peak capacity in LC 316(1984)81
Grushka, E., see Cohen, A.S. 318(1985)221
Gryczko, C., see Daftsios, A.C. 305(1984)145
Guay, D.R.P., see Meatherall, R.C. 307(1984)295
Gübitz, G. and Mihellyes, S.
Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462
Gudéhn, A.
Improved chromatographic procedure for determination of 9-(N-methylaminomethyl)anthracene isocyanate derivatives by HPLC 301(1984)481
Guechot, C. and Nicolle, P.
Purity assay of niflumic acid by reversed-phase HPLC 303(1984)440
Guerra, M.C., Barbaro, A.M., Biagi, G.L., Pietrogrande, M.C., Borea, P.A., Andreani, A. and Cantelli Forti, G.
R_m values, retention times and π values of a series of potentially mutagenic nitroimidazo[2,1-*b*]thiazoles 320(1985)281
Guieze, P. and Williams, J.M.
Determination of the molar mass of petroleum distillation residues using GPC 312(1984)261
Guillemin, C.L., Millet, J.L. and Hamon, E.
Steam GSC: a flexible separation technique 301(1984)11
Guillen, M.D., see Bermejo, J. 318(1985)187
Guilloteau, D., see Maingault, P. 314(1984)445
Guinet, R., see Benbouzid, A. 306(1984)117
Guiochon, G., see Krupčik, J. 312(1984)1
Gulaid, A.A., Houghton, G.W. and Boobis, A.R.
Separation of acebutolol and diacetolol diastereomers by reversed-phase HPLC 318(1985)393
Guldimann, B., see Dell, D. 317(1984)483
Gulyás, H.
Determination of aflatoxins B_1, B_2, G_1, G_2 and M_1 by over-pressurized TLC (in German) 319(1985)105
Gustine, D.L.
Determination of *S*-methyl cysteine sulfoxide in *Brassica* extracts by HPLC 319(1985)450
Gustine, D.L. and McCulloch, J.
Versatile microcomputer-controlled, automated gradient analytical HPLC system 316(1984)407
Gutteridge, C.S., see Aries, R.E. 319(1985)285
Guyon, F., Lecomte-Joulin, V., Falcy, C. and Dupeyron, J.P.
Automated LC of urinary dopamine (in French) 311(1984)160
Gyllenhaal, O. and Hoffmann, K.-J.
Simultaneous determination of metoprolol and metabolites in urine by capillary column GC as oxazolidineone and trimethylsilyl derivatives 309(1984)317
Gyllenhaal, O., see Antonsson, A.-M. 308(1984)181

H

Haag, A., see Schwarz, W. 310(1984)188
Haas, M.J., see Hicks, K.B. 319(1985)159
Haavik, S., Paulsen, B.S. and Wold, J.K.
Purification of a basic glycoprotein allergen from pollen of timothy by HPLC 321(1985)199
Habich, A., see Grob, K. 321(1985)45
Häcker, R., see Punkt, J. 320(1985)105
Haddad, P.R., Alexander, P.W. and Trojanowicz, M.
HPLC of organic acids with potentiometric detection using a metallic copper electrode 315(1984)261
Haddad, P.R., Alexander, P.W. and Trojanowicz, M.
Ion chromatography of inorganic anions with potentiometric detection using a metallic copper electrode 321(1985)363
Haddad, P.R., Alexander, P.W. and Trojanowicz, M.
Application of indirect potentiometric detection with a metallic copper electrode to ion chromatography of transition metal ions 324(1985)319
Haddad, P.R. and Cowie, C.E.
Computer-assisted optimization of eluent concentration and pH in ion chromatography 303(1984)321
Haddad, P.R. and Heckenberg, A.L.
Studies on sample preconcentration in ion chromatography. I. Design of an automated, single pump preconcentration system with direct UV absorbance detection 318(1985)279
Haefelfinger, P. and Hess, B.
Determination of the imidazo quinazoline derivative Ro 13-6438 in biological fluids by HPLC 305(1984)153
Hagedorn, I., see Brown, N.D. 315(1984)389
Hagel, L.
Effect of sample volume on peak width in high-performance gel filtration chromatography 324(1985)422
Hagino, S., see Tanaka, E. 308(1984)393
Haglund, Å.C. and Marsden, N.V.B.
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47
Haglund, Å.C., Marsden, N.V.B. and Östling, S.G.
Partitioning of oligoglucans in Sephadex G-15 in relation to their conformational structure 318(1985)57
Haglund,Å.C. and Marsden, N.V.B.
Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365
Hahn, S., see Skrinska, V. 311(1984)380
Hahnvajanawong, V., see Sattayasai, N. 307(1984)235
Hailey, D.M., see Johnston, M.A. 318(1985)362
Hak, E.W., see Bacic, A. 315(1984)373
Haken, J.K. and Korhonen, I.O.O.
GC of homologous esters. XXVII. Retention increments of C_1-C_{18} primary alkanols and their 2-chloropropanoyl and 3-chloropropanoyl derivatives on SE-30 and OV-351 capillary columns 319(1985)131
Haken, J.K. and Korhonen, I.O.O.
GC of homologous esters. XXVIII. Retention increments of aliphatic C_1-C_{18} *n*-alkyl esters of butanoic acid and its monochloro derivatives on SE-30 and OV-351 capillary columns 320(1985)325
Haken, J.K. and Korhonen, I.O.O.
GC of homologous esters. XXIX. Propanoyl and monochlorpropanoyl esters of lower saturated brached-chain and unsaturated alcohols 324(1985)343
Haken, J.K., Madden, B.G. and Korhonen, I.O.O.
GC of homologous esters. XXXI. Butanoyl and monochlorobutanoyl esters of lower saturated branched chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 325(1985)61
Haken, J.K., see Burford, R.P. 321(1985)295
Haken, J.K., see Chrétien, J.R. 324(1985)355
Haken, J.K., see Vimalasiri, P.A.D.T. 319(1985)121
Haken, J.K., see Wainwright, M.S. 321(1985)287
Haken, J.K., see Zadro, S. 323(1985)305
Haky, J.E. and Stickney, T.M.
Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137
Halbert, M.K. and Baldwin, R.P.
Determination of lidocaine and active metabolites in blood serum by LC with electrochemical detection 306(1984)269
Halket, J.M. and Schulten, H.-R.
Thick-film capillary column GC-field ionization MS. A complementary technique for the rapid analysis of volatiles 322(1985)200
Hallgren, C., see Andersson, K. 312(1984)482
Hallin, P., Madej, A. and Edqvist, L.-E.
Subunits of bovine lutropin. Hormonal and immunological activity after separation with reversed-phase HPLC 319(1985)195
Halloran, B.P., Bikle, D.D. and Whitney, J.O.
Separation of isotopically labeled vitamin D metabolites by HPLC 303(1984)229
Halonen, T., see Herranen, J. 307(1984)241
Halperin, G., Tauber-Finkelstein, M. and Shaltiel, S.
Hydrophobic chromatography of cells: adsorption and resolution on homologous series of alkylagaroses 317(1984)103
Ham, J., see Smith, J.A. 305(1984)295
Hamacher, K., Schmid, G., Sahm, H. and Wandrey, C.
Structural heterogeneity of cellooligomers

homogeneous according to high-resolution size-exclusion chromatography 319(1985)311
Hamon, E., see Guillemin, C.L. 301(1984)11
Hanai, T. and Hubert, J.
Dependence of the retention of phenols upon Van der Waals volume, π-energy and hydrogen-bonding effects 302(1984)89
Hanai, T. and Hubert, J.
Chromatography of aromatic acids on ion exchangers 316(1984)261
Hanibalová, D., Vacík, J. and Fidler, V.
Spectrophotometric detection in capillary isotachophoresis: determination of the composition of mixed zones 320(1985)185
Hanke, F.J., see Kubo, I. 321(1985)246
Hanocq, M., see Duez, P. 303(1984)391
Hansen, D.R. and Hill, H.H. Jr.
Ionization quenching of a hydrogen atmosphere flame as a detection method for GC 303(1984)331
Hansén, L., Sollenberg, J. and Wiberg, K.
Determination of butylamine in air samples by isotachophoresis 312(1984)489
Hantz, A., see Sârbu, C. 322(1985)272
Hara, S. and Ohkuma, T.
Role of mobile phase components for multi-functional solutes in silica gel LSC. Resolution of protected deoxyribonucleotide derivatives for oligomer synthesis 316(1984)389
Hara, S., see Ohkuma, T. 323(1985)227
Harada, K.-I., see Oka, H. 325(1985)265
Hardy, H.L., see Walker, R.F. 301(1984)485
Harris, P.J., see Bacic, A. 315(1984)373
Harris, P.Q., Bacopoulos, N.G. and Brown, S.J.
Measurement of homovanillic acid in human plasma by HPLC with electrochemical detection 309(1984)379
Hart, S.J., Aguilar, M.I., Healey, K., Smail, M.C. and Calder, I.C.
Improved HPLC separation of urinary paracetamol metabolites using radially compressed columns 306(1984)215
Harten, J. van, see Kleinbloesem, C.H. 308(1984)209
Hartigh, J. den, see Den Hartigh, J. 306(1984)444
Hartley, R., Cookman, J.R. and Smith, I.J.
Simultaneous determination of caffeine and its N-demethylated metabolites in umbilical cord plasma using HPLC 306(1984)191
Haruta, M., see Tanaka, A. 306(1984)51
Harvan, D.J., see Albro, P.W. 312(1984)165
Harvan, D.J., see Parker, C.E. 319(1985)273
Harvey, D.J. and Tiffany, J.M.
Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173
Harvey, R.W., see Maskarinec, M.P. 302(1984)51
Harvey, T.G., Matheson, T.W., Pratt, K.C. and Stanborough, M.S.
Determination of carbonyl compounds in an Australian (Rundle) shale oil 319(1985)230
Hasan, T. and Cooperman, B.S.
Reversed-phase HPLC separations of tetracycline derivatives using volatile mobile phases 321(1985)462
Hasegawa, J., see Langendijk, P.N.J. 307(1984)371
Hashimoto, T., see Kato, Y. 318(1985)358, 320(1985)440
Hashimoto, Y., see Yamamoto, S. 303(1984)99
Haškovec, L., see Kysilka, R. 320(1985)414
Hass, J.R., see Parker, C.E. 319(1985)273
Hata, W.Y., see Shiao, S.-Y. 303(1984)13
Hatano, H., see Rokushika, S. 320(1985)335
Hatano, H., see Wada, H. 320(1985)369
Hattori, H., see Ikeda, M. 307(1984)111
Hattori, T., see Yamada, H. 320(1985)403
Hattori, Y., Kobayashi, A., Takemoto, S., Takami, K., Kuge, Y., Sugimae, A. and Nakamoto, M.
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341
Havaši, P. and Kaniansky, D.
Simple UV absorption detector for capillary isotachophoresis 325(1985)137
Havaši, P., see Zelenský, I. 325(1985)161
Haviland, A.M., see Schneider, J.J. 308(1984)359
Hawes, E.M., see Roy, S.D. 310(1984)307
Hawkes, A.D., see Gallagher, R.T. 321(1985)217, 322(1985)159
Hawkins, B.L., see Miller, M.L. 319(1985)9
Hawkins, B.L., see Rudzinski, W.E. 323(1985)281
Hawley-Fedder, R.A., Parsons, M.L. and Karasek, F.W.
Products obtained during combustion of polymers under simulated incinerator conditions. I. Polyethylene 314(1984)263
Hawley-Fedder, R.A., Parsons, M.L. and Karasek, F.W.
Products obtained during combustion of polymers under simulated incinerator conditions. II. Polystyrene 315(1984)201
Hawley-Fedder, R.A., Parsons, M.L. and Karasek, F.W.
Products obtained during combustion of polymers under simulated incinerator conditions. III. Polyvinyl chloride 315(1984)211
Hawley-Fedder, R.A., see Crouch, R.L. 303(1984)53
Hawthorne, S.B., see Hutte, R.S. 302(1984)173
Hayakawa, K. and Tanaka, H.
HPLC determination of bovine and porcine insulins in a commercial injection 312(1984)476
Hayakawa, K., see Miyazaki, M. 323(1985)443
Hayase, K., Shitashima, K. and Tsubota, H.
Application of reversed-phase LC to dissolved organic matter in estuarine seawater 322(1985)358
Hayashi, T., see Konomi, H. 311(1984)375
Hayashi, T., see Todoriki, H. 310(1984)273
Hayashi, T., see Tsuchiya, H. 309(1984)43

Hayes, G.E. and Hillman, D.E.
Determination of 2,6-di-*tert.* -butyl-4-methylphenol in aviation turbine fuel by LC with electrochemical detection 322(1985)376
Hayes, J.M., see Cooper, W.T. 314(1984)111
Hayes, M.J., see Mohamed, G.B. 314(1984)211
Hayes, T.M., see Hsia, J.C. 303(1984)425
Hayman, A.R., Gray, D.O. and Evans, S.V.
New high-performance LC system for the separation of biogenic amines as their Dns derivatives 325(1985)462
Hayward, N.J., see Davies, T.J. 307(1984)11
Hazlett, R.N., see Wright, B.W. 314(1984)253
Head, R.J., see Barone, S. 310(1984)283
Healey, K., see Hart, S.J. 306(1984)215
Heape, A.M., Juguelin, H., Boiron, F. and Cassagne, C.
Improved one-dimensional TLC technique for polar lipids 322(1985)391
Hearn, M.T.W., see Skinner, S.J.M. 308(1984)111
Heath, R.R., see Sonnet, P.E. 321(1985)127
Heckenberg, A.L., see Haddad, P.R. 318(1985)279
Hees, P.A.M. van, see Van Hogezand, R.A. 305(1984)470
Heflich, R.H., see Beland, F.A. 308(1984)121
Hegde, M.V., see Roy, F. 324(1985)489
Heikkinen, L.
Silica capillary GC determination of ibuprofen in serum 307(1984)206
Heimler, D., see Desideri, P.G. 322(1985)107
Hein, C.S., see Marriott, P.J. 301(1984)107
Hein, S.J., see Ramus, T.L. 314(1984)243
Heindze, A.M., see Krishnamurti, C.R. 315(1984)321
Heineman, F.C., see Boppana, V.K. 317(1984)463
Heitz, J.R., see Pimprikar, G.D. 317(1984)413
Heizmann, P. and Geschke, R.
Determination of the antimalarial mefloquine in humans plasma by GC with electron-capture detection 311(1984)411
Heizmann, P., Geschke, R. and Zinapold, K.
Determination of bromazepam in plasma and of its main metabolites in urine by reversed-phase HPLC 310(1984)129
Heizmann, P., Wendt, G., Von Alten, R., Zinapold, K. and Buser, C.
Determination of tiapamil and of its two main metabolites in plasma and in urine by HPLC 310(1984)119
Helland, P., see Tong, H.Y. 324(1985)373
Helm, D. van der, see Jalal, M.A.F. 301(1984)247
Helmlinger, J., see Rausch, T. 318(1985)95
Hemdan, E.S. and Porath, J.
Development of immobilized metal affinity chromatography. I. Comparison of two iminodiacetate gels 323(1985)247
Hemdan, E.S. and Porath, J.
Development of immobilized metal affinity chromatography. II. Interaction of amino acids with immobilized nickel iminodiacetate 323(1985)255
Hemdan, E.S. and Porath, J.
Development of immobilized metal affinity chromatography. III. Interaction of oligopeptides with immobilized nickel iminodiacetate 323(1985)265
Henderson, D.E., see Kirkman, C.M. 317(1984)569
Henion, J.D., see Skrabalak, D.S. 315(1984)359
Henshaw, P.F., see Bovenkamp, J.W. 301(1984)492
Hensler, C.J., see Buchta, R.C. 325(1985)456
Heritage, I.D., see Bond, A.M. 315(1984)313
Hermann, C., see Geoffroy, P. 315(1984)333
Hermann, K., Lang, R.E., Unger, T., Bayer, C. and Ganten, D.
Combined HPLC-radioimmunoassay for the characterization and quantitative measurement of neuropeptides 312(1984)273
Hermansson, J.
Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537
Hermansson, J.
Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379
Hernandez-Gil, N., Wentworth, W.E., Limero, T. and Chen, E.C.M.
Use of electron-capture data at short pulse intervals for study of electron-capture mechanisms 312(1984)31
Hernyes, M., see Gazdag, M. 316(1984)267
Herranen, J., Huhtikangas, A., Tirronen, H., Halonen, T., Huuskonen, M., Reinikainen, K. and Riekkinen, P.
Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241
Herranz, A.S., Lerma, J. and Del Rio, R.M.
Determination of γ-aminobutyric acid in physiological samples by a simple, rapid HPLC method 309(1984)139
Herrmann, K., see Winter, M. 315(1984)243
Hertz, J., see Baltensperger, U. 324(1985)153
Hess, B., see Haefelfinger, P. 305(1984)153
Hess, S.L., see Sloan, D.L. 316(1984)43
Hicks, K.B., Lim, P.C. and Haas, M.J.
Analysis of uronic and aldonic acids, their lactones, and related compounds by HPLC on cation-exchange resins 319(1985)159
Hida, M., see Yoshida, H. 325(1985)179
Hidalgo, I.J. and Muir, K.T.
HPLC method for the determination of labetalol in plasma using UV detection 305(1984)222
Hikino, H., see Kasahara, Y. 324(1985)503
Hilgenberg, W., see Rausch, T. 318(1985)95
Hill, H.H. Jr., see Hansen, D.R. 303(1984)331
Hill, R.A.
Separation of methoxyanilines by

competing-ion reversed-phase HPLC 316(1984)227
Hillebrand, G.G., see Revich, G.G. 317(1984)283
Hillman, D.E., see Hayes, G.E. 322(1985)376
Hine, T., see Kasahara, Y. 324(1985)503
Hines, J.W., see Voyksner, R.D. 323(1985)383
Hinkel, J.-P., see Hornsperger, J.-M. 306(1984)364
Hirama, Y. and Yoshida, H.
Separation of anionic chloro complexes by capillary tube isotachophoresis in non-aqueous systems 322(1985)139
Hirano, H., see Yoza, N. 325(1985)385
Hirata, Y.
Comparison of chromatographic behaviour of silica and chemically bonded phases in supercritical fluid chromatography 315(1984)31
Hirata, Y.
Supercritical fluid chromatography of various samples with different polarities 315(1984)39
Hirokawa, T., Aoki, N. and Kiso, Y.
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11
Hirokawa, T., Kobayashi, S. and Kiso, Y.
Isotachophoretic determination of mobility and pK_a by means of computer simulation. III. Evaluation of mobility and pK_a of fifteen nucleotides and seven phosphorus oxoacids and their isotachophoretic separation 318(1985)195
Hirokawa, T., Takemi, H., Kiso, Y., Takiyama, R., Morio, M., Fujii, K. and Kikuchi, H.
Analytical isotachophoresis utilizing computer simulation. II. Assessment of optimum separation conditions for urinary trifluoroacetic acid metabolized from anaesthetic halothane 305(1984)429
Hirose, S., see Iwamoto, M. 310(1984)151
Hirota, K., Kawase, M. and Kishie, T.
Effect of sodium dodecyl sulphate on the extraction of ubiquinone-10 in the determination of plasma samples 310(1984)204
Hirota, K. and Shimamura, M.
Sepharose derivatives containing citric acid as affinity ligand. Purification of fumarase 319(1985)173
Hirter, P., Walther, H.J. and Dätwyler, P.
Micro column LC-MS using a capillary interface 323(1985)89
Hisayasu, G.H., see Terry, M.D. 311(1984)213
Hiyama, C., see Miwa, H. 321(1985)165
Hjertén, S. and Yang, D.
HPLC separations on dihydroxyboryl-agarose 316(1984)301
Hlaváč, C., see Kovář, J. 319(1985)341
Ho, F.J., see Reeve, J.R. Jr. 321(1985)421
Ho, I.K., see Purser, C. 311(1984)135
Hodges, R.S., see Lau, S.Y.M. 317(1984)129
Hodges, R.S., see Taneja, A.K. 317(1984)1
Hoebus, J., see De Pourcq, P. 321(1985)441
Hoering, T.C. and Freeman, D.H.
Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333
Hoey, L.D., see Slatkavitz, K.J. 302(1984)277
Hofermaas, R., see Josić, D. 317(1984)35
Hoffmann, K.-J., see Gyllenhaal, O. 309(1984)317
Hoffmann, S., Blomberg, L.G., Buijten, J., Markides, K. and Wännman, T.
GC-MS analysis of compounds generated upon thermal degradation of some stationary phases in capillary GC 302(1984)95
Hoffmann, S., see Buijten, J. 301(1984)265
Hoffmann, S., see Markides, K. 302(1984)319
Hofmann, G.E., see Wiehle, R.D. 307(1984)39
Hogezand, R.A. van, see Van Hogezand, R.A. 305(1984)470
Höhn, M., see Van der Greef, J. 323(1985)81
Hoinard, C., see Maingault, P. 314(1984)445
Hojabri, H., Dadgar, D. and Glennon, J.D.
GLC analysis with electron-capture detection of diclofensine in human plasma following derivatization 311(1984)189
Holland, M.L., Rhodes, G.R., Wiesler, D. and Novotny, M.
Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23
Holland, M.L., see Ng, K.-T. 311(1984)418
Holland, O.B., see Risk, M. 317(1984)367
Holler, J.S., see Brooks, J.B. 309(1984)269
Holohan, T.V., see Phillips, T.M. 317(1984)173
Holt, D.W., see Bhamra, R.K. 307(1984)439
Holzer, G., see Andrawes, F. 302(1984)181
Honc, F., see Kochak, G. 310(1984)319
Honda, K., see Miyaguchi, K. 303(1984)173, 316(1984)501
Honda, S., see Kakehi, K. 318(1985)367
Hondo, T., see Hoshino, T. 316(1984)473
Honegger, C.G., Burri, R., Langemann, H. and Kempf, A.
Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53
Honeyman, T.W., see Ryan, P.J. 312(1984)461, 320(1985)421
Hoogewijs, G. and Massart, D.L.
Development of a standardized analysis strategy for basic drugs, using ion-pair extraction and HPLC. VI. Drug level determination in saliva 309(1984)329
Hooghwinkel, G.J.M., see Beem, E.P. 323(1985)439
Hoogmartens, J., see De Pourcq, P. 321(1985)441
Hoogmartens, J., see Dihuidi, K. 325(1985)413
Hoogmartens, J., see Roets, E. 303(1984)117
Hooke, K.F., see Duggan, D.E. 306(1984)69

Hoppel, C.L., see Davis, A.T. 306(1984)79
Hoppel, C.L., see Litowitz, H. 311(1984)443
Hopper, M.L.
Capillary fused-silica on-column injection of chlorinated pesticides with an ultra-low volume rotary valve 302(1984)205
Hori, K., see Matsuhashi, S. 319(1985)79
Horie, M., see Baba, S. 307(1984)1
Horioka, M., see Nakano, Y. 311(1984)390
Horisaka, K., see Yamada, J. 311(1984)385
Horn, M., see Sârbu, C. 322(1985)272
Horna, A., Táborský, J., Dufka, O., Matoušek, P. and Churáček, J.
Chromatography of monomers. II. Glass capillary GC of C_1-C_{18} alkyl esters of acrylic and methacrylic acids 325(1985)367
Hornsperger, J.-M., Wagner, J., Hinkel, J.-P. and Jung, M.J.
Measurement of 3-methoxy-4-hydroxyphenylglycol sulfate ester in brain using reversed-phase LC and electrochemical detection 306(1984)364
Horowitz, P.M.
Rapid fluorescamine based protein assay usable in the presence of interfering substances 319(1985)446
Horvai, G., see Niegreisz, Z. 316(1984)451
Horváth, C., see Chang, J.P. 318(1985)11
Horváth, C., see Chang, J.-P. 319(1985)396
Horváth, C., see Corradini, D. 319(1985)400
Horváth, C., see Huang, J.-X. 316(1984)151
Horváth, C., see Jacobson, J. 316(1984)53
Horváth, C., see Melander, W.R. 317(1984)67, 318(1985)1
Hoshino, T.
Simple device for sequential linear gradient elution with a single, conventional, one-piston reciprocating pump 316(1984)415
Hoshino, T., Senda, M., Hondo, T., Saito, M. and Tohei, S.
Application of photodiode array UV detector to unresolved peak analysis 316(1984)473
Hoskins, J.A., see Bailey, E. 310(1984)199
Hosojima, H., see Imaizumi, N. 308(1984)295
Hostettmann, K., see Domon, B. 315(1984)441
Hostomská, Z., see Mikeš, O. 301(1984)93
Hotta, K., see Iwase, H. 320(1985)426
Houghton, G.W., see Gulaid, A.A. 318(1985)393
Houwelingen, J.C. van, see De Ligny, C.L. 301(1984)311
Howell, J.A., see Priebe, S.R. 324(1985)53
Howell, J.B., see Shea, P.A. 306(1984)358
Howells, R.E., see Coleman, M.D. 308(1984)363
Hradil, J., Svec, F., Gogitidze, N.M. and Andronikashvili, T.T.
GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277
Hruby, V.J., see Cody, W.L. 314(1984)313
Hsia, J.C., Hayes, T.M. and Er, S.S.
Separation of glyoxylated hemoglobin with varying oxygen affinity by polyanion fast protein LC 303(1984)425
Hsia, J.C., see Wong, L.T. 310(1984)19
Hsu, D.-S. and Chen, S.S.
Rapid analysis of adenosine, AMP, ADP, and ATP by anion-exchange CC 311(1984)396
Huang, J.-C.
Linear relationship in the thermodynamics of GLC 321(1985)458
Huang, J.-X., Stuart, J.D., Melander, W.R. and Horváth, C.
HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. I. Interplay of on-column redox reaction and the chromatographic retention process 316(1984)151
Hubball, J.A., Dimauro, P.R., Smith, S.R. and Barry, E.F.
Preliminary investigation of the effects of gamma-radiation on the surface of raw fused-silica tubing 302(1984)341
Hubbard, J.W., see Yeung, P.K.F. 303(1984)412
Hubert, J., see Hanai, T. 302(1984)89, 316(1984)261
Hubert, P., see Mathis, R. 319(1985)43
Hudec, M., see Madajová, V. 320(1985)131
Huhtikangas, A., see Herranen, J. 307(1984)241
Hui, D.S.S., see Hui, K.-S. 325(1985)467
Hui, J.Y. and Taylor, S.L.
Reversed-phase ion-pair HPLC procedure for determination of histamine and its metabolites in rat urine 312(1984)443
Hui, K.-S., Hui, D.S.S. and Chan, K.K.H.
Separation of enkephalins from their metabolites and precursor by a single pump gradient HPLC method 325(1985)467
Hülm, M., see Pötter, H. 319(1985)440
Hulshoff, A., see Barends, D.M. 322(1985)321
Hulshoff, A., see Den Hartigh, J. 306(1984)444
Hulsman, J.A.R.J., see Lemmens, A.A.G. 320(1985)193
Hume, A.S., see Purser, C. 311(1984)135
Hung, C.T., see Taylor, R.B. 316(1984)279
Hung, Y.-L., Kai, M., Nohta, H. and Ohkura, Y.
HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281
Hunt, J.E., see Schaber, P.M. 316(1984)25
Hunter, A.T.
Separation of anthraquinone sulphonic acids by ion-pair LC 319(1985)319
Hunter, K.
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255
Hunter, M.L., see Christie, W.W. 325(1985)473
Hunzicker-Dunn, M. and Sutherland, D.B.
Elution of the catalytic subunits of the type I and type II forms of cyclic AMP-dependent protein kinase within the type I chromatographic peak 312(1984)293
Hurst, W.J., Snyder, K.P. and Martin, R.A. Jr.
Use of microbore HPLC for the determination of caffeine, theobromine and theophylline in cocoa 318(1985)408
Hurtubise, R.J., see Cooper, H.A. 324(1985)1

Hurtubise, R.J., see Ruckmick, S.C. 321(1985)343
Hušek, P. and Felt, V.
Possibilities and limitations in the analysis of amino acid oxazolidinones in the femtomole range by GC with electron-capture detection 305(1984)442
Hutchison, J.S., see Evans, L.J. Jr. 305(1984)163
Hutta, M., see Koval, M. 325(1985)151
Hutte, R.S., Williams, E.J., Staehelin, J., Hawthorne, S.B., Barkley, R.M. and Sievers, R.E.
Chromatographic analysis of organic compounds in the atmosphere 302(1984)173
Huuskonen, M., see Herranen, J. 307(1984)241
Hyun, M.H., see Pirkle, W.H. 316(1984)585, 322(1985)287, 322(1985)295, 322(1985)309
Hyver, K.J., see Parcher, J.F. 302(1984)195

I

Idvall, J., see Björkman, S. 307(1984)481
Ignatiadis, I., Schmitter, J.M. and Arpino, P.
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87
Iizuka, H., see Yoshioka, M. 309(1984)63
Ijima, S., see Okano, Y. 310(1984)251
Ikarashi, Y., Sasahara, T. and Maruyama, Y.
Determination of choline and acetylcholine levels in rat brain regions by LC with electrochemical detection 322(1985)191
Ikeda, M., Hattori, H., Koyama, Y. and Ohmori, S.
Determination of urochloralic acid, the glucuronic acid conjugate of trichloroethanol, by GC with electron-capture detection and its application to urine, plasma and liver 307(1984)111
Ikeda, M., Shimada, K., Sakaguchi, T. and Matsumoto, U.
Fluorometric HPLC of 9-aminophenanthrene-derivatized free fatty acids 305(1984)261
Ikeda, M., see Mikasa, H. 305(1984)204
Ikigai, H., Nakae, T. and Kato, Y.
Analysis of membrane proteins by ion-exchange HPLC 322(1985)212
Imai, K., see Miyaguchi, K. 303(1984)173, 316(1984)501
Imai, K., see Watanabe, Y. 309(1984)279
Imaizumi, N., Morimoto, S., Kigoshi, T., Uchida, K., Hosojima, H. and Yamamoto, I.
Application of reversed-phase HPLC for radioimmunoassay of plasma 18-hydroxycorticosterone 308(1984)295
Imamura, T., Sugihara, J., Yokota, E., Kagimoto, M., Naito, Y. and Yanase, T.
Analytical peptide mapping by ion-exchange HPLC: application to haemoglobin variants 305(1984)456
Imanari, T., see Toida, T. 308(1984)133
Inchauspé, G. and Samain, D.
Use of perfluorinated carboxylic acids in the separation of aminoglycoside antibiotics by ion-pair reversed-phase HPLC 303(1984)277
Ingalls, S.T., see Davis, A.T. 306(1984)79
Inoue, T., see Baba, S. 305(1984)119
Ionescu, T., see Drost, R.H. 310(1984)193
Ireland, P.A. and Dziedzic, S.Z.
Analysis of soybean sapogenins by HPLC 325(1985)275
Irikura, T., see Ishida, R. 305(1984)73
Isaksson, K., Lindquist, J. and Lundström, K.
Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333
Ishida, J., Yamaguchi, M., Kai, M., Ohkura, Y. and Nakamura, M.
Determination of bestatin in serum by HPLC with fluorescence detection 305(1984)381
Ishida, R., Ozaki, T., Uchida, H. and Irikura, T.
GC-MS determination of amitriptyline and its major metabolites in human serum 305(1984)73
Ishida, S., see Tsuruta, Y. 309(1984)309
Ishii, D., Goto, M. and Takeuchi, T.
Multi-channel detectors for micro HPLC: examination of flow cell structures 316(1984)441
Ishii, D., see Takeuchi, T. 321(1985)159
Ishikawa, K., Martinez, J.L. Jr. and McGaugh, J.L.
Simple determination of *p*-hydroxyamphetamine by HPLC with electrochemical detection 306(1984)394
Ishioka, N., see Takahashi, N. 317(1984)11
Ishizaki, T., see Minegishi, A. 308(1984)55, 311(1984)51
Isozaki, S., see Uchino, K. 308(1984)241
Issaq, H.J., Klose, J. and Muschik, G.M.
Separation of polychlorinated biphenyls (Aroclor 1254) by HPLC 302(1984)159
Itkonen, A., see Peltonen, K. 315(1984)412
Ito, K., see Kurosawa, N. 305(1984)485
Ito, S., Kato, T., Maruta, K., Fujita, K. and Kurahashi, T.
Determination of DOPA, dopamine, and 5-S-cysteinyl-DOPA in plasma, urine, and tissue samples by HPLC with electrochemical detection 311(1984)154
Ito, Y.
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. I. Hydrodynamic

distribution of two solvent phases in a helical column subjected to two types of synchronous planetary motion 301(1984)377
Ito, Y.
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. II. Phase distribution diagrams for helical and spiral columns 301(1984)387
Ito, Y. and Conway, W.D.
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405
Ito, Y., see Knight, M. 301(1984)277
Ito, Y., see Nakazawa, H. 307(1984)323
Iwado, A., see Yamamoto, S. 303(1984)99
Iwahashi, H., see Morishita, H. 315(1984)253
Iwamoto, M., Yoshida, S. and Hirose, S.
Fluorescence determination of 5-fluorouracil and 1-(tetrahydro-2-furanyl)-5-fluorouracil in blood serum by HPLC 310(1984)151
Iwasaki, H., see Tanaka, A. 306(1984)51
Iwase, H., Kato, Y. and Hotta, K.
Fractionation of peptic glycopeptides from chicken ovalbumin by reversed-phase HPLC 320(1985)426
Iwatani, A. and Nakamura, H.
Determination of urinary tryptophan, 5-hydroxytryptamine and 5-hydroxyindoleacetic acid in neonatal hyperbilirubinaemic infants using reversed-phase HPLC with fluorescence detection 309(1984)145
Iyengar, J.R., see Lawrence, J.F. 325(1985)299

J

Jack, D.B., see Gregg, M.R. 305(1984)244
Jack, D.B., see Lagerström, P.-O. 307(1984)230
Jackman, G.P., Oddie, C.J., Skews, H. and Bobik, A.
HPLC determination of plasma catecholamines during α-methyldopa therapy 308(1984)301
Jackman, G.P., see Oddie, C.J. 308(1984)370
Jackson, P.J. and Sâmūndsen, J.A.
Improved method for GC determination of phytanic acid 325(1985)336
Jacob, P. III, Elias-Baker, B.A., Jones, R.T. and Benowitz, N.L.
Determination of cocaine in plasma by automated GC 306(1984)173
Jacobs, G.A., see De Beer, J.O. 307(1984)475
Jacobsen, N.W., see Charles, B.G. 306(1984)424
Jacobson, J., Frenz, J. and Horváth, C.
Measurement of adsorption isotherms by LC 316(1984)53
Jähnchen, E., see Bechtold, H. 306(1984)333
Jähnchen, E., see Wiegand, U.-W. 311(1984)218
Jalal, M.A.F., Mocharla, R. and Van der Helm, D.
Separation of ferrichromes and other hydroxamate siderophores of fungal origin by reversed-phase chromatography 301(1984)247
Jamali, F., Russell, A.S., Berry, B.W. and Lehmann, C.
HPLC analysis of tiaprofenic acid and its metabolites in plasma and urine by direct injection 310(1984)327
Jameson, C.W., see Timmons, L. 314(1984)476
Jandera, P.
Reversed-phase LC of homologous series. A general method for prediction of retention 314(1984)13
Jane, I., McKinnon, A. and Flanagan, R.J.
HPLC analysis of basic drugs on silica columns using non-aqueus ionic eluents. II. Application of UV, fluorescence and electrochemical oxidation detection 323(1985)191
Jane, I., see Flanagan, R.J. 323(1985)173
Janjić, T.J., see Vučković, G. 324(1985)431
Jansen, E.H.J.M., Zoontjes, P.W., Van Blitterswijk, H., Both-Miedema, R. and Stephany, R.W.
Fast HPLC screening method for the presence of trenbolone and its major metabolite in urine of slaughter cattle 319(1985)436
Jansen, H., Frei, R.W., Brinkman, U.A.T., Deelder, R.S. and Snellings, R.P.J.
Determination of urea and ammonia using ion-pair LC with on-line post-column derivatization in an enzymatic solid-phase reactor 325(1985)255
Janssen, R.T.P., see Lemmens, A.A.G. 320(1985)193
Jansson, S.-O., see Persson, B.-A. 316(1984)291
Janz, P., see Binder, H. 303(1984)375
Jaramillo, L., see Nazareth, A. 309(1984)357
Jarrott, B., see Drummer, O.H. 305(1984)83
Jastorff, B., see Lüllmann, C. 323(1985)273
Jaworsky, L., see Alton, K.B. 311(1984)319
Jellum, E., see Tong, H.Y. 312(1984)183, 324(1985)373
Jensen, R., see Peuchant, E. 310(1984)297
Jevnikar, J.J., see Lee, T.T. 325(1985)340
Jeyaraj, G.L. and Porter, W.R.
New method for the resolution of racemic warfarin and its analogues using low-pressure LC 315(1984)378
Jezekova, D., see Enger, A. 305(1984)335
Jin, Y., see Takeuchi, T. 321(1985)159
Jin, Z.L., see Xu, X.B. 317(1984)545
Jinno, K. and Kawasaki, K.
Computer-assisted retention prediction

system for reversed-phase micro HPLC 316(1984)1
Johansson, B.-L. and Drevin, I.
Quantitative determination of the ligand content and the negative and positive charges on butylamine-substituted Sepharose 4B 321(1985)335
Johansson, B.-L. and Stafström, N.
Column lifetime in fast protein LC 314(1984)396
Johansson, G.
Determination of ionic charge by liquid-liquid partition 322(1985)425
Johansson, G., Åkerlund, H.-E. and Olde, B.
Liquid-liquid extraction of membranes from calf brain using conventional and centrifugal counter-current distribution techniques 311(1984)277
Johansson, G. and Andersson, M.
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39
Johansson, L., see Antonsson, A.-M. 308(1984)181
Johansson, M.-L., see Persson, B.-A. 316(1984)291
Johnson, D.C., see Polta, J.A. 324(1985)407
Johnson, E.K. and Nurok, D.
Computer simulation as an aid to optimizing continuous-development two-dimensional TLC 302(1984)135
Johnson, E.L., see Daftsios, A.C. 305(1984)145
Johnson, J.S. Jr., see Shiao, S.-Y. 303(1984)13
Johnson, T.C., see Sharifi, B.G. 324(1985)173
Johnston, M.A., Lea, A.R. and Hailey, D.M.
Inter-laboratory study on the separation of cinchona alkaloids by HPLC 318(1985)362
Joly, R., see Dell, D. 317(1984)483
Jones, A.W., Blom, Y., Bondesson, U. and Ånggård, E.
Determination of morphine in biological samples by GC-MS. Evidence for persistent tissue binding in rats twenty-two days post-withdrawal 309(1984)73
Jones, B.A., see Lee, M.L. 302(1984)303
Jones, D.P.
Simple HPLC assay for benzylamine oxidation products in cell suspensions 305(1984)256
Jones, R.S. and Stutte, C.A.
Chromatographic analysis of choline and acetylcholine by UV visualization 319(1985)454
Jones, R.T., see Jacob, P. III 306(1984)173
Jones, V.K. and Tarter, J.G.
Single injection ion chromatographic analysis of both anions and cations 312(1984)456
Jong, J. de, see De Jong, J. 319(1985)23, 322(1985)43
Jonk, R.J.G., see Tas, A.C. 310(1984)243
Joppich, M., see Nazareth, A. 314(1984)201, 319(1985)382
Jordan, N., see Loo, J.C.K. 305(1984)194
Jorgenson, J.W., see Walbroehl, Y. 315(1984)135
Joshi, B.S., see Desai, H.K. 322(1985)223
Joshua, H. and Goetz, M.
Separation of nucleobases on polar amino cyano HPLC columns 303(1984)185
Josić, D., Hofermaas, R., Bauer, C. and Reutter, W.
Automatic amino acid and sugar analysis of glycoproteins 317(1984)35
Jue, D.L., see Moo-Penn, W.F. 318(1985)325
Juenge, E.C., see Allgire, J.F. 325(1985)249
Juergens, U.
Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97
Juguelin, H., see Heape, A.M. 322(1985)391
Julkunen-Tiitto, R.
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129
Jung, M.J., see Hornsperger, J.-M. 306(1984)364
Jusko, W.J., see Carson, S.W. 306(1984)345
Jusko, W.J., see Ebling, W.F. 305(1984)271

K

Kabra, P.M., Bhatnager, P.K. and Nelson, M.A.
LC determination of amikacin in serum with spectrophotometric detection 307(1984)224
Kádár Pauncz, J.
TLC investigation of the antimycin A antibiotic complex 322(1985)386
Kadofuku, T. and Sato, T.
Detection of changes in rabbit serum proteins after partial hepatectomy by means of two-dimensional electrophoresis under non-denaturing conditions 311(1984)93
Kadota, T., see Okano, Y. 310(1984)251
Kadowaki, H., Shiino, M., Uemura, I. and Kobayashi, K.
Sensitive method for the determination of diclofenac in human plasma by GC-MS 308(1984)329
Kadowaki, S., see Oka, H. 325(1985)265
Kafarski, P., see Lejczak, B. 324(1985)455
Kågedal, B. and Källberg, M.
Cysteinylglycine in urine determined by HPLC 308(1984)75
Kågedal, B., Källberg, M. and Mårtensson, J.
Determination of non-protein-bound N-acetylcysteine in plasma by HPLC 311(1984)170
Kagi, R.I., see Rowland, S.J. 312(1984)395

Kagi, R.I., see Wajon, J.E. 319(1985)187
Kagimoto, M., see Imamura, T. 305(1984)456
Kahle, V. and Krejčí, M.
Role of extra-column volume in the efficiency of high-speed LC 321(1985)69
Kahle, V., see Kovář, J. 319(1985)341
Kai, M., Miyazaki, T. and Ohkura, Y.
HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257
Kai, M., Miyazaki, T., Sakamoto, Y. and Ohkura, Y.
Use of benzoin as pre-column fluorescence derivatization reagent for the HPLC of angiotensins 322(1985)473
Kai, M., see Hung, Y.-L. 305(1984)281
Kai, M., see Ishida, J. 305(1984)381
Kakehi, K., Konishi, T., Sugimoto, I. and Honda, S.
Post-column reaction of amino acids with the pentane-2,4-dione-formaldehyde system for their automated analysis 318(1985)367
Kalász, H., Kerecsen, L. and Nagy, J.
Conditions and parameters dominating displacement TLC 316(1984)95
Kalász, H., see Nagy, J. 317(1984)165
Källberg, M., see Kågedal, B. 308(1984)75, 311(1984)170
Kalo, P.
Microanalytical methods in the structure elucidation of sex-specific components in the large pine weevil, *Hylobius abietis* L. (*Coleoptera, Curculionidae*) 323(1985)343
Kamata, K. and Motohashi, N.
Separation of methyl-substituted benz[c]acridines by GLC and HPLC 319(1985)331
Kamblawi, M.O., Stevens, R.G. and Nicholls, P.J.
HPLC assay for aminoglutethimide and its acetylated metabolite in urine 309(1984)431
Kamei, M., see Tokuda, M. 323(1985)434
Kamiyama, F., see Dohji, T. 311(1984)249
Kamp, D., see Kamp, R.M. 317(1984)181
Kamp, R.M., Bosserhoff, A., Kamp, D. and Wittmann-Liebold, B.
Application of HPLC techniques to the separation of ribosomal proteins of different organisms 317(1984)181
Kaneko, M., see Kiba, N. 303(1984)396
Kaniansky, D., see Havaši, P. 325(1985)137
Kaniansky, D., see Koval, M. 325(1985)151
Kaniansky, D., see Madajová, V. 320(1985)131
Kaniansky, D., see Zelenský, I. 325(1985)161
Kanko, S. and Autio, K.
Electrophoretic studies of blood globin preparations 324(1985)395
Kanohta, K., see Kato, S. 324(1985)462
Kapetanovic, I.M. and Kupferberg, H.J.
Analysis of *p*-hydroxyphenytoin in microsomal reactions by HPLC with electrochemical detection 310(1984)418
Kapil, R.P., Abbott, F.S., Kerr, C.R., Edwards, D.J., Lalka, D. and Axelson, J.E.
Simultaneous quantitation of disopyramide and its mono-dealkylated metabolite in human plasma by fused-silica capillary GC using nitrogen-phosphorus specific detection 307(1984)305
Karasek, F.W., see Crouch, R.L. 303(1984)53
Karasek, F.W., see Hawley-Fedder, R.A. 314(1984)263, 315(1984)201, 315(1984)211
Karasek, F.W., see Tong, H.Y. 312(1984)183, 324(1985)373
Karege, F.
Method for total 3-methoxy-4-hydroxyphenylglycol extraction from urine, plasma and brain tissue using bonded-phase materials: comparison with the ethyl acetate extraction method 311(1984)361
Karger, B.L. and Vouros, P.
Chromatographic perspective of HPLC-MS 323(1985)13
Karger, B.L., see Benedek, K. 317(1984)227
Karger, B.L., see Miller, N.T. 316(1984)519
Karger, B.L., see Nazareth, A. 309(1984)357
Karger, B.L., see Yu, T.J. 301(1984)425
Karlsson, L. and Roos, C.
Combination of LC with UV detection and GC with electron-capture detection for the determination of Δ^9-tetrahydrocannabinol-11-oic acid in urine 306(1984)183
Karmen, A., Malikin, G. and Lam, S.
High-sensitivity radioassay in chromatographic effluents 302(1984)31
Karmen, A., see Bock, J.L. 308(1984)354
Karmen, A., see Lam, S. 302(1984)21
Kasahara, Y., Hikino, H. and Hine, T.
Determination of ephedrine alkaloids by isotachophoresis 324(1985)503
Kasai, K.-I., see Shimura, K. 315(1984)161
Kašička, V. and Prusík, Z.
Desorption isotachophoresis: quantitative characterization of sorption and desorption conditions 320(1985)75
Kašička, V., Vacík, J. and Prusík, Z.
Determination of dissociation constants of weak electrolytes by capillary isotachophoresis 320(1985)33
Kašička, V., see Prusík, Z. 320(1985)81
Kataoka, H., Yamamoto, S. and Makita, M.
Quantitative GLC of taurine 306(1984)61
Katayama, K., see Kuwada, M. 308(1984)398
Kates, R.E. and Latini, R.
Simple and rapid HPLC analysis of cyclosporine in human blood and serum 309(1984)441
Kato, S. and Kanohta, K.
Chromatographic studies of the autoxidation products of ethoxyquin and its photochemical conversion 324(1985)462
Kato, T., see Ito, S. 311(1984)154
Kato, Y., Nakamura, K., Yamazaki, Y. and Hashimoto, T.
Comparison of high-performance ion-exchange chromatography and gel

electrophoresis in protein separations 318(1985)358
Kato, Y., Yamasaki, Y., Hashimoto, T., Murotsu, T., Fukushige, S. and Matsubara, K.
Separation of large DNA restriction fragments by high-performance gel filtration on TSKgel DNA-PW 320(1985)440
Kato, Y., see Ikigai, H. 322(1985)212
Kato, Y., see Iwase, H. 320(1985)426
Katoh, K., see Miyazaki, K. 310(1984)219
Katz, E.Z., Granit, L., Drayer, D.E. and Levy, M.
Simultaneous determination of dipyrone metabolites in plasma by HPLC 305(1984)477
Katz, J.J., see Schaber, P.M. 316(1984)25
Katzenstein, G.E., see Sadler, A.J. 317(1984)93
Kauffman, R.E., see Aravind, M.K. 308(1984)350
Kauffman, R.E., see Bowman, D.B. 309(1984)209
Kawabe, T., see Dohji, T. 311(1984)249
Kawaguchi, Y., see Tanaka, M. 301(1984)345, 314(1984)193
Kawai, S., see Nakamura, M. 310(1984)450
Kawamoto, T., Mashimo, I., Yamauchi, S. and Watanabe, M.
Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373
Kawasaki, K., see Jinno, K. 316(1984)1
Kawase, M., see Hirota, K. 310(1984)204
Kay, B., see Keegan, M. 311(1984)223
Keegan, M. and Kay, B.
Detection of nalbuphine in plasma: an improved HPLC assay 311(1984)223
Keeley, F.J., see Daftsios, A.C. 305(1984)145
Kees, F., Grobecker, H. and Naber, K.G.
HPLC analysis of cefotetan epimers in human plasma and urine 305(1984)363
Keiser, H.R., see Goldstein, D.S. 311(1984)148
Kelleher, F.M., see Trumbore, C.N. 322(1985)443
Keller, S., Ghosh, A.K., Ghosh, A.K., Turino, G.M. and Mandl, I.
Separation of the cross-linking amino acids of elastin on thin-layer plates 305(1984)461
Kellert, M., see Tiebach, R. 318(1985)103, 323(1985)121
Kelley, J. and Chrin, L.
TLC of hydroxylysine for collagen analysis 311(1984)400
Kelley, J.L., see Williams, M.C. 308(1984)101
Kempf, A., see Honegger, C.G. 309(1984)53
Kemula, W., see Głód, B. 321(1985)433
Kennedy, L.A., see Fausnaugh, J.L. 317(1984)141
Keough, T., Strife, R.J., Rodriguez, P.A. and Sanders, R.A.
Separation and use of the perdeuterated analogue as an internal standard for the analysis of ethylene dibromide 312(1984)450
Kerecsen, L., see Kalász, H. 316(1984)95
Kerlavage, A.R., Weitzmann, C.J. and Cooperman, B.S.
Application of HPLC to the reconstitution of ribosomal subunits 317(1984)201
Kern, J.W., see Terry, M.D. 311(1984)213
Kerr, C.R., see Kapil, R.P. 307(1984)305
Kershaw, R.A., see Mays, D.C. 311(1984)301
Kester, A.S. and Thompson, R.E.
Computer-optimized normal-phase HPLC separation of *Corynebacterium poinsettiae* carotenoids 310(1984)372
Khaledi, M.G., see Dorsey, J.G. 316(1984)183
Khatib, S., see Poole, C.F. 302(1984)149
Khazanchi, R., see Kulshrestha, G. 318(1985)144
Kholy, A.M. el, see Brooks, J.B. 309(1984)269
Khurana, V.K., see Sharifi, B.G. 324(1985)173
Kiba, N. and Kaneko, M.
Use of immobilized amino acid oxidase as post-column reactor in the HPLC of amino acids 303(1984)396
Kibler, S., see Pike, A.W. 306(1984)39
Kido, R., see Morishita, H. 315(1984)253
Kiel, J.S., Morgan, S.L. and Abramson, R.K.
Effects of amine modifiers on retention and peak shape in reversed-phase HPLC 320(1985)313
Kigoshi, T., see Imaizumi, N. 308(1984)295
Kikuchi, H., see Hirokawa, T. 305(1984)429
Kim, J.H., see Kim, K.R. 325(1985)127
Kim, K.R., Zlatkis, A., Weisner, S., Kim, J.H. and Choi, K.S.
Trace enrichment of indole-3-acetic acid from leaf matrix 325(1985)127
Kim, S.-G., Chatfield, J.M. and Armstrong, D.J.
Trace enrichment of cytokinins using neutral polystyrene resins 321(1985)175
Kim, Y.-H., Woodrow, J.E. and Seiber, J.N.
Evaluation of a GC method for calculating vapor pressures with organophosphorus pesticides 314(1984)37
Kinkel, A.W., see Randinitis, E.J. 307(1984)210, 308(1984)345
Kinkel, J.N. and Unger, K.K.
Role of solvent and base in the silanization reaction of silicas for reversed-phase HPLC 316(1984)193
Kinney, C.D.
Estimation of nadolol levels in plasma using HPLC with recirculating eluent flow 305(1984)489
Kinoshita, T., see Nimura, N. 316(1984)547
Kirchhoefer, R.D., see Allgire, J.F. 325(1985)249
Kirkland, J.J., see Dilks, C.H. Jr. 315(1984)45
Kirkland, J.J., see Głajch, J.L. 318(1985)23
Kirkman, C.M., Zu-Ben, C., Uden, P.C., Stratton, W.J. and Henderson, D.E.
Developments in the HPLC of metallo-organic compounds 317(1984)569
Kishie, T., see Hirota, K. 310(1984)204
Kiso, Y., see Hirokawa, T. 305(1984)429, 312(1984)11, 318(1985)195
Kiss, B., see Oka, K. 308(1984)43

Kissa, E.
Determination of 2H-hexafluoropropanol-2 in air 319(1985)407
Kissinger, L.D. and Robins, R.H.
Silver-modified mobile phase for normal-phase LC determination of prostaglandins and their 5,6-*trans* isomers in prostaglandin bulk drugs and triacetin solutions 321(1985)353
Kissinger, P.T., see Lunte, C.E. 317(1984)407
Kissinger, P.T., see Lunte, S.M. 317(1984)579
Kitayama, M., see Suemitsu, R. 318(1985)139
Kjølholt, J.
Determination of trace amounts of organophosphorus pesticides and related compounds in soils and sediments using capillary GC and a nitrogenphosphorus detector 325(1985)231
Klassen, R., see Stavric, B. 310(1984)107
Klasson, K., see Bergenhem, N. 319(1985)59
Klecker, R.W. Jr. and Collins, J.M.
Quantification of tiazofurin in plasma by HPLC 307(1984)361
Kleijn, E. van der, see Oosterbaan, M.J.M. 306(1984)323
Klein, F., see Bechtold, H. 306(1984)333
Kleinbloesem, C.H., Van Harten, J., Van Brummelen, P. and Breimer, D.D.
LC determination of nifedipine in plasma and of its main metabolite in urine 308(1984)209
Kleinert, T. and Ecknig, W.
Prediction of retention data by using parameters of intermolecular interaction. I. A model for calculation of non-polar and polar interaction parameters 315(1984)75
Kleinert, T., Ecknig, W. and Novák, J.
Prediction of retention data by using parameters of intermolecular interaction. II. Application of the model to GLC systems 315(1984)85
Klesper, E., see Leyendecker, D. 315(1984)19, 321(1985)273
Klett, R.P., Chovan, J.P. and Danse, I.H.R.
Reversed-phase paired-ion HPLC method for the separation and quantification of multiple bleomycin congeners 310(1984)361
Klinger, J.K., see Luzzi, F.A. 311(1984)291
Klose, J., see Issaq, H.J. 302(1984)159
Klus, H., see Begutter, H. 321(1985)475
Kluyev, N.A., see Zamureenko, V.A. 303(1984)109
Knight, M., Tamminga, C.A., Ito, Y., Gardner, J.D. and Chase, T.N.
Purification of a cholecystokinin fragment peptide in the horizontal flow-through coil planet centrifuge 301(1984)277
Knoll, J., see Nagy, J. 317(1984)165
Knox, J.H. and Scott, H.P.
Theoretical models for size-exclusion chromatography and calculation of pore size distribution from size-exclusion chromatography data 316(1984)311
Knox, M.J., see Ashoor, S.H. 324(1985)199
Knutson, D.E., see Bauza, M.T. 310(1984)61
Kobayashi, A., see Hattori, Y. 315(1984)341
Kobayashi, K., see Kadowaki, H. 308(1984)329
Kobayashi, S., Schwartz, S.J. and Lineback, D.R.
Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 319(1985)205
Kobayashi, S., see Hirokawa, T. 318(1985)195
Koch, D.D. and Polzin, G.
Determination of urinary 3-methoxy-4-hydroxyphenylglycol by LC with electrochemical detection 317(1984)327
Kochak, G. and Honc, F.
Improved GLC method for the determination of baclofen in plasma and urine 310(1984)319
Kodama, H., Sasaki, K., Mikasa, H., Cavallini, D. and Ricci, G.
Quantitative analysis of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using isotachophoresis 311(1984)183
Kodama, H., see Mikasa, H. 305(1984)204, 310(1984)401
Kodama, K., Nakata, T. and Aoyama, M.
Direct determination of urinary vanillylmandelic acid and homovanillic acid by HPLC on an anion-exchange column 311(1984)369
Kogan, M.J., Newman, E. and Willson, N.J.
Detection of marijuana metabolite 11-nor-Δ^9-tetrahydrocannabinol-9-carboxylic acid in human urine by bonded-phase adsorption and TLC 306(1984)441
Kohashi, K., see Tsuruta, Y. 309(1984)309
Kohli, J.C. and Badaisha, K.K.
Improved procedure for the TLC of terpenoids on silver ion-silica gel layers 320(1985)455
Kohri, N., see Miyazaki, K. 310(1984)219
Koizumi, K., Utamura, T. and Okada, Y.
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145
Kojima, K., see Oka, K. 308(1984)43
Kojima, M., see Nakajima, K. 301(1984)241
Kok, W.T., Groenendijk, G., Brinkman, U.A.T. and Frei, R.W.
Determination of oxalic acid in biological matrices by LC with amperometric detection 315(1984)271
Kolarovic, L., Traitler, H. and Ducret, P.
On-line GC evaluation of iodine values in edible fats and oils 314(1984)233
Kołodziejczyk, A.M., see Ślebioda, M. 325(1985)282
Komori, T., see Maoka, T. 318(1985)122
Kondo, K., see Nakamura, M. 310(1984)450
König, W.A., Steinbach, E. and Ernst, K.
Phosgene: a versatile reagent for enantiomer separation by capillary GC 301(1984)129
Koningsveld, R., see Glöckner, G. 317(1984)615

Konishi, T., see Kakehi, K. 318(1985)367
Konomi, H., Hayashi, T., Takehana, Y. and Arima, M.
Rapid and sensitive amino acid analysis of human collagens using HPLC 311(1984)375
Koopmans, P.P., Tan, Y., Van Ginneken, C.A.M. and Gribnau, F.W.J.
HPLC determination of hydrochlorothiazide in plasma and urine 307(1984)445
Kopaciewicz, W., Rounds, M.A. and Regnier, F.E.
Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157
Køppen, B. and Dalgaard, L.
Headspace GC determination of β-galactosidase activity using electron-capture detection 321(1985)385
Korhonen, I.O.O.
GLC analyses. XXIX. Separation of free chlorophenol isomers on non-polar and polar quartz capillary columns 303(1984)197
Korhonen, I.O.O.
GLC analyses. XXXI. Retention increments of isomeric chlorophenols on low-polarity (SE-30) and polar (FFAP) capillary columns 315(1984)185
Korhonen, I.O.O.
GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115
Korhonen, I.O.O.
GLC analyses. XXXIII. Retention increments for *o*-chloro and *o*-methoxy substitution in isomeric chloroanisoles 321(1985)467
Korhonen, I.O.O.
GLC analyses. XXXIX. ω-Chloroethanols on low-polarity (SE-30) and polar (OV-351) capillary columns 324(1985)181
Korhonen, I.O.O.
GLC analyses. XL. Incremental effects of a formyl group introduced into the *ortho*- and *para*-positions of isomeric chlorophenols 324(1985)192
Korhonen, I.O.O. and Lind, M.A.
GLC analyses. XXXIV. Separation and retention indices with retention increments of some nitrated polynuclear aromatic hydrocarbons on a low-polarity (SE-30) capillary column 322(1985)71
Korhonen, I.O.O. and Lind, M.A.
GLC analyses. XXXV. Capillary column studies of C_1-C_{12} *n*-alkyl esters of benzoic and monochlorobenzoic acids 322(1985)83
Korhonen, I.O.O. and Lind, M.A.
GLC analyses. XXXVI. Separation of ethyl, 2-chloroethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns with temperature programming 322(1985)97
Korhonen, I.O.O. and Lind, M.A.
GLC analyses. XXXVII. Capillary column studies of benzoyl and monochlorobenzoyl esters of lower saturated branched-chain alcohols 323(1985)331
Korhonen, I.O.O. and Lind, M.A.
GLC Analyses. XXXVIII. Capillary column studies of unsaturated esters of benzoic and monochlorobenzoic acids 324(1985)113
Korhonen, I.O.O. and Lind, M.A.
GLC analyses. XLI. Ethyl and ω-chloroethyl esters of benzoic and monochlorobenzoic acids on SE-30 and OV-351 capillary columns 325(1985)433
Korhonen, I.O.O., see Haken, J.K. 319(1985)131, 320(1985)325, 324(1985)343, 325(1985)61
Koroknay, B., see De Vries, J.X. 309(1984)219
Korpela, T.K. and Saarinen, R.
Affinity chromatography of B_6-vitamin-dependent enzymes: purification of pig-heart branched-chain amino acid transaminase 318(1985)333
Korshevetz, I.K., see Mistryukov, E.A. 322(1985)206
Körtvélyessy, G., Szórádi, S., Sztruhar, I. and Ladányi, L.
GC of 3,7,11-trimethyl-11-hydroxy (or -methoxy-)-2,4-dodecadienoic acids and related compounds as their methyl or trimethylsilyl esters 303(1984)370
Košík, M., see Polonský, J. 320(1985)111
Kostowa, W., see Arsov, A. 322(1985)267
Kou, A.Y., see Chen, S.S.-H. 307(1984)261
Kovács, G.H.
Detection of phosmethylan and its major metabolites by TLC 303(1984)309
Kovács, G.H.
HPLC analysis of the photodegradation products of phosmethylan 322(1985)265
Koval, M., Kaniansky, D., Hutta, M. and Lacko, R.
Analysis of saturated normal fatty acids in hydrocarbon matrices by capillary isotachophoresis 325(1985)151
Kovář, J. and Stejskal, J.
Rapid chromatographic purification of yeast alcohol dehydrogenase 325(1985)359
Kovář, J., Turánek, J., Hlaváč, C., Vala, V. and Kahle, V.
LC separations of dimers of nicotinamide adenine dinucleotide and nicotinamide adenine dinucleotide phosphate 319(1985)341
Koyama, F., see Tokuda, M. 323(1985)434
Koyama, Y., see Ikeda, M. 307(1984)111
Koziol, T., Grayeski, M.L. and Weinberger, R.
Determination of trace levels of steroids in blood plasma by LC with peroxyoxalate chemiluminescence detection 317(1984)355
Kozloski, R.P.
Polychlorinated biphenyl retention time standards obtained by chemical dechlorination of polychlorinated biphenyl isomers 318(1985)211

Kozu, T.
HPLC determination of nitrazepam and its metabolites in human urine 310(1984)213
Kraak, J.C., see Thus, J.L.G. 320(1985)271
Kraak, J.C., see Welling, P. 321(1985)450
Kraal, I., see Peters, G.J. 307(1984)464
Kracht, W.R., see Patrick, D.W. 318(1985)269
Krainhöfner, A., see Schade, W. 319(1985)475
Králová, B., see Šafařík, I. 303(1984)283
Kramer, N.C., see Phillips, T.M. 317(1984)173
Kraml, J., see Kramlová, M. 303(1984)422
Kramlová, M., Přistoupil, T.I., Fričová, V. and Kraml, J.
Fast protein liquid chromatofocusing of lyophilized native haemoglobin and its chemically modified derivatives 303(1984)422
Kraus, G., see Moegel, H-J. 324(1985)29
Kraus, G., see Soják, L. 323(1985)414
Krebs, H.A., Cheng, L.K. and Wright, G.J.
Determination of fenfluramine and norfenfluramine in plasma using a nitrogen-sensitive detector 310(1984)412
Krejčí, M., see Kahle, V. 321(1985)69
Krishnamurti, C.R., Heindze, A.M. and Galzy, G.
Application of reversed-phase HPLC using pre-column derivatization with *o*-phthaldialdehyde for the quantitative analysis of amino acids in adult and fetal sheep plasma, animal feeds and tissues 315(1984)321
Krivit, W., see Tuchman, M. 307(1984)172
Krohn, J., see Boo, A.T. 301(1984)335
Krol, G.J., Noe, A.J., Yeh, S.C. and Raemsch, K.D.
GC and LC analyses of nimodipine calcium antagonist in blood plasma and cerebrospinal fluid 305(1984)105
Krüger, J., Rabe, H., Reichmann, G. and Rüstow, B.
Separation and determination of diacylglycerols as their naphthylurethanes by HPLC 307(1984)387
Krüger, R., Mengel, I. and Kuss, H.J.
Determination of benperidol in human plasma by HPLC 311(1984)109
Krull, I.S., see Driscoll, J.N. 302(1984)43
Krummen, K., see Buck, R.H. 303(1984)238, 315(1984)279
Krupčík, J., Garaj, J., Čellár, P. and Guiochon, G.
Calculation of the peak capacity in capillary GC 312(1984)1
Krupčík, J., see Matisová, E. 303(1984)151
Kubo, I., Matsumoto, A., Hanke, F.J. and Ayafor, J.F.
Analytical droplet counter-current chromatography isolation of 20-hydroxyecdysone from *Vitex thyrsiflora* (Verbenaceae) 321(1985)246
Kubota, Y., see Shoji, S. 319(1985)359
Kucharski, M.J., see Dihuidi, K. 325(1985)413
Kuei, J.C., see Lee, M.L. 302(1984)303
Kuge, Y., see Hattori, Y. 315(1984)341
Kuge, Y., see Tanaka, M. 301(1984)345
Kuhn, S., see Grob, K. Jr. 301(1984)1
Kulshrestha, G. and Khazanchi, R.
Reversed-phase HPLC method for the simultaneous microquantitative determination of the herbicide isoproturon and its possible degradation products in soil 318(1985)144
Kuo, K.C., see Gehrke, C.W. 301(1984)199
Kupferberg, H.J., see Kapetanovic, I.M. 310(1984)418
Kurahashi, T., see Ito, S. 311(1984)154
Kurihara, H., see Ujimoto, K. 301(1984)57
Kuroiwa, Y., see Tanaka, E. 308(1984)393
Kurono, M., see Baba, S. 305(1984)119
Kurosawa, N., Morishima, S., Owada, E. and Ito, K.
Reversed-phase HPLC determination of salbutamol in rabbit plasma 305(1984)485
Kushwaha, R.S., see Williams, M.C. 308(1984)101
Kuss, E., see Dibbelt, L. 322(1985)371
Kuss, H.J., see Krüger, R. 311(1984)109
Küsters, E., see Bayer, E. 320(1985)393
Kuwada, M. and Katayama, K.
Improved method for the determination of γ-carboxyglutamic acid in human urine by HPLC 308(1984)398
Kuwamoto, T., see Okada, T. 325(1985)327
Kwan, T.K., Taylor, N.F. and Gower, D.B.
Use of steroid profiling in the resolution of pregnenolone metabolites from porcine testicular preparations 301(1984)189
Kwoka, M., see Ray, G. 317(1984)455
Kylberg-Hanssen, K., see Antonsson, A.-M. 308(1984)181
Kysilka, R., Wurst, M., Pacáková, V., Štulík, K. and Haškovec, L.
HPLC determination of hallucinogenic indoleamines with simultaneous UV photometric and voltammetric detection 320(1985)414

L

Labadarios, D., Moodie, I.M. and Shephard, G.S.
GC analysis of amino acids in physiological fluids: a critique (Review) 310(1984)223
Labadarios, D., Shephard, G.S., Moodie, I.M. and Botha, E.
Effect of the presence of glucose on the determination of amino acids by GC 309(1984)375
LaBrosse, J.L. and Anderegg, R.J.
The mass spectrometer as a chlorine-selective chromatographic detector. I.

Description and evaluation of the technique 314(1984)83
LaBrosse, J.L. and Anderegg, R.J.
The mass spectrometer as a chlorine-selective chromatographic detector. II. Applications to chemical systems 314(1984)93
Lacey, E. and Brady, R.L.
Separation of colchicine and related hydrolysis and photodecomposition products by HPLC, using copper ion complexation 315(1984)233
Lachatre, G., see Nicot, G. 306(1984)279
Lackner, B., see Rembold, H. 323(1985)355
Lacko, R., see Koval, M. 325(1985)151
Lacroix, B.V., see Bovenkamp, J.W. 301(1984)492
Lacroix, C., Laine, G. and Goulle, J.P.
Determination of isoniazid and acetylisoniazid in plasma or serum by HPLC (in French) 307(1984)137
Lacroix, C., Levert, P., Laine, G. and Goulle, J.P.
Microdetermination of two antifibrinolytic agents (ε-aminocaproic acid and tranexamic acid) by LC and fluorimetric detection (in French) 309(1984)183
Ladányi, L., see Körtvélyessy, G. 303(1984)370
Lagerström, P.-O., Carlebom, P., Clarke, A.F. and Jack, D.B.
Determination of prenalterol in plasma and urine by LC with electrochemical detection 307(1984)230
Lagerström, P.-O. and Persson, B.-A.
Determination of omeprazole and metabolites in plasma and urine by LC 309(1984)347
Lagerström, P.-O., see Persson, B.-A. 316(1984)291
Laguzzi, B., see Carlucci, G. 311(1984)141
Lahr, L.M., see Spittler, T.D. 317(1984)527
Lai, C.-M., see Stampfli, H.F. 309(1984)203
Lai, S.-T. and Locke, D.C.
Pyrolysis-fluorescence spectroscopy, pyrolysis-MS and pyrolysis-LC of linear and cross-linked polystyrenes 314(1984)283
Lai, S.-T. and Sangermano, L.
Characterization of epoxy novolac resins using phenyl-bonded phase HPLC 321(1985)325
Lai, S.-T. and Sangermano, L.
Molecular weight determination of polychromophore polymers using HPLC and size-exclusion chromatography with UV detection 322(1985)336
Lai, S.-T., Sangermano, L., Chou, C.-K. and Locke, D.C.
Multi-mode LC of polystyrenes on Partisil PAC 324(1985)436
Lai, S.-T., Sangermano, L. and Locke, D.C.
Phenyl-bonded phase LC of styrene oligomers 312(1984)313
Laine, G., see Lacroix, C. 307(1984)137, 309(1984)183
Lakings, D.B. and Wozniak, J.M.
HPLC methods for the determination of cefpimizole in plasma and urine 308(1984)261
Lalka, D., see Kapil, R.P. 307(1984)305
Lam, S., Azumaya, H. and Karmen, A.
HPLC of amino acids in urine and cerebrospinal fluid 302(1984)21
Lam, S., see Bock, J.L. 308(1984)354
Lam, S., see Karmen, A. 302(1984)31
Lamb, J.H., see Bailey, E. 310(1984)199
Lamiable, D., Millart, H., Lemeignan, M. and Vistelle, R.
HPLC determination of 3,4-diaminopyridine in human plasma 309(1984)222
Landa, B.L., see Narasimhachari, N. 311(1984)424
Landis, J.B., see Asmus, P.A. 316(1984)461
Landon, M., see Poynter, D. 325(1985)333
Landy, J.S., see Dorsey, J.G. 316(1984)183
Lang, J.F., see Chan, K.Y. 306(1984)249
Lang, R.E., see Hermann, K. 312(1984)273
Langemann, H., see Honegger, C.G. 309(1984)53
Langenberg, J.P. and Tjaden, U.R.
Determination of (endogenous) vitamin K_1 in human plasma by reversed-phase HPLC using fluorometric detection after post-column electrochemical reduction. Comparison with UV, single and dual electrochemical detection 305(1984)61
Langendijk, P.N.J., Smith, P.C., Hasegawa, J. and Benet, L.Z.
Simultaneous determination of zomepirac and its major metabolite zomepirac glucuronide in human plasma and urine 307(1984)371
Langer, K., see Frank, H. 309(1984)261
Langer, K., see Schwarz, W. 310(1984)188
Langseth, W.
Speciation of alkyltin compounds by HPLC with a cyanopropyl-bonded silica column 315(1984)351
Lant, M.S., Martin, L.E. and Oxford, J.
Qualitative and quantitative analysis of ranitidine and its metabolites by HPLC-MS 323(1985)143
Lanz, E., see Wiegand, U.-W. 311(1984)218
Larcher, A.V., see Rowland, S.J. 312(1984)395
Larson, B.J., see Timm, D.C. 316(1984)343
Larsson, L., see De la Cochetière-Collinet, M.-F. 305(1984)178
Lartigue-Mattei, C., Chabard, J.L., Touzet, C., Bargnoux, H., Petit, J. and Berger, J.A.
Plasma cyclophosphamide assay by selective ion monitoring 310(1984)407
Laschet, J., see Bontemps, J. 311(1984)59
Lascu, I., see Porumb, T. 319(1985)218
Lasley, S.M., Michaelson, I.A., Greenland, R.D. and McGinnis, P.M.
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover

in discrete rat brain regions by LC with electrochemical detection 305(1984)27
Latini, R., see Kates, R.E. 309(1984)441
Lau, H.T., see Parkin, J.E. 314(1984)488
Lau, S.Y.M., Taneja, A.K. and Hodges, R.S. Effects of HPLC solvents and hydrophobic matrices on the secondary and quaternary matrices on the secondary and quaternary structure of a model protein. Reversed-phase and size-exclusion HPLC 317(1984)129
Lau, S.Y.M., see Taneja, A.K. 317(1984)1
Laudová, Z., see Šafařík, I. 303(1984)283
Laufer, R., Wormser, U., Selinger, Z., Chorev, M. and Gilon, C. Ion-exchange chromatographic assay of peptidases acting on the C-terminal hexapeptide sequence of substance P 301(1984)415
Lauren, D.R. and Agnew, M.P. Anomalies in the HPLC determination of diquat in water samples 303(1984)206
Laurensse, E., see Peters, G.J. 307(1984)464
Laurent, P., see Airaudo, C.B. 314(1984)349
Law, B. and Gill, R. Construction of a simple wall-jet electrochemical flow-cell for HPLC 325(1985)294
Law, B., Gill, R. and Moffat, A.C. HPLC retention data for 84 basic drugs of forensic interest on a silica column using an aqueous methanol eluent 301(1984)165
Lawley, C.W., see Bear, G.R. 302(1984)65
Lawrence, J.F., Iyengar, J.R. and Sun, W.F. GC patterns of some apple surface wax constituents 325(1985)299
Le Moing, J.P., see Vajta, S. 311(1984)329
Lea, A.R., see Johnston, M.A. 318(1985)362
Lecaillon, J.-B., Febvre, N. and Souppart, C. Influence of solute polarity in column-switching chromatography for the assay of drugs in plasma and urine 317(1984)493
Lecaillon, J.-B., Godbillon, J., Abadie, F. and Gosset, G. Determination of metoprolol and its α-hydroxylated metabolite in human plasma by HPLC 305(1984)411
Lechat, P., see Fels, J.-P. 308(1984)273
Lecomte-Joulin, V., see Guyon, F. 311(1984)160
Ledden, D.J., see Feldhoff, R.C. 311(1984)267
Lee, M.G. Determination of pseudomorphine in morphine injection by HPLC 312(1984)473
Lee, M.L., Kuei, J.C., Adams, N.W., Tarbet, B.J., Nishioka, M., Jones, B.A. and Bradshaw, J.S. Polarizable polysiloxane stationary phases for capillary column GC 302(1984)303
Lee, R.M., see McDowall, R.D. 305(1984)214
Lee, T.L. and Brooks, M.A. HPLC determination of amoxicillin in human plasma using a bonded-phase extraction 306(1984)429
Lee, T.T., Starratt, A.N. and Jevnikar, J.J. Separation of conjugates and oxidative metabolites of indole-3-acetic acid by HPLC 325(1985)340
Leeuw, I. de, see Vandewoude, M. 311(1984)176
Lefèbvre, M.-A., see Fourtillan, J.-B. 309(1984)391
Legrand, M., see Geoffroy, P. 315(1984)333
Lehmann, C., see Jamali, F. 310(1984)327
Lehtonen, P. Reversed-phase retention behaviour of Dns-amides as a function of eluent composition and molecular structure in ethanol-water and methanol-water 314(1984)141
Leidy, H.L., see Musson, D.G. 308(1984)251
Leitner, C., see Lindner, W. 316(1984)605
Leitold, A., see Vigh, G. 312(1984)345
Leitz, F., see Alton, K.B. 311(1984)319
Lejczak, B., Kafarski, P. and Mastalerz, P. Chromatographic assignment of the configuration of 1-aminoalkanephosphonic acids 324(1985)455
Lejeune, P.-J., see Marriq, C. 323(1985)395
Lemberkovics, É. and Verzár-Petri, G. GC characterization of frequently occuring sesquiterpenes in essential oils 318(1985)125
Lemeignan, M., see Lamiable, D. 309(1984)222
Lemmens, A.A.G., Reijenga, J.C., Everaerts, F.M., Janssen, R.T.P., Hulsman, J.A.R.J. and Meijers, C.A.M. Isotachophoresis as a candidate reference method in analytical chemistry. Determination of sodium in serum 320(1985)193
Lemmens, A.A.G., see Reijenga, J.C. 320(1985)67, 320(1985)245
Lemmer, B., Ohm, T. and Winkler, H. Determination of the beta-adrenoceptor blocking drug sotalol in plasma and tissues of the rat by HPLC with UV detection 309(1984)187
Lemmer, B., see Winkler, H. 309(1984)193
Lenczyk, M., see Schmitz, G. 307(1984)81
Lenten, F.J. van, see Buchta, R.C. 325(1985)456
Leone, L., Monteleone, M., Gabutti, V. and Amione, C. Reversed-phase HPLC of human haemoglobin chains 321(1985)407
Leonhardt, B.A. and DeVilbiss, E.D. Separation and double-bond determination on nanogram quantities of aliphatic monounsaturated alcohols, aldehydes and carboxylic acid methyl esters 322(1985)484
Lepom, P. and Robowsky, K.-D. Determination of the alkaloid perloline in grasses by HPLC 322(1985)261
Lepri, L., Desideri, P.G. and Coas, W. HPTLC of fluorescent whitening agents and their identification in detergents 322(1985)363
Lepri, L., see Desideri, P.G. 322(1985)107
Lercker, G., see Caboni, M.F. 315(1984)223
Lerma, J., see Herranz, A.S. 309(1984)139
Leroy, P. and Nicolas, A. Determination of secondary amino drugs as their metal dithiocarbamate complexes by

reversed-phase HPLC with electrochemical detection 317(1984)513
Lertora, J.J.L., see Rege, A.B. 309(1984)397
Lethbridge, J.W. and Reeve, R.N.
HPLC in the study of some reactions of chromium(III) complexes 314(1984)387
Levert, P., see Lacroix, C. 309(1984)183
Levin, J.-O., see Andersson, K. 312(1984)482
Levine, S.P., see Strang, C. 314(1984)482
Levsen, K., Wagner-Redeker, W., Schäfer, K.H. and Dobberstein, P.
On-line LC-MS analysis of non-ionic surfactants 323(1985)135
Levsen, K., see Schmelzeisen-Redeker, G. 323(1985)127
Levy, M., see Katz, E.Z. 305(1984)477
Lewis, H.F., see Dulitzky, M. 317(1984)403
Lewis, R.V., see Sadler, A.J. 317(1984)93
Lewis, S. and Wilson, I.D.
Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133
Leyendecker, D., Schmitz, F.P. and Klesper, E.
Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19
Leyendecker, D., Schmitz, F.P., Leyendecker, D. and Klesper, E.
Chromatography with sub- and supercritical eluents: influence of temperature, pressure and flow-rate on the behaviour of lower alkanes 321(1985)273
Leyendecker, D., see Leyendecker, D. 321(1985)273
Leyva, A., see Peters, G.J. 307(1984)464
Liebich, H.M. and Först, C.
Hydroxycarboxylic and oxocarboxylic acids in urine: products from branched-chain amino acid degradation and from ketogenesis 309(1984)225
Liggins, G.C., see Skinner, S.J.M. 308(1984)111
Ligny, C.L. de, see Brandts, P.M. 322(1985)399
Ligny, C.L. de, see De Ligny, C.L. 301(1984)311
Lim, C.K. and Peters, T.J.
Ammonium acetate: a general purpose buffer for clinical applications of HPLC 316(1984)397
Lim, C.K., Rideout, J.M. and Peters, T.J.
HPLC of dicarboxylic porphyrins and metalloporphyrins: retention behaviour and biomedical applications 317(1984)333
Lim, C.K., see Bottiglieri, T. 311(1984)354
Lim, P.C., see Hicks, K.B. 319(1985)159
Limentani, G.B. and Uden, P.C.
High-resolution GC and LC of organoarsenic compounds 325(1985)53
Limero, T., see Hernandez-Gil, N. 312(1984)31
Lin, J.-L., see Dorsey, J.G. 316(1984)183
Lind, M.A., see Korhonen, I.O.O. 322(1985)71, 322(1985)83, 322(1985)97, 323(1985)331, 324(1985)113, 325(1985)433
Lindberg, R.L.P. and Pihlajamäki, K.K.
HPLC determination of bupivacaine in human serum 309(1984)369
Lindblad, W.J. and Diegelmann, R.F.
Chromatographic separation of preparative quantities of the stereoisomers of *trans*-2,3-*cis*-3,4-dihydroxy-L-proline 315(1984)447
Lindfors, A., see Eksborg, S. 306(1984)446
Lindner, W., Leitner, C. and Uray, G.
LC separation of enantiomeric alkanolamines via diastereomeric tartaric acid monoesters 316(1984)605
Lindquist, J., see Isaksson, K. 324(1985)333
Lineback, D.R., see Kobayashi, S. 319(1985)205
Linton, R.W., see Miller, M.L. 319(1985)9
Lion, C., see Chrétien, J.R. 324(1985)355
Lippi, A., see Grossi, G. 309(1984)214
Lisman, J.J.W., see Beem, E.P. 323(1985)439
Lissitzky, S., see Marriq, C. 323(1985)395
Litowitz, H., Olanoff, L. and Hoppel, C.L.
Determination of ibuprofen in human plasma by HPLC 311(1984)443
Little, C.J. and Stahel, O.
Role of column switching in semi-preparative LC. Isolation of the sweetener stevioside 316(1984)105
Lizotte, R., see Ryan, J.J. 303(1984)351
Lo, J.-G. and Nieh, J.-C.
Radio-GC of tritiated compounds 322(1985)332
Loche, S., Porcelli, F., Rosen, M., Feffer, M., Stoner, E. and New, M.I.
Clinical application of the rapid HPLC determination of serum cortisol 317(1984)377
Locke, D.C., see Lai, S.-T. 312(1984)313, 314(1984)283, 324(1985)436
Lockwood, F.E., see Sprissler, B. 319(1985)222
Lodge, B.A. and Vincent, A.
Analysis of flumethasone pivalate formulations by HPLC 301(1984)477
Lodge, B.A., see Carignan, G. 301(1984)292, 315(1984)470
Lombet, J., see Bontemps, J. 307(1984)283
Long, A.R., see Chism, G.W. III 317(1984)263
Loo, J.C.K., Jordan, N. and Ngoc, A.H.
Application of a high-performance gel permeation LC procedure to the determination of binding of prednisolone to high-affinity binding sites in human serum 305(1984)194
Looker, M.R., see Yeung, P.K.F. 303(1984)412
López-Garzón, F.J., see Domingo-García, M. 324(1985)19
López-Garzón, R., see Domingo-García, M. 324(1985)19
Lorenzin, F., see Castagnola, M. 307(1984)91
Louis, W.J., see Drummer, O.H. 305(1984)83
Lovering, E.G. and Curran, N.M.
HPLC determination of cimetidine and related compounds (raw material and tablets) 319(1985)235
Low, T.L.K. and Mercer, R.C.
Isolation and structural studies of porcine,

ovine and murine thymosin β_4 by HPLC 301(1984)221
Lowe, J.R., see Dixon, J.S. 310(1984)455
Lozier, J.N., see Takahashi, N. 317(1984)11
Lugari, M.T., Maggiali, C.A. and Morini, G. HPLC of phytotoxic substances. II. Behaviour of imidazopyrimidines 315(1984)384
Luker, R.G., see Weaver, K.M. 301(1984)288
Lüllmann, C., Genieser, H.-G. and Jastorff, B. Structural investigations on reversed-phase silicas. Determination of ligand functionality 323(1985)273
Luna, M., see Felici, M. 320(1985)435
Lundanes, E. and Greibrokk, T. Reversed-phase separation of alkanes in high-boiling fractions of crude oil 322(1985)347
Lundh, T., see Dalene, M. 322(1985)169
Lundmo, P.I. and Sunde, A. Rapid analysis of C_{19}-steroid metabolism by HPLC and in-line monitoring of radioactivity 308(1984)289
Lundström, K., see Isaksson, K. 324(1985)333
Lundy, A.M., see Bellon, G. 311(1984)405
Lunte, C.E. and Kissinger, P.T. Determination of quinonoid dihydrobiopterin by LC and electrochemical detection 317(1984)407
Lunte, S.M. and Kissinger, P.T. Use of LC with dual-electrode electrochemical detection in the investigation of glutathione oxidation during benzene metabolism 317(1984)579
Luoma, G.A. Fingerprinting epoxy resin systems using two-dimensional reversed-phase TLC 318(1985)75
Lurie, I.S. and Allen, A.C. Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427
Luzzi, F.A., Wenger, T.L., Klinger, J.K., Barchowsky, A. and Strauss, H.C. Simultaneous determination of lidocaine and its metabolites in plasma and myocardium 311(1984)291
Lynn, R.K., see Boppana, V.K. 317(1984)463

M

McAllister, C.B., see Wedlund, P.J. 307(1984)121
McAllister, T., see Early, R.J. 310(1984)1
McBurney, A. and George, S. HPLC of haloperidol in serum at the concentrations achieved during chronic therapy 308(1984)387
McCall, J.T., see Reece, P.A. 306(1984)417
McCauley, F.A., see Wilson, T.W. 306(1984)351
McCormick, M.J. and Dixon, L.M. Determination of sulphite in fixers and photographic effluents by ion chromatography 322(1985)478
McCown, M., see Cheng, W. 318(1985)173
McCulloch, J., see Gustine, D.L. 316(1984)407
McCune, R.A., see Gehrke, C.W. 301(1984)199
McDermott, J.R., see Smith, A.I. 306(1984)99
McDowall, M.A., see Schmelzeisen-Redeker, G. 323(1985)127
McDowall, R.D., Murkitt, G.S. and Lee, R.M. Improved assay procedure for oxmetidine and its metabolites in plasma, urine and bile samples 305(1984)214
McDowall, R.D., Murkitt, G.S. and Walford, J.A. HPLC determination of oxmetidine in human plasma: comparison of liquid-liquid and liquid-solid extraction techniques 317(1984)475
McGaugh, J.L., see Ishikawa, K. 306(1984)394
MacGee, J., see Allen, K.G. 309(1984)33
McGinnis, P.M., see Lasley, S.M. 305(1984)27
McGoldrick, K., see Ryan, P.J. 320(1985)421
McGregor, J.L., see Clezardin, P. 319(1985)67
McGregor, W.C., see Roy, S.K. 303(1984)225
Maciel, G.E., see Miller, M.L. 319(1985)9
Maciel, G.E., see Rudzinski, W.E. 323(1985)281
McKay, G., see Roy, S.D. 310(1984)307
McKay, L., Bradberry, C. and Oke, A. Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167
MacKenzie, S.L. and Tenaschuk, D. Analysis of γ-carboxyglutamic acid by GLC 319(1985)404
MacKenzie, S.L. and Tenaschuk, D. GLC assay for asparagine and glutamine 322(1985)228
McKinnon, A., see Jane, I. 323(1985)191
McLaughlin, L.W., see Bischoff, R. 317(1984)251
McLean, S., see Davies, N.W. 310(1984)179
McNally, M.E., see Alexander, J.N. IV 318(1985)289
McNicholas, P.A., Batley, M. and Redmond, J.W. Reversed-phase HPLC of hydrazones of 3-deoxy-D-*manno*-oct-2-ulosonic acid and neuraminic acids 315(1984)451
Macrae, R., see Aries, R.E. 319(1985)285
Macrae, R., see Robinson, J.L. 303(1984)386, 324(1985)35
Madajová, V., Kaniansky, D., Čižmárová, E. and Hudec, M. Isotachophoretic determination of lysine in fermentation broths 320(1985)131
Madden, B.G., see Haken, J.K. 325(1985)61
Madej, A., see Hallin, P. 319(1985)195
Maes, R.A.A., see Drost, R.H. 310(1984)193

Maes, R.A.A., see Meering, P.G. 310(1984)159
Maggiali, C.A., see Lugari, M.T. 315(1984)384
Mailhe, L., see Bauer, M. 315(1984)457
Maingault, P., Hoinard, C., Crouzat-Reynes, G. and Guilloteau, D.
Immobilization of ligands for affinity chromatography. Coupling on a spacer arm gel with N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline as condensation agent: study of coupling conditions by means of a radioimmunologic method 314(1984)445
Maisano, F., Belew, M. and Porath, J.
Synthesis of new hydrophobic adsorbents based on homologous series of uncharged alkyl sulphide agarose derivatives 321(1985)305
Maitra, A.M., see Dilli, S. 318(1985)350
Majumdar, S.G., see Ray, B. 315(1984)401
Makino, K., see Wada, H. 320(1985)369
Makita, M., see Kataoka, H. 306(1984)61
Makita, M., see Yamamoto, S. 303(1984)99
Makuch, B., see Gazda, K. 320(1985)397
Malcolme-Lawes, D.J., see Blatchford, C. 321(1985)227
Malgras, A., see Bellon, G. 311(1984)405
Malhotra, R.C., see Tripathi, D.N. 315(1984)417
Malikin, G., see Karmen, A. 302(1984)31
Mallet, A.I., Cunningham, F.M. and Daniel, R.
Rapid isocratic HPLC purification of platelet activating factor (PAF) and *lyso*-PAF from human skin 309(1984)160
Mallet, A.I., see Woollard, P.M. 306(1984)1
Manach, M., see Clezardin, P. 319(1985)67
Manca, M., see Cossu, G. 307(1984)103
Mandl, I., see Keller, S. 305(1984)461
Mangani, F., see Bruner, F. 302(1984)167
Mangia, A., see Casoli, A. 303(1984)404
Manning, D.L., see Maskarinec, M.P. 302(1984)51
Månsson, J.-E., Rosengren, B. and Svennerholm, L.
Separation of gangliosides by anion-exchange chromatography on Mono Q™ 322(1985)465
Maoka, T., Komori, T. and Matsuno, T.
Direct diastereomeric resolution of carotenoids. I. 3-Hydroxy-4-oxo β-end group 318(1985)122
Marafioti, R.A., see Spittler, T.D. 317(1984)527
Marciniak, W. and Witkiewicz, Z.
Distribution of the liquid crystal stationary phase on the surface of diatomite supports 324(1985)299
Marciniak, W. and Witkiewicz, Z.
Phase transitions of 4-cyano-4'-*n*-alkoxyformyloxyazobenzenes on silanized supports 324(1985)309
Marcucci, F., Colombo, L., De Ponte, G., Cani, D., Cobelli, L., Frigerio, A. and Mussini, E.
GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132
Marcucci, F., see Mussini, E. 305(1984)450
Marenchic, I.G., see Taylor, R.F. 317(1984)193
Margolis, S.A. and Dizdaroglu, M.
Separation and purification of diastereomers of angiotensin I by weak anion-exchange HPLC 322(1985)117
Maris, F.A., Geerdink, R.B., Frei, R.W. and Brinkman, U.A.T.
On-line trace enrichment for improved sensitivity in LC with direct liquid introduction MS detection 323(1985)113
Markey, S.P., see Goldstein, D.S. 311(1984)148
Markides, K., Blomberg, L., Hoffmann, S., Buijten, J. and Wännman, T.
Cyanosilicones as stationary phases in GC. III. Synthesis, characterization and evaluation 302(1984)319
Markides, K., see Buijten, J. 301(1984)265
Markides, K., see Hoffmann, S. 302(1984)95
Markovič, O. and Slezárik, A.
Product analysis of the action of *Trichoderma reesei* polygalacturonase by TLC 312(1984)492
Marks, E.S., see Goldstein, D.S. 311(1984)148
Marks, V., see Burton, N.K. 309(1984)409
Marlet, B. and Pommier, C.
GC analysis of the extraction solvent used in nuclear fuel reprocessing plants 325(1985)75
Marriott, P.J., Gill, J.P., Evershed, R.P., Hein, C.S. and Eglinton, G.
Computerised GC-MS analysis of complex mixtures of alkyl porphyrins 301(1984)107
Marriq, C., Lejeune, P.-J., Rolland, M. and Lissitzky, S.
Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395
Marsden, N.V.B.
Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.*-butanol 319(1985)247
Marsden, N.V.B., see Haglund, Å.C. 301(1984)47, 318(1985)57
Marsden, N.V.B., see Haglund, Å.C. 301(1984)365
Marsh, P.G., see Pike, A.W. 306(1984)39
Martelli, E., see Grossi, G. 309(1984)214
Martelli, P., Bovalini, L., Ferri, S. and Franchi, G.G.
Rapid separation and quantitative determination of khellin and visnagin in *Ammi visnaga* (L.) Lam. fruits by HPLC 301(1984)297
Martens, M.A., see De Beer, J.O. 307(1984)475
Mårtensson, J., see Kågedal, B. 311(1984)170
Martin, L.E., see Lant, M.S. 323(1985)143
Martin, R.A. Jr., see Hurst, W.J. 318(1985)408
Martínez, E., see Tusell, J.M. 306(1984)338
Martinez, J.L. Jr., see Ishikawa, K. 306(1984)394
Maruta, K., see Ito, S. 311(1984)154
Maruyama, Y., see Ikarashi, Y. 322(1985)191
Marzo, A., Treffner, E., Neggiani, P.P. and Staibano, G.

GLC evaluation of lofemizole in biological samples for pharmacokinetic investigations. Comparison of two analytical methods 310(1984)51
Masaki, H., see Tanaka, A. 306(1984)51
Mascher, H., Nitsche, V. and Schütz, H.
Separation, isolation and identification of optical isomers of 1,4-benzodiazepine glucuronides from biological fluids by reversed-phase HPLC 306(1984)231
Mashimo, I., see Kawamoto, T. 305(1984)373
Maskarinec, M.P., Manning, D.L., Harvey, R.W., Griest, W.H. and Tomkins, B.A.
Determination of munitions components in water by resin adsorption and HPLC electrochemical detection 302(1984)51
Mason, P.F., see Snook, M.E. 324(1985)141
Massart, D.L., see Hoogewijs, G. 309(1984)329
Mastalerz, P., see Lejczak, B. 324(1985)455
Masuda, S., see Baba, S. 305(1984)119
Matas, R., see Méndez, E. 323(1985)373
Matheson, T.W., see Harvey, T.G. 319(1985)230
Mathews, R.A., see Said, H.M. 324(1985)65
Mathiasson, L., see Audunsson, G. 315(1984)299
Mathiasson, L., see Dalene, M. 322(1985)169
Mathis, R., Hubert, P. and Dellacherie, E.
Sepharose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 319(1985)43
Matisová, E., Krupčík, J., Čellár, P. and Garaj, J.
Quantitative analysis of hydrocarbons in gasolines by capillary GLC. I. Isothermal analysis 303(1984)151
Matlis, R. and Greenblatt, D.J.
Rapid HPLC analysis of oxaprozin, a non-steroidal anti-inflammatory agent 310(1984)445
Maton, P.N., see Selden, C. 305(1984)43
Matoušek, P., see Horna, A. 325(1985)367
Matsubara, K., see Kato, Y. 320(1985)440
Matsuda, A., see Okano, Y. 310(1984)251
Matsuhashi, S., Mukai, T. and Hori, K.
Determination of protein spots separated by two-dimensional polyacrylamide gel electrophoresis 319(1985)79
Matsumoto, A., see Kubo, I. 321(1985)246
Matsumoto, H., see Oka, H. 325(1985)265
Matsumoto, U., see Ikeda, M. 305(1984)261
Matsuno, T., see Maoka, T. 318(1985)122
Matsushita, S.
Simultaneous determination of anions and metal cations by single-column ion chromatography with ethylenediaminetetraacetate as eluent and conductivity and UV detection 312(1984)327
Maurer, H. and Pfleger, K.
Screening procedure for detection of antidepressants and their metabolites in urine using a computerized GC-MS technique 305(1984)309
Maurer, H. and Pfleger, K.
Screening procedure for detection of phenothiazine and analogous neuroleptics and their metabolites in urine using a computerized GC-MS technique 306(1984)125
Maurizis, J.C., see Bernard, D. 308(1984)322
Maurs, M., Ducrocq, C., Righini-Tapie, A. and Azerad, R.
GC resolution of substituted glutamic acid enantiomers 325(1985)444
Mays, D.C., Sharp, D.E., Beach, C.A., Kershaw, R.A., Bianchine, J.R. and Gerber, N.
Improved method for the determination of aspirin and its metabolites in biological fluids by HPLC: applications to human and animal studies 311(1984)301
Meagher, M.M., see Nikolov, Z.L. 319(1985)51, 321(1985)393
Meatherall, R.C. and Guay, D.R.P.
Isothermal GC analysis of diphenhydramine after direct injection onto a fused-silica capillary column 307(1984)295
Meek, J.L. and Eva, C.
Enzymes adsorbed on an ion exchanger as a post-column reactor: application to acetylcholine measurement 317(1984)343
Meering, P.G., Baumann, R.A., Zijp, J.J. and Maes, R.A.A.
Determination of misonidazole and desmethylmisonidazole in plasma by HPLC with reductive electrochemical detection 310(1984)159
Meffin, P.J., see Veenendaal, J.R. 307(1984)432
Meijerhoff, G., see Van der Greef, J. 323(1985)81
Meijerink, N.L.J., see Glöckner, G. 317(1984)615
Meijers, C.A.M., see Lemmens, A.A.G. 320(1985)193
Meister, W., see Dell, D. 317(1984)483
Melander, W.R., Chen, B.-K. and Horváth, C.
Mobile phase effects in reversed-phase chromatography. VII. Dependence of retention on mobile phase composition and column temperature 318(1985)1
Melander, W.R., Corradini, D. and Horváth, C.
Salt-mediated retention of proteins in hydrophobic interaction chromatography. Application of solvophobic theory 317(1984)67
Melander, W.R., see Chang, J.P. 318(1985)11
Melander, W.R., see Huang, J.-X. 316(1984)151
Mellbin, G. and Smith, B.E.F.
Trace determination of aliphatic amines using HPLC with chemiluminescence excitation and photon counting 312(1984)203
Meltzer, N.M. and Sternson, L.A.
Analysis of riboxamide in urine by HPLC 307(1984)216
Men, L.-C., Na, H. and Rogers, L.B.
Solvent selectivity studies using isomers of polystyrene 314(1984)55

Méndez, E., Matas, R. and Soriano, F.
Complete automatization of peptide maps by reversed-phase LC using *o*-phthalaldehyde pre-column derivatization 323(1985)373
Menendez, V., see Fernández-Sánchez, E. 312(1984)69
Menge, G. and Dubois, J.P.
Determination of aminoglutethimide and N-acetylaminoglutethimide in human plasma by HPLC 310(1984)431
Mengel, I., see Krüger, R. 311(1984)109
Mentasti, E., Gennaro, M.C., Sarzanini, C., Baiocchi, C. and Savigliano, M.
Derivatization, identification and separation of carboxylic acids in wines and beverages by HPLC 322(1985)177
Mercer, R.C., see Low, T.L.K. 301(1984)221
Merigan, T.C., see Smith, G.A. 307(1984)410
Merkel, K.E., see Polta, J.A. 324(1985)407
Merle, L., see Nicot, G. 306(1984)279
Merrill, E.W., see Chaumont, P.R. 303(1984)177
Mesrob, B., see Arsov, A. 322(1985)267
Metelitsa, V.I., see Piotrovskii, V.K. 309(1984)421
Meyer, V.R.
Preparative HPLC as an aid in organic synthesis 316(1984)113
Micallef, S., see Pace-Asciak, C.R. 310(1984)233
Micanovic, R., see Sadler, A.J. 317(1984)93
Miceli, J.N., see Aravind, M.K. 308(1984)350
Miceli, J.N., see Bowman, D.B. 309(1984)209
Micera, G., see Deiana, S. 320(1985)450
Michaelson, I.A., see Lasley, S.M. 305(1984)27
Middaugh, C.R., see Sadler, A.J. 317(1984)93
Midha, K.K., see Roy, S.D. 310(1984)307
Midha, K.K., see Yeung, P.K.F. 303(1984)412
Midskov, C.
Rapid GC determination of pyrimethamine in human plasma and urine 306(1984)388
Midskov, C.
HPLC assay of pyrimethamine, sulfadoxine and its N^4-acetyl metabolite in serum and urine after ingestion of Suldox 308(1984)217
Mihaly, G.W., see Coleman, M.D. 308(1984)363
Mihara, S. and Enomoto, N.
Calculation of retention indices of pyrazines on the basis of molecular structure 324(1985)428
Mihellyes, S., see Gübitz, G. 314(1984)462
Mijnsbergen, C.W., see Barends, D.M. 322(1985)321
Mikasa, H., Arata, J. and Kodama, H.
Measurement of prolidase activity in erythrocytes using isotachophoresis 310(1984)401
Mikasa, H., Sasaki, K., Kodama, H., Arata, J. and Ikeda, M.
Isotachophoretic analysis of iminopeptides in the urine of patients with iminopeptiduria 305(1984)204
Mikasa, H., see Kodama, H. 311(1984)183
Mikaya, A.I., see Zaikin, V.G. 301(1984)77
Mikeš, O., Štrop, P., Hostomská, Z., Smrž, M., Slováková, S. and Čoupek, J.
Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93
Mikešová, M., see Vejrosta, J. 324(1985)269
Milborrow, B.V., see Devine, P.L. 325(1985)323
Millart, H., see Lamiable, D. 309(1984)222
Miller, D.D., see Abernethy, D.R. 307(1984)194
Miller, G.C., see Tamma, R.V. 322(1985)236
Miller, K.J., Gal, J. and Ames, M.M.
HPLC resolution of enantiomers of 1-phenyl-2-aminopropanes (amphetamines) with four chiral reagents 307(1984)335
Miller, M.L., Linton, R.W., Maciel, G.E. and Hawkins, B.L.
Characterization of silanol reactivity and acidity on octadecyl-bonded chromatographic supports by ^{29}Si solid-state nuclear magnetic resonance spectroscopy and surface titration 319(1985)9
Miller, N.T., Feibush, B. and Karger, B.L.
Wide-pore silica-based ether-bonded phases for separation of proteins by high-performance hydrophobic-interaction and size-exclusion chromatography 316(1984)519
Miller, S., see Dilli, S. 324(1985)75
Millet, J.L., see Guillemin, C.L. 301(1984)11
Minárik, M., see Nondek, L. 324(1985)261
Minegishi, A. and Ishizaki, T.
Rapid and simple method for the simultaneous determination of 3,4-dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by HPLC with electrochemical detection 308(1984)55
Minegishi, A. and Ishizaki, T.
Determination of free 3-methoxy-4-hydroxyphenylglycol with several other monoamine metabolites in plasma by HPLC with amperometric detection 311(1984)51
Mink, C.J.K., see Wielders, J.P.M. 310(1984)379
Minor, J.M., see Glajch, J.L. 318(1985)23
Mistryukov, E.A. and Korshevetz, I.K.
"Inverse" injection system for GLC with capillary columns 322(1985)206
Mitchell, F., see D'Agostino, G. 305(1984)13
Mitchell, J.D., see Ersser, R.S. 307(1984)393
Mitchell, S.D., see Albert, L.S.R. 312(1984)357
Miwa, H., Hiyama, C. and Yamamoto, M.
HPLC of short- and long-chain fatty acids as 2-nitrophenylhydrazides 321(1985)165
Miyaguchi, K., Honda, K. and Imai, K.
Sub-picomol chemiluminescence detection of Dns-amino acids separated by HPLC with gradient elution 303(1984)173
Miyaguchi, K., Honda, K. and Imai, K.
Microbore HPLC and chemiluminescence detection of Dns-amino acids 316(1984)501
Miyamoto, E., see Shoji, S. 319(1985)359
Miyata, T., see Okano, Y. 310(1984)251
Miyazaki, K., Kohri, N., Arita, T., Shimono, H., Katoh, K., Nomura, A. and Yasuda, H.

HPLC determination of nifedipine in plasma 310(1984)219
Miyazaki, M., Hayakawa, K. and Choi, S.-G. Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443
Miyazaki, T., see Kai, M. 311(1984)257, 322(1985)473
Miyazaki, T., see Yoshioka, M. 309(1984)63
Mizutani, T. and Tachibana, Y. Rapid fractionation of tRNA on benzoylated DEAE-porous glass 324(1985)480
Moats, W.A. Determination of penicillin G anmd cloxacillin residues in beef and pork tissue by HPLC 317(1984)311
Mocharla, R., see Jalal, M.A.F. 301(1984)247
Möckel, H.J. Retention of sulphur and sulphur organics in reversed-phase LC 317(1984)589
Moegel, H-J., Kraus, G. and Novak, M. Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29
Moerloose, P. de, see De Croo, F. 325(1985)395
Moerloose, P. de, see De Spiegeleer, B. 315(1984)481
Moerloose, P. de, see Van den Eeckhout, E. 318(1985)343
Moffat, A.C., see Gill, R. 301(1984)155
Moffat, A.C., see Law, B. 301(1984)165
Mohamed, G.B., Nazareth, A., Hayes, M.J., Giese, R.W. and Vouros, P. GC-MS characteristics of methylated perfluoroacyl derivatives of cytosine and 5-methylcytosine 314(1984)211
Mohiuddin, G., see Shiao, S.-Y. 303(1984)13
Möhrke, W., see Spahn, H. 310(1984)167
Möhrke, W., see Weber, H. 307(1984)145
Moing, J.P. le, see Vajta, S. 311(1984)329
Molle, L., see Duez, P. 303(1984)391
Monteleone, M., see Leone, L. 321(1985)407
Montgomery, T.L., see Rudzinski, W.E. 323(1985)281
Moo-Penn, W.F. and Jue, D.L. Separation of hemoglobin variants by ion-exchange chromatography on monobead resins 318(1985)325
Moodie, I.M., see Labadarios, D. 309(1984)375, 310(1984)223
Moore, R.M. and Walters, R.R. Protein separations on reversed-phase HPLC minicolumns 317(1984)119
More, N.S., see Phillips, T.M. 317(1984)173
Moreno-Castilla, C., see Domingo-García, M. 324(1985)19
Morgan, S.L., see Kiel, J.S. 320(1985)313
Mori, M., see Dohji, T. 311(1984)249
Mori, S. and Suzuki, M. Semi-micro size-exclusion chromatography of polymers. Calibration of columns and comparison with conventional columns 320(1985)343
Mori, Y., see Baba, S. 307(1984)1
Morimoto, S., see Imaizumi, N. 308(1984)295
Morini, G., see Lugari, M.T. 315(1984)384
Morio, M., see Hirokawa, T. 305(1984)429
Morishima, S., see Kurosawa, N. 305(1984)485
Morishita, H., Iwahashi, H., Osaka, N. and Kido, R. Chromatographic separation and identification of naturally occurring chlorogenic acids by ^1H nuclear magnetic resonance spectroscopy and MS 315(1984)253
Morita, K., see Funazo, K. 319(1985)143
Moro, F., see Bettero, A. 310(1984)390
Morreal, C.E. and Clift, R.G. Identification of phenols of 7,12-dimethylbenz[a]anthracene by conversion into their methyl ethers 319(1985)153
Morrison, R.A. and Fung, H.-L. Determination of the partitioning, stability, and metabolite formation of isosorbide dinitrate in human and rat blood using and improved GLC assay 308(1984)153
Morrissey, E.G., see Berridge, J.C. 316(1984)69
Morselli, P.L., see Padovani, P. 308(1984)229
Mosher, R.A., Thormann, W. and Bier, M. Computer aided analysis of electric field gradients within isotachophoretic boundaries between weak electrolytes 320(1985)23
Motohashi, N., see Kamata, K. 319(1985)331
Mourey, T.H., Noh, I. and Yu, H. High-performance liquid-solid adsorption chromatography of spiropyran-end-labelled polystyrenes 303(1984)361
Mourey, T.H., see Oppenheimer, L.E. 323(1985)297
Moynihan, C., see Pike, A.W. 306(1984)39
Mudřík, Z., see Vrátný, P. 322(1985)352
Muir, K.T., see Hidalgo, I.J. 305(1984)222
Mukai, T., see Matsuhashi, S. 319(1985)79
Müller, K., see Blasius, E. 313(1984)161
Mullin, B.R., Poore, C.M.B. and Rupp, B.H. Quantitation of gangliosides in the picomolar range 305(1984)512
Mumby, S.J., Cotts, P.M. and Russell, T.P. Preparative fractionation of fluorescent-dye-labeled poly(ethylene oxide) by aqueous size-exclusion chromatography 319(1985)241
Munk, P., see Card, T.W. 301(1984)261
Murkitt, G.S., see McDowall, R.D. 305(1984)214, 317(1984)475
Murotsu, T., see Kato, Y. 320(1985)440
Murphy, R.C., see Strife, R.J. 305(1984)3
Murray, G.R., Calvey, T.N., Williams, N.E. and Chan, K. Quantitative capillary column GC method for the determination of glycopyrronium in human plasma 308(1984)143
Murrill, E., see Ray, G. 317(1984)455
Murrill, E., see Timmons, L. 314(1984)476

Musacchia, X.J., see Feldhoff, R.C. 311(1984)267
Muschik, G.M., see Issaq, H.J. 302(1984)159
Muselasová, I., see Vacík, J. 320(1985)199, 320(1985)233
Mushtaq, M., see Yang, S.K. 316(1984)569
Musial, S.P., Freeman, C.J. and Sinsheimer, J.E.
Mitotane (*o,p'*-DDD) emulsion and tablet analysis by HPLC 319(1985)467
Mussini, E., Colombo, L., De Ponte, G. and Marcucci, F.
Determination of creatine in body fluids and muscle 305(1984)450
Mussini, E., see Marcucci, F. 309(1984)132
Musson, D.G., Vincek, W.C., Dobrinska, M.R., Vickers, S. and Leidy, H.L.
Sensitive HPLC assay using electrochemical detection for a novel prodrug ester of methyldopa, pivaloyloxyethyl 3-(3,4-dihydroxyphenyl)-2-methylalaninate, in plasma and urine 308(1984)251
Musson, D.G., see Gravallese, D.A. 310(1984)71
Mutschler, E., see Prinoth, M. 305(1984)508
Mutschler, E., see Spahn, H. 310(1984)167
Mutschler, E., see Weber, H. 307(1984)145
Muusze, R.G., see De Jong, J. 319(1985)23

N

Na, H., see Men, L.-C. 314(1984)55
Naber, K.G., see Kees, F. 305(1984)363
Nagasawa, H.T., see Demaster, E.G. 308(1984)83
Nagashima, H. and Nakamura, K.-I.
Microdetermination of iodine in organic compounds by reversed-phase HPLC 324(1985)498
Nagata, J., see Okano, Y. 310(1984)251
Nagatsu, T., see Oka, K. 308(1984)43
Nagy, J., Várady, L., Kalász, H. and Knoll, J.
Molecular weight measurement of human satietin 317(1984)165
Nagy, J., see Kalász, H. 316(1984)95
Naito, Y., see Imamura, T. 305(1984)456
Nakae, T., see Ikigai, H. 322(1985)212
Nakagawa, F., see Uchino, K. 308(1984)241
Nakajima, K., Kojima, M., Fujita, M. and Fujita, J.
Chromatographic separation of the diastereomers of the bis(2,4-pentanedionato)(2-(methylseleno)ethylamine)cobalt(III) ion on a column of sulphoethyl-Toyopearl and determination of the rate of inversion at the selenium atom 301(1984)241
Nakamoto, M., see Hattori, Y. 315(1984)341
Nakamura, A., see Yoshioka, M. 309(1984)63
Nakamura, H., see Iwatani, A. 309(1984)145
Nakamura, K., see Kato, Y. 318(1985)358
Nakamura, K.-I., see Nagashima, H. 324(1985)498
Nakamura, M., Kondo, K., Nishioka, R. and Kawai, S.
Improved procedure for the HPLC determination of valproic acid in serum as its phenacyl ester 310(1984)450
Nakamura, M., see Ishida, J. 305(1984)381
Nakano, Y., Yamaguchi, M., Tsuruta, Y., Ohkura, Y., Aoyama, T. and Horioka, M.
Assay for histidine decarboxylase in rat stomach and brain by HPLC with fluorescence detection 311(1984)390
Nakata, T., see Kodama, K. 311(1984)369
Nakatani, K., see Yoneda, H. 319(1985)35
Nakazawa, H., Riggs, C.E. Jr., Egorin, M.J., Redwood, S.M., Bachur, N.R., Bhatnagar, R. and Ito, Y.
Continuous extraction of urinary anthracycline antitumor antibiotics with the horizontal flow-through coil planet centrifuge 307(1984)323
Nambara, T., see Shimada, K. 307(1984)23
Narang, P.K., Tourville, J.F., Chatterji, D.C. and Gallelli, J.F.
Quantitation of flupirtine and its active acetylated metabolite by reversed-phase HPLC using fluorometric detection 305(1984)135
Narasimhachari, N., Dorey, R.C., Landa, B.L. and Friedel, R.O.
Improved HPLC method for the quantitation of *cis*-thiothixene in plasma samples using *trans*-thiothixene as internal standard 311(1984)424
Narayanan, T.K. and Greenbaum, L.M.
Detection and quantitation of fluorescamine-labeled bradykinin, its analogues and metabolites using HPLC 306(1984)109
Naruse, H., see Todoriki, H. 310(1984)273
Nazareth, A., Jaramillo, L., Karger, B.L., Giese, R.W. and Snyder, L.R.
Automated analysis of antiepileptic drugs in serum by column-switching HPLC 309(1984)357
Nazareth, A., Joppich, M., Abdel-Baky, S., O'Connell, K., Sentissi, A. and Giese, R.W.
Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201
Nazareth, A., Joppich, M., Panthani, A., Fisher, D. and Giese, R.W.
Alkylation with dialkylsulfate and diisopropylethylamine 319(1985)382
Nazareth, A., O'Connell, K., Sentissi, A. and Giese, R.W.
Evidence for surface effects in an electron-capture detector 314(1984)219
Nazareth, A., see Mohamed, G.B. 314(1984)211
Neale, M.E., see Weaver, K.M. 301(1984)288

Nebinger, P.
Comparison of gel permeation and ion-exchange chromatographic procedures for the separation of hyaluronate oligosaccharides 320(1985)351
Needham, L.L., see Cline, R.E. 307(1984)420
Neggiani, P.P., see Marzo, A. 310(1984)51
Nelson, C., see Randinitis, E.J. 307(1984)210, 308(1984)345
Nelson, L.D., Brown, N.D. and Wiesmann, W.P.
Simultaneous assay of choline kinase and choline oxidase in tissue by high-performance cation-exchange chromatography and continuous radioactive detection 324(1985)203
Nelson, M.A., see Kabra, P.M. 307(1984)224
Nelson, P.E., see Cook, S.E. 325(1985)450
Nene, S., Anjaneyulu, B. and Rajagopalan, T.G.
Determination of diethylcarbamazine in blood using GC with alkali flame ionization detection 308(1984)334
Neukom, H.P., see Grob, K. Jr. 323(1985)237
Neumann, H.
Analysis of opium and crude morphine samples by capillary GC. Comparison of impurity profiles 315(1984)404
Neupert, M., see Bauer, S. 309(1984)243
New, M.I., see Loche, S. 317(1984)377
Newman, E., see Kogan, M.J. 306(1984)441
Newsom, A.E., see Said, H.M. 324(1985)65
Newsome, W.H., see Ryan, J.J. 303(1984)351
Ng, K.-T., Holland, M.L. and Plutte, J.A.
Determination of linogliride in biological fluids by HPLC 311(1984)418
Ng, K.-T., Plutte, J.A. and Galante, L.J.
Determination of bepridil in biological fluids by HPLC 309(1984)125
Ng, L.-K.
Separation and analysis of ethylenediamines as their *tert.*-butyldimethylsilyl derivatives by GLC and MS 314(1984)455
Ngoc, A.H., see Loo, J.C.K. 305(1984)194
N'Guyen Công, H., see Valencia, R. 325(1985)207
N'Guyen, N.-T., see Bonnaud, B. 318(1985)398
Nguyen-Huu, J.J., see Diquet, B. 311(1984)430
Nicholls, P.J., see Kamblawi, M.O. 309(1984)431
Nicholson, B.C., Bursill, D.B. and Couche, D.J.
Rapid method for the analysis of trihalomethanes in water 325(1985)221
Nicholson, G.J., see Bayer, E. 320(1985)393
Nickless, G.
Trace metal determination by chromatography (Review) 313(1984)129
Nicolaisen, E.M.
Characterization of a heterogeneous chicken plasma protein, HEF, by analytical isotachophoresis in agarose gel 320(1985)119
Nicolas, A., see Leroy, P. 317(1984)513
Nicoletti, I., see Grassini-Strazza, G. 322(1985)149
Nicolle, P., see Guechot, C. 303(1984)440
Nicot, G., Lachatre, G., Gonnet, C., Valette, J.-P., Merle, L., Nouaille, Y. and Bromet, N.
HPLC method for the determination of amineptine and its main metabolite in human plasma 306(1984)279
Nieass, C.S., see Wainwright, M.S. 321(1985)287
Niegreisz, Z., Szücs, L., Fekete, J., Horvai, G., Tóth, K. and Pungor, E.
Modifications of the wall-jet electrochemical detector for LC and flow analysis 316(1984)451
Nieh, J.-C., see Lo, J.-G. 322(1985)332
Nielen, M.W.F., Frei, R.W. and Brinkman, U.A.T.
Use of cation exchangers for the on-line preconcentration of polar anilines in LC 317(1984)557
Nielen, M.W.F., see Goewie, C.E. 301(1984)325
Nieman, R.H. and Clark, R.A.
Measurement of plant nucleotides by HPLC 317(1984)271
Niemelä, K., see Alén, R. 301(1984)273
Nierenberg, D.W.
Determination of serum and plasma concentrations of retinol using HPLC 311(1984)239
Niessen, W.M.A. and Poppe, H.
Open-tubular LC-MS using direct liquid introduction 323(1985)37
Niinae, T., see Tanaka, M. 314(1984)193
Nikolov, Z.L., Meagher, M.M. and Reilly, P.J.
HPLC of disaccharides on amine-bonded silica columns 319(1985)51
Nikolov, Z.L., Meagher, M.M. and Reilly, P.J.
HPLC of trisaccharides on amine-bonded silica columns 321(1985)393
Nikolov, Z.L. and Reilly, P.J.
Retention of carbohydrates on silica and amine-bonded silica stationary phases: application of the hydration model 325(1985)287
Nill, K., see Breyer-Pfaff, U. 309(1984)107
Nilsson, C.-A., see Andersson, K. 312(1984)482
Nimura, N., Toyama, A. and Kinoshita, T.
Optical resolution of amino acid enantiomers by HPLC 316(1984)547
Nishidate, K., see Yoshioka, M. 309(1984)63
Nishioka, M., see Lee, M.L. 302(1984)303
Nishioka, R., see Nakamura, M. 310(1984)450
Nissinen, E.
Determination of monoamine oxidase B activity by HPLC 309(1984)156
Nitsch, E., see Praznik, W. 303(1984)417
Nitsche, V., see Mascher, H. 306(1984)231
Noe, A.J., see Krol, G.J. 305(1984)105
Noever de Brauw, M.C. ten, see Tas, A.C. 310(1984)243
Noever de Brauw, M.C. ten, see Van der Greef, J. 323(1985)81
Noguchi, H., see Tokuma, Y. 311(1984)339
Noguchi, K., see Wada, H. 320(1985)369
Noh, I., see Mourey, T.H. 303(1984)361
Nohta, H., Noma, S., Ohkura, Y. and Yoo, B.-T.
Assay for catechol-O-methyltransferase in

erythrocytes using a new fluorogenic substrate, 2-(3,4-dihydroxyphenyl)naphtho[1,2-*d*]thiazole 308(1984)93
Nohta, H., see Hung, Y.-L. 305(1984)281
Nolan, C., see Chubb, C. 308(1984)11
Noma, S., see Nohta, H. 308(1984)93
Nomura, A., see Miyazaki, K. 310(1984)219
Nondek, L. and Chvalovský, V.
On-line pre-concentration and LC of azaarenes using nitroaromatic sorbents 312(1984)303
Nondek, L. and Minárik, M.
Chromatographic selectivity in LC of polycondensed aromatic hydrocarbons on 3-(2,4-Dinitroanilino)propyl silica 324(1985)261
Nora, M.V., Parkhurst, G.W., Thomas, R.W. and Carson, P.E.
HPLC-electrochemical assay method for primaquine in plasma and urine 307(1984)451
Nordholm, L. and Dalgaard, L.
Determination of trimethoprim metabolites including conjugates in urine using HPLC with combined UV and electrochemical detection 305(1984)391
Nose, N., see Tanaka, A. 306(1984)51
Nouaille, Y., see Nicot, G. 306(1984)279
Nousiainen, U. and Raatikainen, O.
HPLC of propanidid in rat plasma 308(1984)376
Novák, J., see Kleinert, T. 315(1984)85
Novák, J., see Vejrosta, J. 324(1985)269
Novak, M., see Moegel, H-J. 324(1985)29
Novotny, M., see Alasandro, M. 308(1984)1
Novotný, M., see Gluckman, J. 317(1984)443
Novotny, M., see Gluckman, J.C. 314(1984)103
Novotny, M., see Holland, M.L. 306(1984)23
Novotny, M., see Raymer, J. 325(1985)13
Nowakowska, E., see Goldberg, A.P. 316(1984)241
Nurok, D., see Johnson, E.K. 302(1984)135
Nwaiwu, J., see Agarwal, S.P. 323(1985)424

O

Obuchi, A., Aoyama, H., Ohi, A. and Ohuchi, H.
Determination of polycyclic aromatic hydrocarbons in diesel exhaust particulate matter and diesel fuel oil 312(1984)247
Ochs, H.R., see Abernethy, D.R. 307(1984)194
O'Connell, K., see Nazareth, A. 314(1984)201, 314(1984)219
Oddie, C.J., Jackman, G.P. and Bobik, A.
Measurement of xamoterol in plasma and urine by HPLC 308(1984)370
Oddie, C.J., see Jackman, G.P. 308(1984)301
Odink, J., see Tas, A.C. 310(1984)243
Ogawa, S., see Amagaya, S. 320(1985)430
Ogihara, Y., see Amagaya, S. 320(1985)430
Ogihara, Y., see Shimizu, K. 307(1984)488
Ogliaruso, M.A., see Brubaker, W.F. Jr. 324(1985)450
O'Hare, M.J., see D'Agostino, G. 305(1984)13
O'Hare, M.J., see Smith, J.A. 305(1984)295
Ohashi, S., see Baba, Y. 318(1985)319
Ohashi, S., see Yoza, N. 325(1985)385
Ohi, A., see Obuchi, A. 312(1984)247
Ohkuma, T. and Hara, S.
Mobile phase optimization for multi-functional solutes in LSC. Resolution of synthetic intermediates of oligodeoxyribonucleotides 323(1985)227
Ohkuma, T., see Hara, S. 316(1984)389
Ohkura, Y., see Hung, Y.-L. 305(1984)281
Ohkura, Y., see Ishida, J. 305(1984)381
Ohkura, Y., see Kai, M. 311(1984)257, 322(1985)473
Ohkura, Y., see Nakano, Y. 311(1984)390
Ohkura, Y., see Nohta, H. 308(1984)93
Ohkura, Y., see Tsuruta, Y. 309(1984)309
Ohlweiler, D.F., see Chan, K.Y. 306(1984)249
Ohm, T., see Lemmer, B. 309(1984)187
Ohmori, S., see Ikeda, M. 307(1984)111
Ohnishi, J., see Shoji, S. 319(1985)359
Ohuchi, H., see Obuchi, A. 312(1984)247
Oka, H., Matsumoto, H., Uno, K., Harada, K.-I., Kadowaki, S. and Suzuki, M.
Improvement of chemical analysis of antibiotics. VIII. Application of prepacked C_{18} Cartridge for the analysis of tetracycline residues in animal liver 325(1985)265
Oka, H. and Suzuki, M.
Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303
Oka, K., Kojima, K., Togari, A., Nagatsu, T. and Kiss, B.
Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43
Okada, K., see Amagaya, S. 320(1985)430
Okada, T. and Kuwamoto, T.
Elimination of the peak of a major ion in anion chromatography with UV detection 325(1985)327
Okada, Y., see Koizumi, K. 321(1985)145
Okano, Y., Kadota, T., Nagata, J., Matsuda, A., Ijima, S., Takahama, K. and Miyata, T.
Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251
Oke, A., see McKay, L. 311(1984)167
Okerholm, R.A., see Chan, K.Y. 306(1984)249
O'Kruk, R.J., Adams, M.A. and Philp, R.B.
Rapid and sensitive determination of

acetylsalicylic acid and its metabolites using reversed-phase HPLC 310(1984)343
Olanoff, L., see Litowitz, H. 311(1984)443
Olde, B., see Johansson, G. 311(1984)277
Oldfield, E.H., see Yeager, R.L. 305(1984)496
Olek, M., Blanchard, F. and Sudraud, G. Application of electrochemical detection to the determination of residues of some carbamate insecticides by HPLC (in French) 325(1985)239
Oliver, R.W.A., see Ager, R.P. 309(1984)1
Olkkonen, C., see Sjöberg, A.-M. 318(1985)149
Olsen, I., see Brondz, I. 308(1984)19, 308(1984)282, 310(1984)261, 311(1984)31, 311(1984)347
Olson, J.A., see Furr, H.C. 309(1984)299
Ontyd, J. and Schrader, J. Measurement of adenosine, inosine, and hypoxanthine in human plasma 307(1984)404
Ooijen, R.D. van, see Drost, R.H. 310(1984)193
Oort, W.J. van, see Den Hartigh, J. 306(1984)444
Oostelbos, M.C.J.M., see Visser, T. 309(1984)81
Oosterbaan, M.J.M., Dirks, R.J.M., Vree, T.B. and Van der Kleijn, E. Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323
Oppenheimer, L.E. and Mourey, T.H. Examination of the concentration response of evaporative light-scattering mass detectors 323(1985)297
Ord, D., see Schmitz, G. 307(1984)81
Origitano, T.C. and Collins, M.A. GC analysis of endogenous catecholamines, phenolic amines and derived isoquinolines using short glass capillary columns and electron-capture detection 311(1984)17
Orlan, B.P., see Strang, C. 314(1984)482
Orme, M.L'E., see Ward, S.A. 305(1984)239
Orme, P., see Perlstein, P. 325(1985)87
Oro, J., see Andrawes, F. 302(1984)181
Orr, J.D., see Weber, S.G. 322(1985)433
Ortel, T.L., see Takahashi, N. 317(1984)11
Osaka, N., see Morishita, H. 315(1984)253
Osek, J., see Oszczapowicz, J. 315(1984)95
Oshima, R., see Du, Y. 318(1985)378
Ossicini, L., see Fanali, S. 318(1985)440
Østensen, E.T., see Øyehaug, E. 308(1984)199
Östling, S.G., see Haglund, Å.C. 318(1985)57
Ostrovský, I., see Soják, L. 323(1985)414
Ostrowksi, J., Theumer, J., Gärtner, W., Resag, K. and Passing, H. Determination of pirlindole in plasma and urine by HPLC 309(1984)115
Oszczapowicz, J., Osek, J. and Dolecka, E. Retention indices of dimethylformamidines, dimethylacetamidines and tetramethylguanidines on a non-polar column 315(1984)95
Otterburn, M.S. and Gargan, P.E. Determination of dityrosine in protein fractions isolated from merino wool using reversed-phase HPLC 303(1984)429
Overdijk, B., see Beem, E.P. 323(1985)439
Owada, E., see Kurosawa, N. 305(1984)485
Owens, J.A. and Robinson, J.S. HPLC determination of plasma lactate specific radioactivity 307(1984)380
Oxford, J., see Lant, M.S. 323(1985)143
Øyehaug, E., Østensen, E.T. and Salvesen, B. HPLC determination of citalopram and four of its metabolites in plasma and urine samples from psychiatric patients 308(1984)199
Ozaki, T., see Ishida, R. 305(1984)73

P

Pacáková, V., see Kysilka, R. 320(1985)414
Pace-Asciak, C.R. and Micallef, S. GC-MS profiling with negative-ion chemical ionization detection of prostaglandins and their 15-keto and 15-keto-13,14-dihydro catabolites in rat blood 310(1984)233
Pacholec, F. and Poole, C.F. Evaluation of a calibration marker scheme for open tubular column GC with on-column injection and electron-capture detection 302(1984)289
Padovani, P., Deves, C., Bianchetti, G., Thenot, J.P. and Morselli, P.L. Determination of progabide and its main acid metabolite in biological fluids using HPLC and electrochemical detection. Application to the measurement of blood/plasma partition ratio 308(1984)229
Palmer, J.K., see Pierotti, J.A. 306(1984)377
Palmisano, F., Desimoni, E. and Zambonin, P.G. HPLC with polarographic and voltammetric anodic detection: simultaneous determination of allopurinol, oxipurinol and uric acid in body fluids 306(1984)205
Panthani, A., see Nazareth, A. 319(1985)382
Panu, A.M., see Desai, H.K. 322(1985)223
Paphassarang, S., Raynaud, J., Godeau, R.P. and Binsard, A.M. Qualitative and quantitative analysis of hyoscyamine-atropine and scopolamine in mother tinctures of Solanaceae by HPLC (in French) 319(1985)412
Papp, O., see Valkó, K. 301(1984)355
Parcher, J.F. and Hyver, K.J. GC determination of binary adsorption isotherms 302(1984)195
Parekh, N.J., Fatmi, A.A., Tshabalala, M.A. and Rogers, L.B. Effects of various strengths of acids on the electrophoretic behavior of weak bases in a non-aqueous solvent 314(1984)65

Parente, E.S. and Wetlaufer, D.B.
Effects of urea-thermal denaturation on the high-performance cation-exchange chromatography of α-chymotrypsinogen-A 314(1984)337
Parke, D.V., see Amoah, A.G.B. 305(1984)401
Parker, C.E., Yamaguchi, K., Harvan, D.J., Smith, R.W. and Hass, J.R.
LC-chloride-attachment negative chemical ionization MS 319(1985)273
Parker, T.D. III, see Randinitis, E.J. 307(1984)210
Parkhurst, G.W., see Nora, M.V. 307(1984)451
Parkin, J.E.
HPLC assay of menthol using indirect photometric detection 303(1984)436
Parkin, J.E. and Lau, H.T.
HPLC of low-molecular-weight aliphatic alcohols using indirect photometric detection 314(1984)488
Parsons, M.L., see Crouch, R.L. 303(1984)53
Parsons, M.L., see Hawley-Fedder, R.A. 314(1984)263, 315(1984)201, 315(1984)211
Partos, C., see Dell, D. 317(1984)483
Paryjczak, T., Zieliński, P. and Farbotko, M.J.
Effect of activated sorption processes on the results of thermoprogrammed reduction investigations on Fe-Al_2O_3 catalysts 301(1984)270
Passing, H., see Ostrowksi, J. 309(1984)115
Patel, A.H., see Vinson, J.A. 307(1984)493
Patel, V., see Sindhuphak, T. 314(1984)408
Patience, R.L. and Rees, L.H.
Comparison by gel filtration chromatography and reversed-phase HPLC of the immunoreactive growth hormone composition of a human pituitary extract 324(1985)385
Patoiseau, J.-F., see Bonnaud, B. 318(1985)398
Patrick, D.W. and Kracht, W.R.
Single-valve, single-pump, HPLC column-switching analysis of hydrolyzed wood components 318(1985)269
Patrick, J., see Alton, K.B. 311(1984)319
Patsalides, E., see Dilli, S. 318(1985)350
Patterson, S.E.
Rapid and sensitive analysis of terazosin in plasma, peritoneal dialysis solution, and urine using HPLC with fluorescence detection 311(1984)206
Pauliukonis, L.T., see Gravallese, D.A. 310(1984)71
Paulsen, B.S., see Haavik, S. 321(1985)199
Pauncz, J. Kádár, see Kádár Pauncz, J. 322(1985)386
Peal, J.A., see Bailey, E. 310(1984)199
Peetre, I.-B., Ellren, O. and Smith, B.E.F.
GC investigation of organometallic compounds and their carbon analogues. VI. Determination, calculation and correlation of Kováts retention indices for alkylalkoxysilanes 318(1985)41
Pekarovičová, A., see Polonský, J. 320(1985)111
Peleran, J.C. and Bories, G.F.
GC determination and MS confirmation of traces of indole and 3-methylindole (skatole) in pig back fat 324(1985)469
Pelletier, S.W., see Desai, H.K. 322(1985)223
Pellinen, J. and Salkinoja-Salonen, M.
Aqueous size exclusion chromatography of industrial lignins 322(1985)129
Pellizzari, E.D., see Voyksner, R.D. 312(1984)221
Peltonen, K., Pfäffli, P. and Itkonen, A.
Determination of aldehydes in air as dimethone derivatives by GC with electron-capture detection 315(1984)412
Peña, M.I. arranz, see Arranz Peña, M.I. 305(1984)210
Peng, C.T., see Buchman, O. 312(1984)75
Peng, D.-R., see Petters, I. 306(1984)241
Perlstein, P. and Orme, P.
Determination of polymeric hindered-amine light stabilisers in plastics by pyrolysis-GC 325(1985)87
Perrone, P.A. and Brown, P.R.
Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53
Perrone, P.A. and Brown, P.R.
Ion-pair chromatography of nucleotides 317(1984)301
Persson, B. and Wendsjö, S.
Oxygen effects in amperometric LC detection of omeprazole at a mercury electrode 321(1985)375
Persson, B.-A., Jansson, S.-O., Johansson, M.-L. and Lagerström, P.-O.
Influence of mobile phase additives in LC of phenoxypropanolamines 316(1984)291
Persson, B.-A., see Lagerström, P.-O. 309(1984)347
Pertierra-Rimada, E., see Fernández-Sánchez, E. 312(1984)69
Petchey, M. and Crabbe, M.J.C.
Analysis of carbohydrates in lens, erythrocytes, and plasma by HPLC of nitrobenzoate derivatives 307(1984)180
Peters, G.J., Kraal, I., Laurensse, E., Leyva, A. and Pinedo, H.M.
Separation of 5-fluorouracil and uracil by ion-pair reversed-phase HPLC on a column with porous polymeric packing 307(1984)464
Peters, T.J., see Bottiglieri, T. 311(1984)354
Peters, T.J., see Lim, C.K. 316(1984)397, 317(1984)333
Peterson, E.A., see Torres, A.R. 316(1984)125
Petit, J., see Lartigue-Mattei, C. 310(1984)407
Petters, I., Peng, D.-R. and Rane, A.
Quantitation of clonazepam and its 7-amino and 7-acetamido metabolites in plasma by HPLC 306(1984)241
Pettersson, C.
Chromatographic separation of enantiomers of acids with quinine as chiral counter ion 316(1984)553
Peuchant, E., Covi, G. and Jensen, R.
Faecal lipid chromatography. I.

Quantitative determination with Chromarods 310(1984)297
Pévet, P., see Beck, O. 311(1984)1
Peyton, A.L., see Toren, P.C. 305(1984)171
Pezzuto, J.M., see Ratko, T.A. 324(1985)484
Pfäffli, P., see Peltonen, K. 315(1984)412
Pfleger, K., see Maurer, H. 305(1984)309, 306(1984)125
Philippe, P., see Bontemps, J. 307(1984)283
Phillips, T.M., More, N.S., Queen, W.D., Holohan, T.V., Kramer, N.C. and Thompson, A.M.
High-performance affinity chromatography: a rapid technique for the isolation and quantitation of IgG from cerebral spinal fluid 317(1984)173
Philp, R.B., see O'Kruk, R.J. 310(1984)343
Pierotti, J.A., Dickinson, A.G., Palmer, J.K. and Driskell, J.A.
LC separation and quantitation of B_6 vitamers in selected rat tissues 306(1984)377
Pietrogrande, M.C., see Guerra, M.C. 320(1985)281
Pifferi, G., see Crolla, T. 310(1984)139
Pihlajamäki, K.K., see Lindberg, R.L.P. 309(1984)369
Pike, A.W., Moynihan, C., Kibler, S., Marsh, P.G. and Fennessey, P.V.
Semi-micro quantitative analysis of complex urinary steroid mixtures in healthy and diseased states 306(1984)39
Pimprikar, G.D., Coign, M.J., Sakurai, H. and Heitz, J.R.
HPLC determination of ecdysteroid titers in the house fly 317(1984)413
Pinczewski, W.V., see Zadro, S. 323(1985)305
Pinedo, H.M., see Elferink, F. 320(1985)379
Pinedo, H.M., see Peters, G.J. 307(1984)464
Pinedo, H.M., see Van der Vijgh, W.J.F. 310(1984)335
Pinkerton, T.C., see Cook, S.E. 325(1985)450
Piotrovskii, V.K., Zhirkov, Y.A. and Metelitsa, V.I.
Ion-exchange HPLC in drug assay in biological fluids. IV. Nadolol diastereomers: demonstration of pharmacokinetic and binding equivalence 309(1984)421
Piovetti, L., see Serve, L. 322(1985)257
Pirastru, M.G., see Cossu, G. 307(1984)103
Pirkle, W.H. and Hyun, M.H.
Reversed-phase chromatographic resolution of N-(3,5-dinitrobenzoyl)-α-amino acids on chiral stationary phases 322(1985)287
Pirkle, W.H. and Hyun, M.H.
α-Arylalkylamine-derived chiral stationary phases. Evaluation of urea linkages 322(1985)295
Pirkle, W.H. and Hyun, M.H.
Preparation and use of hydantoin-based chiral stationary phases 322(1985)309
Pirkle, W.H., Hyun, M.H. and Bank, B.
Rational approach to the design of highly-effective chiral stationary phases 316(1984)585
Pirkle, W.H., Tsipouras, A. and Sowin, T.J.
Preparative separation of enantiomers by flash chromatography 319(1985)392
Plagne, R., see Bernard, D. 308(1984)322
Plaisted, S.M., DeZwaan, J. and Snider, B.G.
HPLC determination of acid-catalyzed degradation products of methyl carboprost in a polymeric controlled-release device 314(1984)369
Plaisted, S.M., see Theis, D.L. 321(1985)209
Plakunov, I., see Toren, P.C. 305(1984)171
Platt, D., see Rieck, W. 305(1984)419
Plutte, J.A., see Ng, K.-T. 309(1984)125, 311(1984)418
Podcasy, J.A. and Weber, S.G.
Non-aqueous LC of some metalloporphyrins on a reversed-phase column 315(1984)111
Pollack, G.M., Slaughter, R.L., Buchanan, J.F. and Shen, D.D.
HPLC procedure for the determination of di-(2-ethylhexyl)phthalate in human blood specimens. Problems of variable-extraction yield and the use of standard addition for calibration 311(1984)101
Polonský, J., Repášová, Ľ., Pekarovičová, A. and Košík, M.
Isotachophoretic analysis of saccharinic acids 320(1985)111
Polta, J.A., Johnson, D.C. and Merkel, K.E.
LC separation of aminoglycosides with pulsed amperometric detection 324(1985)407
Polyakova, N.P., see Bazyl'chick, V.V. 321(1985)240
Polzin, G., see Koch, D.D. 317(1984)327
Pommier, C., see Marlet, B. 325(1985)75
Pommier, F., see Sioufi, A. 305(1984)95
Ponchet, M., see Abad, P. 318(1985)417
Ponte, G. de, see Marcucci, F. 309(1984)132
Ponte, G. de, see Mussini, E. 305(1984)450
Poole, C.F., Butler, H.T., Coddens, M.E., Khatib, S. and Vandervennet, R.
Comparison of methods for separating polycyclic aromatic hydrocarbons by HPTLC 302(1984)149
Poole, C.F., see Dhanesar, S.C. 324(1985)415
Poole, C.F., see Pacholec, F. 302(1984)289
Poon, B.T., see Brown, N.D. 309(1984)426
Poore, C.M.B., see Mullin, B.R. 305(1984)512
Poppe, H., see Niessen, W.M.A. 323(1985)37
Poppe, H., see Welling, P. 321(1985)450
Porath, J., see Hemdan, E.S. 323(1985)247, 323(1985)255, 323(1985)265
Porath, J., see Maisano, F. 321(1985)305
Porath, J., see Ramadan, N. 321(1985)81, 321(1985)93, 321(1985)105
Porcelli, F., see Loche, S. 317(1984)377
Porsch, B.
HPLC of oligosaccharides in water on a reversed phase doped with primary amino groups 320(1985)408
Porter, W.R., see Jeyaraj, G.L. 315(1984)378
Porumb, H., see Porumb, T. 319(1985)218
Porumb, T., Porumb, H., Lascu, I. and Proinov, I.

Cibacron Blue 3G-A-substituted cross-linked agar beads: an inexpensive, fast-flow affinity medium for large scale applications 319(1985)218
Pospíchal, J., Deml, M., Žemlová, Z. and Boček, P.
Determination of relative ionic mobilities by capillary isotachophoresis 320(1985)139
Postma, G.J., see Van der Vijgh, W.J.F. 310(1984)335
Potěšil, T. and Potěšilová, H.
TLC of enamino ketones. II 312(1984)387
Potěšilová, H., see Potěšil, T. 312(1984)387
Pötter, H., Hülm, M. and Schumann, S.
Partition chromatography of ergot alkaloids on formamide-saturated silica gel columns (in German) 319(1985)440
Poulter, C.D., see Yang, T.-H. 312(1984)121
Poupet, A., see Abad, P. 318(1985)417
Pourcq, P. de, see De Pourcq, P. 321(1985)441
Pourcq, P. de, see Roets, E. 303(1984)117
Powell, J.R., see Wainer, I.W. 306(1984)405
Powell, S.C., Friedlander, E.R. and Shihabi, Z.K.
Myoglobin determination by HPLC 317(1984)87
Powis, G., see Reece, P.A. 306(1984)417
Powles, P., Robinson, D.T., Andrews, R.S. and Robinson, P.R.
Determination of trilostane and ketotrilostane in human plasma by HPLC 311(1984)434
Poynter, D. and Landon, M.
Chloral hydrate is compatible with isoelectric focusing 325(1985)333
Pratt, K.C., see Harvey, T.G. 319(1985)230
Praznik, W., Beck, R.H.F. and Nitsch, E.
Determination of fructan oligomers of degree of polymerization 2-30 by HPLC 303(1984)417
Predieri, G., see Casoli, A. 303(1984)404
Price, C., see Styring, M.G. 319(1985)115
Price, G.J., see Ashworth, A.J. 324(1985)231
Priebe, S.R. and Howell, J.A.
Post-column reaction detection system for the determination of organophosphorus compounds by LC 324(1985)53
Prinoth, M. and Mutschler, E.
Fluorimetric determination of maprotiline in urine and plasma after TLC separation 305(1984)508
Přistoupil, T.I., see Kramlová, M. 303(1984)422
Privat, G., see Andary, C. 303(1984)312
Proinov, I., see Porumb, T. 319(1985)218
Prosperi, T., see Fanali, S. 318(1985)440
Prueksaritanont, T., Chen, M.-L. and Chiou, W.L.
Simple and micro HPLC method for simultaneous determination of *p*-aminohippuric acid and iothalamate in biological fluids 306(1984)89
Prusík, Z. and Kašička, V.
Micropreparative isotachophoretic electrodesorption of monoclonal antibodies from an affinity adsorbent 320(1985)81
Prusík, Z., see Kašička, V. 320(1985)33, 320(1985)75
Prusoff, W.H., see Brubaker, W.F. Jr. 322(1985)455
Pullen, R.H., see Cox, J.W. 307(1984)155
Pungor, E., see Niegreisz, Z. 316(1984)451
Punkt, J. and Häcker, R.
Advantages and limitations of isotachophoretic determinations of metabolites in tissue extracts 320(1985)105
Purnell, J.H., Rodriguez, M. and Williams, P.S.
Approximate approach to correction for carrier compressibility effects in coupled packed GC columns 323(1985)402
Purnell, J.H. and Williams, P.S.
General theory for correction for compressibility effects in binary coupled GC columns and the procedure for window diagram optimisation of relative lengths 321(1985)249
Purnell, J.H. and Williams, P.S.
Theory of the efficiency of serially connected GC columns and the effect on optimized speed of analysis 325(1985)1
Purnell, J.H., see Ali, S.G.A.H. 302(1984)119
Purser, C., Baltar, A., Ho, I.K. and Hume, A.S.
New rapid method of analysis of cefoxitin in serum and bone, by HPLC 311(1984)135
Putman, L.J. and Butler, L.G.
Fractionation of condensed tannins by counter-current chromatography 318(1985)85
Putnam, F.W., see Takahashi, N. 317(1984)11
Pyysalo, H., see Sjöberg, A.-M. 319(1985)90

Q

Qadri, F., see Yasmin, T. 315(1984)425
Qiu, Z.Y., see Rokushika, S. 320(1985)335
Queen, W.D., see Phillips, T.M. 317(1984)173

R

Raat, W.K. de, see De Raat, W.K. 310(1984)41
Raatikainen, O., see Nousiainen, U. 308(1984)376
Rabe, H., see Krüger, J. 307(1984)387
Raddats, M.A., see Burch, J.E. 308(1984)165
Raemsch, K.D., see Krol, G.J. 305(1984)105
Rajagopalan, T.G., see Nene, S. 308(1984)334

Rajgopal, S. and Vijayalakshmi, M.A.
Metal-ion mediated interaction of luciferase with tetraiodofluorescein 315(1984)175
Ramadan, N. and Porath, J.
α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81
Ramadan, N. and Porath, J.
Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93
Ramadan, N. and Porath, J.
Separation of serum proteins on a Fe^{3+}-monohydroxamate adsorbent 321(1985)105
Ramjit, H.G., see Duggan, D.E. 306(1984)69
Ramsey, E.D., see Games, D.E. 323(1985)67
Ramus, T.L., Hein, S.J. and Thomas, L.C.
Determinations of chlorinated hydrocarbons by GC using response factor calibration 314(1984)243
Randinitis, E.J., Kinkel, A.W., Nelson, C. and Parker, T.D. III
GC determination of gemfibrozil and its metabolites in plasma and urine 307(1984)210
Randinitis, E.J., Nelson, C. and Kinkel, A.W.
GC determination of bevantolol in plasma 308(1984)345
Randolph, W.C., see Boppana, V.K. 317(1984)463
Rane, A., see Petters, I. 306(1984)241
Rantwijk, F. van, see Stikkelman, R.M. 322(1985)220
Rapp, U., see Van der Greef, J. 323(1985)81
Rascol, J.P., see Andary, C. 303(1984)312
Raso, M.A., see García-Raso, A. 302(1984)107
Rassi, Z. el, see Chang, J.-P. 319(1985)396
Ratko, T.A. and Pezzuto, J.M.
Improved fluorometric method for quantitative HPLC analysis of methylated guanine derivatives in DNA 324(1985)484
Ratnayake, W.M.N., see Banerjee, A.K. 319(1985)215
Ratti, B., see Bernareggi, A. 309(1984)415
Rausch, T., Helmlinger, J. and Hilgenberg, W.
High-performance LC separation and some properties of (*E*)- and (*Z*)-3-indoleacetaldoxime 318(1985)95
Ravenscroft, P.J., see Charles, B.G. 306(1984)424
Rawski, V., see Daftsios, A.C. 305(1984)145
Ray, B., Ghosal, P.K., Thakur, S. and Majumdar, S.G.
Sulphosalicylic acid as spray reagent for the detection of sugars on thin-layer chromatograms 315(1984)401
Ray, G., Crook, M., West, N., Kwoka, M., Rehagen, G., Cox, J., Murrill, E. and Flora, K.
Comparison of the analysis of Δ^9-tetrahydrocannabinol capsules by HPLC and capillary GC 317(1984)455
Raymer, J., Wiesler, D. and Novotny, M.
Structure-retention studies of model ketones by capillary GC 325(1985)13
Raynaud, J., see Paphassarang, S. 319(1985)412
Redfern, B., see Demaster, E.G. 308(1984)83
Redmond, J.W., see McNicholas, P.A. 315(1984)451
Redwood, S.M., see Nakazawa, H. 307(1984)323
Reece, P.A., McCall, J.T., Powis, G. and Richardson, R.L.
Sensitive HPLC assay for platinum in plasma ultrafiltrate 306(1984)417
Reeck, G.R., see Wen, L. 314(1984)436
Rees, L.H., see Patience, R.L. 324(1985)385
Reeve, J.R. Jr., Ho, F.J., Walsh, J.H., Ben-Avram, C.M. and Shively, J.E.
Rapid high-yield purification of canine intestinal motilin and its complete sequence determination 321(1985)421
Reeve, R.N., see Lethbridge, J.W. 314(1984)387
Rege, A.B., Lertora, J.J.L., White, L.E. and George, W.J.
Rapid analysis of valproic acid by GC 309(1984)397
Regnier, F.E., see Fausnaugh, J.L. 317(1984)141
Regnier, F.E., see Kopaciewicz, W. 318(1985)157
Rehagen, G., see Ray, G. 317(1984)455
Reichmann, G., see Krüger, J. 307(1984)387
Reid, R., see Taylor, R.B. 316(1984)279
Reijenga, J.C., Aben, G.V.A., Lemmens, A.A.G., Verheggen, T.P.E.M., De Bruijn, C.H.M.M. and Everaerts, F.M.
Determination of quinine in beverages, pharmaceutical preparations and urine by isotachophoresis 320(1985)245
Reijenga, J.C., Lemmens, A.A.G., Verheggen, T.P.E.M. and Everaerts, F.M.
Aspects of detection and identification in isotachophoresis 320(1985)67
Reijenga, J.C., see De Bruijn, C.H.M.M. 320(1985)205
Reijenga, J.C., see Everaerts, F.M. 320(1985)263
Reijenga, J.C., see Lemmens, A.A.G. 320(1985)193
Reijenga, J.C., see Verheggen, T.P.E.M. 320(1985)99
Reilly, P.J., see Nikolov, Z.L. 319(1985)51, 321(1985)393, 325(1985)287
Reineke, F.J. and Bächmann, K.
GC determination of C_2-C_8 hydrocarbons and halocarbons in ambient air by simultaneous use of three detectors 323(1985)323
Reiner, J., see Thoma, H. 309(1984)17
Reinikainen, K., see Herranen, J. 307(1984)241
Rembold, H. and Lackner, B.
Convenient method for the determination of picomole amounts of juvenile hormone 323(1985)355
Remelli, M., see Dondi, F. 315(1984)67
Repášová, L., see Polonský, J. 320(1985)111
Resag, K., see Ostrowksi, J. 309(1984)115
Reuning, R.H., see Bockbrader, H.N. 310(1984)85
Reutter, W., see Josić, D. 317(1984)35
Revich, G.G., Hillebrand, G.G. and Beattie, K.L.

HPLC purification of deoxynucleoside 5'-triphosphates and their use in a sensitive electrophoretic assay of misincorporation during DNA synthesis 317(1984)283
Rhodes, G.R., see Holland, M.L. 306(1984)23
Ribeiro, J.A., see Figueira, M.A. 325(1985)317
Ricci, G., see Kodama, H. 311(1984)183
Rice, L.G.
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523
Richards, G.N., see Blake, J.D. 312(1984)211
Richardson, R.L., see Reece, P.A. 306(1984)417
Riddle, R.M., see Bear, G.R. 302(1984)65
Rideout, J.M., see Lim, C.K. 317(1984)333
Rieck, W. and Platt, D.
Determination of 3,7-dimethyl-1-(5-oxohexyl)-xanthine (pentoxifylline) and its 3,7-dimethyl-1-(5-hydroxyhexyl)xanthine metabolite in the plasma of patients with multiple diseases using HPLC 305(1984)419
Riekkinen, P., see Herranen, J. 307(1984)241
Riggs, C.E. Jr., see Nakazawa, H. 307(1984)323
Righetti, P.G., see Cossu, G. 307(1984)103
Righini-Tapie, A., see Maurs, M. 325(1985)444
Rio, R.M. del, see Herranz, A.S. 309(1984)139
Rios, J.L., see Villar, A. 303(1984)306
Risk, M., Holland, O.B. and Brown, H.
Rapid chromatographic purification of urinary steroid glucuronates for determination of aldosterone secretion rate 317(1984)367
Risk, M., see Benedict, C.R. 317(1984)27
Rispe, R., see Fels, J.-P. 308(1984)273
Riutta, A., Seppälä, E. and Vapaatalo, H.
Rapid, sensitive method for detecting different arachidonic acid metabolites by TLC: the use of autoradiography 307(1984)185
Rix, C.J., see Cookson, D.J. 312(1984)237
Robards, K., see Dilli, S. 312(1984)109, 322(1985)1, 322(1985)23, 324(1985)75
Roberts, S.G., see Burch, J.E. 308(1984)165
Robins, R.H., see Kissinger, L.D. 321(1985)353
Robinson, D.T., see Powles, P. 311(1984)434
Robinson, J.L. and Macrae, R.
Comparison of detection systems for the HPLC analysis of complex triglyceride mixtures 303(1984)386
Robinson, J.L., Tsimidou, M. and Macrae, R.
Evaluation of the mass detector for quantitative detection of triglycerides and fatty acid methyl esters 324(1985)35
Robinson, J.S., see Owens, J.A. 307(1984)380
Robinson, P.R., see Powles, P. 311(1984)434
Robinson, W.S., see Smith, G.A. 307(1984)410
Robinson, W.T., see Smith, H.T. 305(1984)353
Robowsky, K.-D., see Lepom, P. 322(1985)261
Rodriguez, G.P. and Claus-Walker, J.
Measurement of hydroxylysine glycosides in urine and its application to spinal cord injury 308(1984)65
Rodriguez, M., see Purnell, J.H. 323(1985)402
Rodriguez, P.A., see Keough, T. 312(1984)450
Roedder, E., see Andrawes, F. 302(1984)181
Roeraade, J., Blomberg, S. and Flodberg, G.
Miniature connector with a low dead volume for fused-silica capillary columns 301(1984)454
Roeraade, J., Flodberg, G. and Blomberg, S.
New injection syringe for GC 322(1985)55
Roets, E., De Pourcq, P., Toppet, S., Hoogmartens, J., Vanderhaeghe, H., Williams, D.H. and Smith, R.J.
Isolation and structure elucidation of ampicillin and amoxicillin oligomers 303(1984)117
Roets, E., see De Pourcq, P. 321(1985)441
Roets, E., see Dihuidi, K. 325(1985)413
Rogers, H.J., see Adam, A.M. 307(1984)129
Rogers, H.J., see Talha, M.R.Z. 311(1984)194
Rogers, L.B., see Alexander, J.N. IV 318(1985)289
Rogers, L.B., see Men, L.-C. 314(1984)55
Rogers, L.B., see Parekh, N.J. 314(1984)65
Rokushika, S., Qiu, Z.Y. and Hatano, H.
Microbore packed-column anion-exchange LC of nucleobases, nucleosides and nucleotides 320(1985)335
Rolland, M., see Marriq, C. 323(1985)395
Rolland, P.H., see Santoni, Y. 306(1984)165
Rolle, R., see Chism, G.W. III 317(1984)263
Röllgen, F.W., see Schmelzeisen-Redeker, G. 323(1985)127
Ronchi, S., see Biondi, P.A. 309(1984)151
Rooney, L., see Brodie, R.R. 310(1984)353
Roos, C., see Karlsson, L. 306(1984)183
Rose, S.M., see Tsao, C.S. 308(1984)306
Rosen, M., see Loche, S. 317(1984)377
Rosenberg, C. and Savolainen, H.
Detection of urinary amine metabolites in toluene diisocyanate exposed rats 323(1985)429
Rosenberg, M. and Choi, R.L.
HPLC analysis of flordipine in human plasma 308(1984)382
Rosenfeld, J.M., Brown, G.M., Walker, C.H. and Sprung, C.
Use of impregnated reagents in the pentafluorobenzylation of indoleamine metabolites 325(1985)309
Rosengren, B., see Månsson, J.-E. 322(1985)465
Rosset, R., see Foucault, A. 317(1984)41
Rosset, R., see Souteyrand, C. 316(1984)373
Rossetti, D., see Castagnola, M. 307(1984)91
Rössner, B., see Schmuckler, G. 302(1984)15
Roth, E.F. Jr., see Stump, D.D. 306(1984)371
Rounds, M.A., see Kopaciewicz, W. 318(1985)157
Roussel, J.L., see Andary, C. 303(1984)312
Rovei, V., see Vajta, S. 311(1984)329
Rowland, S.J., Larcher, A.V., Alexander, R. and Kagi, R.I.
GC and GC-MS analysis of diastereomeric acyclic isoprenoid esters using fused-silica capillary columns 312(1984)395
Roy, F. and Hegde, M.V.
Rapid procedure for purification of β-amylase from *Ipomea batata* 324(1985)489

Roy, S.D., McKay, G., Hawes, E.M. and Midha, K.K.
GC quantitation of methoxyphenamine and three of its metabolites in plasma 310(1984)307
Roy, S.K., Weber, D.V. and McGregor, W.C.
High-performance immunosorbent purification of recombinant leukocyte A interferon 303(1984)225
Ruckmick, S.C. and Hurtubise, R.J.
LC systems for the separation of polycyclic aromatic hydrocarbon and nitrogen heterocycle compounds present in coal liquids 321(1985)343
Rudland, P.S., see Smith, J.A. 305(1984)295
Rudolph, F.B., see Greenhut, J. 319(1985)461
Rudzinski, W.E., Montgomery, T.L., Frye, J.S., Hawkins, B.L. and Maciel, G.E.
Synthesis, solid-state nuclear magnetic resonance characterization and chromatographic evaluation of a diphenylphosphine-octadecylsilane hybrid bonded phase 323(1985)281
Ruijten, H.M., Van Amsterdam, P.H. and De Bree, H.
Conical precolumn as loading buffer for the main column 314(1984)183
Rupp, B.H., see Mullin, B.R. 305(1984)512
Rusk, M.L., see Theis, D.L. 321(1985)209
Russell, A.S., see Jamali, F. 310(1984)327
Russell, T.P., see Mumby, S.J. 319(1985)241
Rüstow, B., see Krüger, J. 307(1984)387
Ruud-Christensen, M. and Salvesen, B.
Separation of (*R*)- and (*S*)-proxyphylline as diastereoisomeric camphanates by reversed-phase LC 303(1984)433
Ryan, J.J., Lizotte, R. and Newsome, W.H.
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351
Ryan, P.J. and Honeyman, T.W.
Determination of fatty acids by HPLC of Dns-ethanolamine derivatives 312(1984)461
Ryan, P.J., McGoldrick, K., Stickney, D. and Honeyman, T.W.
Formation and separation of fluorescent derivatives of phosphatidic acid 320(1985)421
Rydzik, R., Terragno, A. and Tackett, R.
Picogram detection of eicosanoids by UV absorbance after narrow-bore HPLC. Comparison with conventional-bore column 308(1984)31

S

Saarinen, R., see Korpela, T.K. 318(1985)333
Sabljić, A.
Calculation of retention indices by molecular topology: chlorinated alkanes 314(1984)1
Sabljić, A.
Calculation of retention indices by molecular topology. Chlorinated benzenes 319(1985)1
Sadler, A.J., Micanovic, R., Katzenstein, G.E., Lewis, R.V. and Middaugh, C.R.
Protein conformation and reversed-phase HPLC 317(1984)93
Sadler, G., see Cook, S.E. 325(1985)450
Šafařík, I.
Isolation of trypsin by column chromatography on tea particles 315(1984)478
Šafařík, I., Laudová, Z. and Králová, B.
Purification of chymotrypsin and trypsin by CC on agar gel particles 303(1984)283
Sahm, H., see Hamacher, K. 319(1985)311
Said, H.M., Newsom, A.E., Tippins, B.L. and Mathews, R.A.
Reversed-phase HPLC of some ultra-heterogeneous and covalently-modified proteins from human hair 324(1985)65
Saito, M., see Hoshino, T. 316(1984)473
Saitoh, Y., see Uchino, K. 308(1984)241
Sakabe, Y., see Terada, H. 318(1985)299
Sakaguchi, T., see Ikeda, M. 305(1984)261
Sakamoto, Y., see Kai, M. 322(1985)473
Sakurai, H., see Pimprikar, G.D. 317(1984)413
Salama, A.I., see Saller, C.F. 309(1984)287
Salkinoja-Salonen, M., see Pellinen, J. 322(1985)129
Saller, C.F. and Salama, A.I.
Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287
Salmon, S.E., see Berens, M.E. 307(1984)251
Salotto, A.W., see Weiser, E.L. 303(1984)1
Salvesen, B., see Øyehaug, E. 308(1984)199
Salvesen, B., see Ruud-Christensen, M. 303(1984)433
Salvi, M.L., see Castagnola, M. 307(1984)91
Samain, D., see Inchauspé, G. 303(1984)277
Samperi, R., see Andreolini, F. 310(1984)208
Samperi, R., see Borra, C. 311(1984)9
Såmūndsen, J.A., see Jackson, P.J. 325(1985)336
Sander, L.C. and Wise, S.A.
Influence of substrate parameters on column selectivity with alkylbonded-phase sorbents 316(1984)163
Sander, L.C., see Diamondstone, B.I. 321(1985)319
Sanders, R.A., see Keough, T. 312(1984)450
Saner, W.A., see Strang, C. 314(1984)482

Sangermano, L., see Lai, S.-T. 312(1984)313, 321(1985)325, 322(1985)338, 324(1985)436
Sano, H., see Baba, S. 305(1984)119
Sano, T., see Suemitsu, R. 318(1985)139
Santini, F., see Crolla, T. 310(1984)139
Santoni, Y., Rolland, P.H. and Cano, J.-P.
Determination of isosorbide dinitrate and its mononitrate metabolites in human plasma using Extrelut purification and capillary column GLC 306(1984)165
Sappa, E., see Casoli, A. 303(1984)404
Sârbu, C., Horn, M. and Hantz, A.
Uranyl salts as fluorescence reagents for the TLC detection of complex-forming compounds (in German) 322(1985)272
Sarkissian, T.M., see Tameesh, A.H.H. 321(1985)59
Sarzanini, C., see Mentasti, E. 322(1985)177
Sasagawa, T., Ericsson, L.H., Teller, D.C., Titani, K. and Walsh, K.A.
Separation of peptides on a polystyrene resin column 307(1984)29
Sasahara, T., see Ikarashi, Y. 322(1985)191
Sasaki, K., see Kodama, H. 311(1984)183
Sasaki, K., see Mikasa, H. 305(1984)204
Sato, M., see Tsuchiya, H. 309(1984)43
Sato, T., see Kadofuku, T. 311(1984)93
Sattayasai, J., see Sattayasai, N. 307(1984)235
Sattayasai, N., Sattayasai, J. and Hahnvajanawong, V.
HPLC determination of gossypol in plasma 307(1984)235
Saura-Calixto, F., García-Raso, A. and García-Raso, J.
Heats of formation-GC retention relationships for aliphatic hydrocarbons and alkylbenzenes 322(1985)35
Saura-Calixto, F., see García-Raso, A. 302(1984)107
Sauter, A.
HPLC determination of adenine nucleotides in biological materials. Improvements and adaptations to routine analysis 325(1985)314
Sauvaire, D., see Cociglio, M. 307(1984)351
Savić, J., see Žunić, G. 311(1984)69
Savigliano, M., see Mentasti, E. 322(1985)177
Savolainen, H., see Rosenberg, C. 323(1985)429
Sawodny, W., see Grünes, R. 322(1985)63
Saxton, W.L.
Emergence temperatures as physical constants for measuring analyte retention in programmed temperature GC 312(1984)59
Schaber, P.M., Hunt, J.E., Fries, R. and Katz, J.J.
HPLC study of the chlorophyll allomerization reaction 316(1984)25
Schade, W. and Krainhöfner, A.
Determination of 9α-hydroxyandrost-4-en-3,17-dione in fermentation solutions by TLC (in German) 319(1985)475
Schäfer, K.H., see Levsen, K. 323(1985)135
Schaffner, C. and Giger, W.
Determination of nitrilotriacetic acid in water by high-resolution GC 312(1984)413
Schaik, A. van, see Van Hogezand, R.A. 305(1984)470
Schechter, P.J., see Grove, J. 306(1984)383
Schieffer, G.W.
Limitation of assessing HPLC peak purity with photodiode array detectors 319(1985)387
Schiffer, M., see Short, M.T. 323(1985)418
Schlicht, F., see De Vries, J.X. 309(1984)219
Schmelzeisen-Redeker, G., McDowall, M.A., Giessmann, U., Levsen, K. and Röllgen, F.W.
Studies with a laboratory-constructed thermospray liquid chromatograph-mass spectrometer interface 323(1985)127
Schmid, G., see Hamacher, K. 319(1985)311
Schmitter, J.M., see Ignatiadis, I. 324(1985)87
Schmitz, F.P., see Leyendecker, D. 315(1984)19, 321(1985)273
Schmitz, G. and Assmann, G.
Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65
Schmitz, G., Borgmann, U. and Assmann, G.
Analytical capillary isotachophoresis: a routine technique for the analysis of lipoproteins and lipoprotein subfractions in whole serum 320(1985)253
Schmitz, G., Lenczyk, M., Ord, D., Bowyer, D.E. and Assmann, G.
Quantitative densitometric method for the rapid separation and quantitation of the major lipids of tissues and lipoproteins by HPTLC. II. Reduction of the densitometric data 307(1984)81
Schmuckler, G.
Recent developments in ion chromatography (Review) 313(1984)47
Schmuckler, G., Rössner, B. and Schwedt, G.
Methods for the analysis of inorganic anions. IV. Reversed-phase HPLC with aqueous hydrophobic ion pairs as eluents 302(1984)15
Schneider, J.J., Triggs, E.J., Bourne, D.W.A., Stephens, I.D. and Haviland, A.M.
Determination of oxycodone in human plasma by HPLC with electrochemical detection 308(1984)359
Schoenmakers, P.J.
Thermodynamic model for supercritical fluid chromatography 315(1984)1
Schoffeniels, E., see Bontemps, J. 307(1984)283
Scholte, T.G., see Glöckner, G. 317(1984)615
Schöneshöfer, M., see Eibs, G. 310(1984)386
Schouten, J.P., see De Jong, J. 319(1985)23
Schrader, J., see Ontyd, J. 307(1984)404
Schreurs, W.H.P., see Speek, A.J. 301(1984)441, 305(1984)53
Schrijver, J., see Speek, A.J. 301(1984)441, 305(1984)53
Schrijver, J., see Tas, A.C. 310(1984)243
Schröder, W., see Svensson, P. 323(1985)363

Schroeder, J.S., see Albro, P.W. 312(1984)165
Schulten, H.-R., see Halket, J.M. 322(1985)200
Schumann, S., see Pötter, H. 319(1985)440
Schütz, H., see Mascher, H. 306(1984)231
Schwartz, H.A., see Yu, T.J. 301(1984)425
Schwartz, S.J., see Kobayashi, S. 319(1985)205
Schwarz, W., Langer, K. and Haag, A.
HPLC determination of (*Z*)- and (*E*)-urocanic acid in human skin 310(1984)188
Schwedt, G., see Schmuckler, G. 302(1984)15
Scott, B.S. and Dunn, D.L.
HPLC analysis of epinephrine enantiomers using a UV detector in series with an optical activity detector 319(1985)419
Scott, H.P., see Fell, A.F. 316(1984)423
Scott, H.P., see Knox, J.H. 316(1984)311
Secchi, C., see Biondi, P.A. 309(1984)151
Sedman, A.J. and Gal, J.
Simultaneous determination of the enantiomers of tocainide in blood plasma using GLC with electron-capture detection 306(1984)155
Sedman, A.J., see Gal, J. 314(1984)275
Seiber, J.N., see Kim, Y.-H. 314(1984)37
Sekharan, P. chandra, see Annapoorani, K.S. 303(1984)296
Seki, T. and Yamaguchi, Y.
New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188
Selden, C., Maton, P.N. and Chadwick, V.S.
Reversed-phase HPLC and separation of peptide hormones in human tissues and plasma with reference to cholecystokinins 305(1984)43
Selinger, Z., see Laufer, R. 301(1984)415
Senda, M., see Hoshino, T. 316(1984)473
Sentissi, A., see Nazareth, A. 314(1984)201, 314(1984)219
Seppälä, E., see Riutta, A. 307(1984)185
Serve, L., Combaut, G. and Piovetti, L.
Isolation, identification and determination of 3-(4-hydroxyphenyl)propionic acid by HPLC of the phenolic compounds of *Loiseleuria procumbens* Desf. (Ericaceae) (in French) 322(1985)257
Severin, G.
Rapid HPLC procedure for nitroglycerin and its degradation products 320(1985)445
Shader, R.I., see Abernethy, D.R. 307(1984)194
Shah, J.A. and Weber, D.J.
HPLC assay of pirlimycin in human serum and urine using 9-fluorenylmethylchloroformate 309(1984)95
Shaltiel, S., see Halperin, G. 317(1984)103
Shamoon, J.M. and Davies, I.A.I.
Separation of the enkephalins from proteins in an aqueous medium by chromatofocusing 314(1984)499
Shanbhag, V.D., see Bhatia, S.C. 305(1984)325
Shanks, C.A., see Avram, M.J. 306(1984)398
Sharifi, B.G., Bascom, C.C., Khurana, V.K. and Johnson, T.C.
Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173
Sharma, S.K. and Babitch, J.A.
Affinity chromatography of membrane fragments. Separation of carbonic anhydrase-enriched and -depleted brain membranes 312(1984)435
Sharp, D.E., see Mays, D.C. 311(1984)301
Shaw, I.C., see Sidau, B. 311(1984)234
Shaw, I.M., see Cookson, D.J. 312(1984)237
Shea, P.A. and Howell, J.B.
HPLC method for determining plasma and urine 3-methoxy-4-hydroxyphenylglycol by amperometric detection 306(1984)358
Shelly, D., see Gluckman, J. 317(1984)443
Shen, D.D., see Pollack, G.M. 311(1984)101
Shephard, G.S., see Labadarios, D. 309(1984)375, 310(1984)223
Shepherd, M.J., see Gilbert, J. 320(1985)361
Shiao, S.-Y., Johnson, J.S. Jr., Mohiuddin, G., Hata, W.Y., Tolan, J.S. and Doerr, W.W.
Ion-exchange properties of activated carbon filled with hydrous Zr(IV) and Zr(IV)-P(V) oxides 303(1984)13
Shih, F.F.
Analysis of glutamine, glutamic acid and pyroglutamic acid in protein hydrolysates by HPLC 322(1985)248
Shihabi, Z.K., see Powell, S.C. 317(1984)87
Shiino, M., see Kadowaki, H. 308(1984)329
Shimada, K., Tanaka, M. and Nambara, T.
Studies on steroids. CC. Determination of 17-ketosteroid sulphates in serum by HPLC with electrochemical detection using pre-column derivatization 307(1984)23
Shimada, K., see Ikeda, M. 305(1984)261
Shimamura, M., see Hirota, K. 319(1985)173
Shimizu, K., Amagaya, S. and Ogihara, Y.
Quantitative analysis of the metabolites of saikosaponin a using HPLC 307(1984)488
Shimono, H., see Miyazaki, K. 310(1984)219
Shimura, K. and Kasai, K.-I.
Introduction of ω-carboxyl spacers onto cross-linked agarose gel beads by O-alkylation for the preparation of affinity adsorbents 315(1984)161
Shinohara, Y., see Baba, S. 305(1984)119
Shiozaki, Y., see Tokuma, Y. 311(1984)339
Shirota, F.N., see Demaster, E.G. 308(1984)83
Shitashima, K., see Hayase, K. 322(1985)358
Shively, J.E., see Reeve, J.R. Jr. 321(1985)421
Shoji, S., Ohnishi, J., Funakoshi, T., Kubota, Y., Fukunaga, K., Miyamoto, E. and Ueki, H.
Simple and rapid identification of phosphorylated peptides from bovine brain myelin basic protein by reversed-phase HPLC 319(1985)359
Shono, T., see Funazo, K. 319(1985)143
Shono, T., see Tanaka, M. 301(1984)345, 314(1984)193
Short, M.T., Westholm, F.A., Schiffer, M. and Stevens, F.J.
Comparison of chromatographic

characteristics of a series of homologous Bence-Jones proteins during size-exclusion chromatography by HPLC and by Sephadex 323(1985)418
Shukla, U.A., Stetson, P.L. and Wagner, J.G. Sensitive GLC-electron-capture detection method for determination of sobrerol in biological fluids 308(1984)189
Sidau, B. and Shaw, I.C. Determination of sodium 2-mercaptoethanesulphonate by HPLC using post-column reaction colorimetry or electrochemical detection 311(1984)234
Siegel, S.A., Woods, K.L. and Dreyer, R.N. Pyrimidine nucleoside phosphorylase assay by automated HPLC 319(1985)351
Sienko, M., see Cantwell, A.M. 316(1984)133
Sievers, R.E., see Hutte, R.S. 302(1984)173
Sigvardson, K.W. and Birks, J.W. Detection of nitro-polycyclic aromatic hydrocarbons in LC by zinc reduction and peroxyoxalate chemiluminescence 316(1984)507
Simeón, S., see Villar, A. 303(1984)306
Simon, C.G., see Bidleman, T.F. 301(1984)448
Simon, R. and Wells, G. Bipolar pulsed microvolume electron-capture detector for capillary GC 302(1984)221
Simon, R.H., see Sura, D.J. 314(1984)471
Simonic, T., see Biondi, P.A. 309(1984)151
Simpson, R., see Gaffney, M.H. 306(1984)303
Šimuničová, E., see Zelenský, I. 325(1985)161
Sindhuphak, T., Patel, V. and Svensson, I. Temperature gradient chromatography of transfer RNA on hydroxyapatite 314(1984)408
Singh, A.K. Improved analysis of acephate and methamidophos in biological samples by selective ion monitoring GC-MS 301(1984)465
Singh, A.K., Zeleznikar, R.J. Jr. and Drewes, L.R. Analysis of soman and sarin in blood utilizing a sensitive GC-MS method 324(1985)163
Singh, H., see Slifkin, M.A. 303(1984)190
Sinsheimer, J.E., see Musial, S.P. 319(1985)467
Sinsheimer, J.E., see Van den Eeckhout, E. 318(1985)343
Sioufi, A. and Pommier, F. Determination of the two mononitrate metabolites of isosorbide dinitrate in human plasma and urine by GC with electron-capture detection 305(1984)95
Sisco, W.R. and DiFeo, T.J. Stability-indicating reversed-phase HPLC assay for fenretinide drug substances 322(1985)380
Sjöberg, A.-M. and Olkkonen, C. Determination of synthetic organic colours in lipsticks by TLC and HPLC 318(1985)149
Sjöberg, A.-M. and Pyysalo, H. Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90
Sjöström, E., see Alén, R. 301(1984)273
Sjövall, J., see Tandon, R. 302(1984)1
Skakum, W., see Carignan, G. 301(1984)292, 315(1984)470
Skarping, G., Smith, B.E.F. and Dalene, M. Trace analysis of amines and isocyanates using glass capillary GC and selective detection. III. Determination of aliphatic and alicyclic amines as perfluoro fatty acid amides using electron-capture and nitrogen-selective detection 303(1984)89
Skews, H., see Jackman, G.P. 308(1984)301
Skinner, S.J.M., Grego, B., Hearn, M.T.W. and Liggins, G.C. Separation of collagen α-chains by reversed-phase HPLC. Comparison of column alkyl stationary phases and temperature effects 308(1984)111
Skrabalak, D.S., Covey, T.R. and Henion, J.D. Qualitative detection of corticosteroids in equine biological fluids and the comparison of relative dexamethasone metabolite/dexamethasone concentration in equine urine by micro-LC-MS 315(1984)359
Skrinska, V. and Hahn, S. HPLC of 5-hydroxyindole-3-acetic acid in urine with direct sample injection 311(1984)380
Slatkavitz, K.J., Uden, P.C., Hoey, L.D. and Barnes, R.M. Atmospheric-pressure microwave-induced helium plasma spectroscopy for simultaneous multielement GC detection 302(1984)277
Slaughter, R.L., see Pollack, G.M. 311(1984)101
Ślebioda, M. and Kołodziejczyk, A.M. Prediction of retention data of protected dipeptides in normal phase HPLC 325(1985)282
Śledzińska, B. Investigation of conditions of the preparative LC separation of cardiac glycosides present in *Digitalis lanata* 303(1984)179
Slegers, G., see De Spiegeleer, B. 315(1984)481
Slezárik, A., see Markovič, O. 312(1984)492
Slifkin, M.A. and Singh, H. Masking effects in "charge-transfer" TLC 303(1984)190
Sloan, D.L., Ali, L.Z., Aybar-Batista, D., Yan, C. and Hess, S.L. Enzymatic assay procedures that employ HPLC: competition between phosphoribosyltransferases for a common substrate 316(1984)43
Sloan, D.L., see Ashton, R.W. 317(1984)245
Slováková, S., see Mikeš, O. 301(1984)93
Smail, M.C., see Hart, S.J. 306(1984)215
Smith, A.I. and McDermott, J.R. HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99
Smith, B.E., see Cookson, D.J. 312(1984)237

Smith, B.E.F., see Mellbin, G. 312(1984)203
Smith, B.E.F., see Peetre, I.-B. 318(1985)41
Smith, B.E.F., see Skarping, G. 303(1984)89
Smith, C.T., see Smith, G.A. 307(1984)410
Smith, G.A., Giacomini, K.M., Smith, C.T., Gregory, P.B., Robinson, W.S., Merigan, T.C. and Blaschke, T.F.
HPLC analysis of hypoxanthine arabinoside in plasma 307(1984)410
Smith, H.T. and Robinson, W.T.
Semi-automated HPLC method for the determination of cyclosporine in plasma and blood using column switching 305(1984)353
Smith, I.J., see Hartley, R. 306(1984)191
Smith, J.A., Ham, J., Winslow, D.P., O'Hare, M.J. and Rudland, P.S.
Use of HPLC in the isolation and characterisation of mouse and rat epidermal growth factors and examination of apparent heterogeneity 305(1984)295
Smith, L.L., see Teng, J.I. 322(1985)240
Smith, P.C. and Benet, L.Z.
HPLC method for the determination of indomethacin and its two primary metabolites in urine 306(1984)315
Smith, P.C., see Langendijk, P.N.J. 307(1984)371
Smith, R.D., see Wright, B.W. 314(1984)253
Smith, R.J., see Roets, E. 303(1984)117
Smith, R.M.
The use of retention indices to measure the solvent selectivity of ternary eluents in reversed-phase LC 324(1985)243
Smith, R.V., see Bauza, M.T. 310(1984)61
Smith, R.W., see Parker, C.E. 319(1985)273
Smith, S.R., see Hubball, J.A. 302(1984)341
Smrž, M., see Mikeš, O. 301(1984)93
Smyth, M.H., see Anders, J.C. 311(1984)117
Snellings, R.P.J., see Jansen, H. 325(1985)255
Snider, B.G., see Plaisted, S.M. 314(1984)369
Snider, B.G., see Theis, D.L. 321(1985)209
Snook, M.E., Mason, P.F., Arrendale, R.F. and Chortyk, O.T.
Capillary GC of dihydroxybenzoic, -phenylacetic and -phenylpropionic acids 324(1985)141
Snyder, K.P., see Hurst, W.J. 318(1985)408
Snyder, L.R., see Antle, P.E. 321(1985)1
Snyder, L.R., see Goldberg, A.P. 316(1984)241
Snyder, L.R., see Nazareth, A. 309(1984)357
Snyder, L.R., see Weiser, E.L. 303(1984)1
Soják, L., Kraus, G. and Ostrovský, I.
High-performance GC on liquid crystal capillaries. VIII. Separation of alkylbenzenes up to C_9 (in German) 323(1985)414
Soldin, S.J., see Tan, Y.K. 311(1984)311
Solinas, V., see Deiana, S. 320(1985)450
Sollenberg, J., see Hansén, L. 312(1984)489
Sonnet, P.E. and Heath, R.R.
Aryl substituted diastereomeric alkenes: GC behavior on a non-polar *versus* a liquid crystal phase 321(1985)127
Sonnini, S., see Gazzotti, G. 315(1984)395
Soriano, F., see Méndez, E. 323(1985)373
Soula, G., see Bellon, P. 309(1984)170
Souppart, C., see Lecaillon, J.-B. 317(1984)493
Souteyrand, C., Thibert, M., Caude, M. and Rosset, R.
Influence of water in LSC systems on retention data. II. Fixation of monolocalized solutes with class N solvents and silica gels 316(1984)373
Sova, O.
Autofocusing: a method for isoelectric focusing without carrier ampholytes 320(1985)15
Sova, O.
Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213
Sowin, T.J., see Pirkle, W.H. 319(1985)392
Spahn, H., Weber, H., Mutschler, E. and Möhrke, W.
α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167
Spahn, H., see Weber, H. 307(1984)145
Spanjer, M.C., see De Ligny, C.L. 301(1984)311
Speek, A.J., Schrijver, J. and Schreurs, W.H.P.
Fluorimetric determination of menadione sodium bisulphite (vitamin K_3) in animal feed and premixes by HPLC with post-column derivatization 301(1984)441
Speek, A.J., Schrijver, J. and Schreurs, W.H.P.
Fluorometric determination of total vitamin C in whole blood by HPLC with pre-column derivatization 305(1984)53
Speranza, M., see Felici, M. 320(1985)435
Spiegeleer, B. de, see De Spiegeleer, B. 315(1984)481
Spiteller, G., see Bauer, S. 309(1984)243
Spiteller, G., see Thoma, H. 309(1984)17
Spittler, T.D., Marafioti, R.A. and Lahr, L.M.
Determination of benomyl and its metabolites by cation-exchange HPLC 317(1984)527
Sportoletti, G., see Cremonesi, P. 308(1984)310
Sprissler, B. and Lockwood, F.E.
Analysis of mechanical lubricant contaminants in aluminium cold rolling oil by reversed-phase LC 319(1985)222
Sprung, C., see Rosenfeld, J.M. 325(1985)309
Sreenivasan, K.
Determination of polymer molecular weights by PC 325(1985)363
Staehelin, J., see Hutte, R.S. 302(1984)173
Stafström, N., see Johansson, B.-L. 314(1984)396
Stahel, O., see Little, C.J. 316(1984)105
Staibano, G., see Marzo, A. 310(1984)51
Staiger, C., see De Vries, J.X. 309(1984)219
Stampfli, H.F., Lai, C.-M., Yacobi, A. and Sum, C.Y.
HPLC assay for the major blood metabolite of esmolol: an ultra short acting beta blocker 309(1984)203
Stan, H.-J. and Goebel, H.
Evaluation of automated splitless and

manual on-column injection techniques using capillary GC for pesticide residue analysis 314(1984)413
Stanborough, M.S., see Harvey, T.G. 319(1985)230
Stancher, B. and Zonta, F.
Quantitative HPLC method for determining the isomer distribution of retinols (vitamin A_1) and 3-dehydroretinol (vitamin A_2) in fish oils 312(1984)423
Stancher, B., see Zonta, F. 301(1984)65
Standard, S.A., Vaux, P. and Bray, C.M.
HPLC of nucleotides and nucleotide sugars extracted from wheat embryo and vegetable seed 318(1985)433
Stanimirović, S., see Žunić, G. 311(1984)69
Stark, B.C. and Chooi, W.Y.
Affinity chromatography of *Drosophila melanogaster* ribosomal proteins to 5S rRNA 319(1985)367
Starratt, A.N., see Lee, T.T. 325(1985)340
Startin, J.R., see Gilbert, J. 319(1985)376
Staszewski, R., see Wardencki, W. 325(1985)430
Stavric, B., Klassen, R. and Gilbert, S.G.
Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107
Steele, D., see Timmons, L. 314(1984)476
Steffen, J.M., see Feldhoff, R.C. 311(1984)267
Steffen, M.C., see Feldhoff, R.C. 311(1984)267
Steinauer, R., Chen, F.M.F. and Benoiton, N.L.
Determination of epimeric peptides for assessing enantiomeric purity of starting materials and studying racemization in peptide synthesis using HPLC 325(1985)111
Steinbach, E., see König, W.A. 301(1984)129
Steinmeyer, S., see Tiebach, R. 318(1985)103
Stejskal, J., see Kovář, J. 325(1985)359
Stenhagen, G., see Alborn, H. 323(1985)47
Stephany, R.W., see Jansen, E.H.J.M. 319(1985)436
Stephens, I.D., see Schneider, J.J. 308(1984)359
Stermer-Cox, M.G., see Brown, N.D. 309(1984)426, 315(1984)389
Sternson, L.A., see Meltzer, N.M. 307(1984)216
Stetson, P.L., see Shukla, U.A. 308(1984)189
Stevens, F.J., see Short, M.T. 323(1985)418
Stevens, R.G., see Kamblawi, M.O. 309(1984)431
Stewart, J.M., see Gallagher, R.T. 321(1985)217
Stickney, D., see Ryan, P.J. 320(1985)421
Stickney, T.M., see Haky, J.E. 321(1985)137
Stikkelman, R.M., Tjioe, T.T., Van der Wiel, J.P. and Van Rantwijk, F.
HPLC separation of glucose-1-phosphate, fructose, sucrose and inorganic orthophosphate 322(1985)220
Stitzel, R.E., see Barone, S. 310(1984)283
Stoll, A.L., Baldessarini, R.J., Cohen, B.M. and Finklestein, S.P.
Assay of plasma thioridazine and metabolites by HPLC with amperometric detection 307(1984)457
Stoner, E., see Loche, S. 317(1984)377
Stover, F.S.
Enhancing isotachophoresis sensitivity by low-concentration electrolyte cascading 320(1985)45
Stowe, M., see Trumbore, C.N. 322(1985)443
Strang, C., Levine, S.P., Orlan, B.P., Gouda, T.A. and Saner, W.A.
High resolution GC analysis of cytochrome P-448 inducing polychlorinated biphenyl congeners in hazardous waste 314(1984)482
Stránský, Z.
Isotachophoresis of cationic herbicides in waters and soils 320(1985)219
Stratton, W.J., see Kirkman, C.M. 317(1984)569
Strauss, H.C., see Luzzi, F.A. 311(1984)291
Strife, R.J. and Murphy, R.C.
Preparation of pentafluorobenzyl esters of arachidonic acid lipoxygenase metabolites. Analysis by GC and negative-ion chemical ionization MS 305(1984)3
Strife, R.J., see Keough, T. 312(1984)450
Štrop, P., see Mikeš, O. 301(1984)93
Stuart, J.D., see Huang, J.-X. 316(1984)151
Stuart, J.F.B., see Cummings, J. 311(1984)125
Štulík, K., see Kysilka, R. 320(1985)414
Stull, R., see Goldstein, D.S. 311(1984)148
Stump, D.D., Roth, E.F. Jr. and Gilbert, H.S.
Simultaneous determination by HPLC of tocopherol isomers, α-tocopheryl quinone, and cholesterol in red blood cells and plasma 306(1984)371
Sturani, C., see Grossi, G. 309(1984)214
Stutte, C.A., see Jones, R.S. 319(1985)454
Styring, M.G., Price, C. and Booth, C.
Effect of solute size and shape on the elution volume in size-exclusion chromatography 319(1985)115
Su, H. and Bernauer, K.
Quantitation of the enantiomers of β-hydroxyamino acids by GC resolution on an optically active packed column 301(1984)461
Sudduth, R.D., see Timm, D.C. 316(1984)343
Sudraud, G., see Olek, M. 325(1985)239
Suemitsu, R., Yamada, Y., Sano, T. and Kitayama, M.
Analysis of naturally occurring reduced anthraquinones obtained from *Alternaria porri* (Ellis) Ciferri by HPLC 318(1985)139
Sugihara, J., see Imamura, T. 305(1984)456
Sugimae, A., see Hattori, Y. 315(1984)341
Sugimoto, I., see Kakehi, K. 318(1985)367
Sugimoto, Y., see Yamada, J. 311(1984)385
Sugishita, E., see Amagaya, S. 320(1985)430
Sugita, N., see Dohji, T. 311(1984)249
Sulaiman, S.M., see Ali, H.M. 305(1984)465
Sullivan, G.M., see Allgire, J.F. 325(1985)249
Sum, C.Y., see Stampfli, H.F. 309(1984)203
Sun, W.F., see Lawrence, J.F. 325(1985)299
Sunaga, H., see Suzuki, K.T. 303(1984)131
Sunde, A., see Lundmo, P.I. 308(1984)289
Suñol, C., see Tusell, J.M. 306(1984)338
Suortti, T.
Improved analytical and preparative

methods for necatorin from *Lactarius necator* (Fr.) Karst mushroom 301(1984)303
Sura, D.J., Simon, R.H. and Beyerlein, F.H.
Simultaneous determination of pynamin forte, resmethrin and piperonyl butoxide in insecticide formulations by GC 314(1984)471
Sutherland, D.B., see Hunzicker-Dunn, M. 312(1984)293
Suzuki, A., see Crétin, C. 315(1984)430
Suzuki, K.T., Sunaga, H. and Yajima, T.
Separation of metallothionein into isoforms by column switching on gel permeation and ion-exchange columns with HPLC-atomic-absorption spectrophotometry 303(1984)131
Suzuki, M.
Selected ion monitoring MS method for the determination of hexachlorocyclohexane isomers in soils 312(1984)403
Suzuki, M., see Mori, S. 320(1985)343
Suzuki, M., see Oka, H. 314(1984)303, 325(1985)265
Svec, F., see Hradil, J. 324(1985)277
Svennerholm, L., see Månsson, J.-E. 322(1985)465
Svensson, I., see Sindhuphak, T. 314(1984)408
Svensson, P., Schröder, W., Åkerlund, H.-E. and Albertsson, P.-Å.
Fractionation of thylakoid membrane components by extraction in aqueous polymer two-phase systems containing detergent 323(1985)363
Sweeney, A.G. and Cheh, A.M.
Production of mutagenic artifacts by the action of residual chlorine on XAD-4 resin 325(1985)95
Sweetman, B.J., see Wedlund, P.J. 307(1984)121
Sweetman, J.A., see Tong, H.Y. 312(1984)183
Szefler, S.J., see Ebling, W.F. 305(1984)271
Szepesi, G., see Gazdag, M. 316(1984)267
Szórádi, S., see Körtvélyessy, G. 303(1984)370
Sztruhar, I., see Körtvélyessy, G. 303(1984)370
Szücs, L., see Niegreisz, Z. 316(1984)451
Szulc, J., see Witkiewicz, Z. 312(1984)141, 315(1984)145
Szymczyński, G.A., see Waliszewski, S.M. 321(1985)480
Szymoniak, K., see Chrétien, J.R. 324(1985)355

T

Táborský, J., see Horna, A. 325(1985)367
Tachibana, Y., see Mizutani, T. 324(1985)480
Tackett, R., see Rydzik, R. 308(1984)31
Taga, M., see Yoshida, H. 325(1985)179
Takagi, N., see Tsuchiya, H. 309(1984)43
Takahama, K., see Okano, Y. 310(1984)251
Takahashi, N., Takahashi, Y., Ortel, T.L., Lozier, J.N., Ishioka, N. and Putnam, F.W.
Purification of glycopeptides of human plasma proteins by HPLC 317(1984)11
Takahashi, Y., see Takahashi, N. 317(1984)11
Takami, K., see Hattori, Y. 315(1984)341
Takehana, Y., see Konomi, H. 311(1984)375
Takemi, H., see Hirokawa, T. 305(1984)429
Takemoto, S., see Hattori, Y. 315(1984)341
Takeuchi, T., Jin, Y. and Ishii, D.
Development of an on-line high-pressure pre-column concentration method for micro HPLC and its application to the rapid determination of dibutyl phthalate in water 321(1985)159
Takeuchi, T., see Ishii, D. 316(1984)441
Takeuchi, T., see Wada, H. 320(1985)369
Takiyama, R., see Hirokawa, T. 305(1984)429
Talha, M.R.Z. and Rogers, H.J.
Rapid GC determination of ifosfamide in biological fluids 311(1984)194
Tam, Y.K., Ferguson, S.M., Yau, M.L. and Wyse, D.G.
Simple and rapid HPLC method for the analysis of sulfinpyrazone and four of its metabolites in human plasma 310(1984)438
Tamchyna, J., Zuska, J. and Vacík, J.
The performance of the UVM-4 UV analyser in isotachophoresis 320(1985)241
Tameesh, A.H.H., Bender, A.O. and Sarkissian, T.M.
GC study of the analysis and elution mechanism of hydrogen sulphide, carbonyl sulphide and light mercaptans in petroleum gases 321(1985)59
Tamma, R.V., Miller, G.C. and Everett, R.
HPLC analysis of coumarins and flavonoids from section Tridentatae of *Artemisia* 322(1985)236
Tammilehto, S.A.
HPTLC determination of trimethoprim and sulphamethoxazole in pharmaceutical dosage forms 323(1985)456
Tamminga, C.A., see Knight, M. 301(1984)277
Tamura, Z., see Uchino, K. 308(1984)241
Tamura, Z., see Yoshioka, M. 309(1984)63
Tan, Y., see Koopmans, P.P. 307(1984)445
Tan, Y.K. and Soldin, S.J.
Determination of salbutamol in human serum by reversed-phase HPLC with amperometric detection 311(1984)311
Tanabe, S., see Toida, T. 308(1984)133
Tanaka, A., Nose, N., Masaki, H., Iwasaki, H. and Haruta, M.
Determination of the stable isotope of nitrite flux in the blood of mice by GC-MS with selected ion monitoring 306(1984)51
Tanaka, E., Hagino, S., Yoshida, T. and Kuroiwa, Y.
Simultaneous determination of trimethadione and its metabolite in rat and human serum by HPLC 308(1984)393
Tanaka, H., see Hayakawa, K. 312(1984)476
Tanaka, M., Kawaguchi, Y., Niinae, T. and Shono, T.
Preparation and retention behaviour of

chemically bonded methylated-cyclodextrin stationary phases for LC 314(1984)193
Tanaka, M., Kawaguchi, Y., Shono, T., Uebori, M. and Kuge, Y.
Unmodified and acylated cyclodextrin stationary phases for LC separation of aromatic compounds 301(1984)345
Tanaka, M., see Funazo, K. 319(1985)143
Tanaka, M., see Shimada, K. 307(1984)23
Tanaka, N., see Uchino, K. 308(1984)241
Tandon, R., Axelson, M. and Sjövall, J.
Selective LC isolation and GC-MS analysis of ketonic bile acids in faeces 302(1984)1
Taneja, A.K., Lau, S.Y.M. and Hodges, R.S.
Separation of hydrophobic peptide polymers by size-exclusion and reversed-phase HPLC 317(1984)1
Taneja, A.K., see Lau, S.Y.M. 317(1984)129
Tangerås, A.
Separation of haem compounds by reversed-phase ion-pair HPLC and its application in the assay of ferrochelatase activity 310(1984)31
Tangerman, A., see Van Hogezand, R.A. 305(1984)470
Tanizawa, O., see Dohji, T. 311(1984)249
Tarbet, B.J., see Lee, M.L. 302(1984)303
Tarter, J.G., see Jones, V.K. 312(1984)456
Tarutani, T.
Chromatographic studies in the field of silicate chemistry (Review) 313(1984)33
Tas, A.C., Odink, J., Noever de Brauw, M.C. ten, Schrijver, J. and Jonk, R.J.G.
Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243
Tas, A.C., see Van der Greef, J. 323(1985)81
Tatsumi, M., see Tsuchiya, H. 309(1984)43
Tauber-Finkelstein, M., see Halperin, G. 317(1984)103
Tayar, N. el, see El Tayar, N. 320(1985)293, 320(1985)305
Taylor, N.F., see Kwan, T.K. 301(1984)189
Taylor, R.B., Reid, R. and Hung, C.T.
Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279
Taylor, R.F. and Marenchic, I.G.
Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193
Taylor, S.L., see Hui, J.Y. 312(1984)443
Teja, E. de la, see Dulitzky, M. 317(1984)403
Teller, D.C., see Sasagawa, T. 307(1984)29
Templeton, J.L., see Bardalaye, P.C. 314(1984)450
Tenaschuk, D., see MacKenzie, S.L. 319(1985)404, 322(1985)228
Teng, J.I. and Smith, L.L.
Improved HPLC of sphingomyelin 322(1985)240
Terada, H., Asanoma, M. and Sakabe, Y.
Studies on residual antibacterials in foods. III. HPLC determination of penicillin G in animal tissues using an on-line pre-column concentration and purification system 318(1985)299
Terada, M.
Determination of methamphetamine and its metabolites in rat tissues by GC with a nitrogen-phosphorus detector 318(1985)307
Terragno, A., see Rydzik, R. 308(1984)31
Terry, M.D., Hisayasu, G.H., Kern, J.W. and Cohen, J.L.
HPLC analysis of naloxone in human serum 311(1984)213
Tešić, Ž.L., see Vučković, G. 324(1985)431
Testa, B., see El Tayar, N. 320(1985)293, 320(1985)305
Thakur, S., see Ray, B. 315(1984)401
Theis, D.L., Rusk, M.L., Plaisted, S.M. and Snider, B.G.
HPLC method for the separation of isomers of 9β-methylcarbacyclin (ciprostene) and the analysis of pharmaceutical samples 321(1985)209
Thenot, J.P., see Padovani, P. 308(1984)229
Theoharides, A.D., see Anders, J.C. 311(1984)117
Theumer, J., see Ostrowksi, J. 309(1984)115
Thibert, M., see Souteyrand, C. 316(1984)373
Thiel, D., see Frank, H. 309(1984)261
Thieme, R.E., see Eichholtz, P.C.N. 305(1984)438
Thiessen, J.J., see Yip, M.S. 307(1984)343
Thoma, H., Reiner, J. and Spiteller, G.
Profiles of strongly polar and less polar acids obtained from human blood, plasma and serum by two-step ultrafiltration 309(1984)17
Thomas, E.W.
HPLC determination of a new calcium antagonist, fostedil, in plasma and urine using fluorescence detection 305(1984)233
Thomas, L.C., see Adams, A.K. 303(1984)341
Thomas, L.C., see Ramus, T.L. 314(1984)243
Thomas, L.M., see Anders, J.C. 311(1984)117
Thomas, R.W., see Nora, M.V. 307(1984)451
Thompson, A.M., see Phillips, T.M. 317(1984)173
Thompson, H.J. and Brown, S.A.
Separations of some coumarins of higher plants by LC 314(1984)323
Thompson, J.R., see Early, R.J. 310(1984)1
Thompson, R.E., see Kester, A.S. 310(1984)372
Thompson, T.A., see Dehaas Vermeulen, J. 306(1984)412
Thormann, W.
Description and detection of moving sample zones in zone electrophoresis: Zone spreading due to the sample as a necessary discontinuous element 318(1985)455
Thormann, W., see Mosher, R.A. 320(1985)23
Thorsrud, A.K., see Tong, H.Y. 312(1984)183
Thus, J.L.G. and Kraak, J.C.
Comparison of phenyl- and octadecyl-modified silica gel as stationary phase for

the prediction of *n*-octanol-water partition coefficients by HPLC 320(1985)271
Thyssen, K.
Determination of chloroacetanilides and chloroanilines in sulphanilamides and waste water by HPLC 319(1985)99
Tian, W.M., see Timm, D.C. 316(1984)343
Tibbetts, D., see Biffar, S. 318(1985)404
Tibbetts, D.L., see Cohn, L.J. 321(1985)401
Tiebach, R., Blaas, W. and Kellert, M.
Design and construction of an interface for direct liquid introduction coupling of a microbore high-performance liquid chromatograph to a quadrupole mass spectrometer 323(1985)121
Tiebach, R., Blaas, W., Kellert, M., Steinmeyer, S. and Weber, R.
Confirmation of nivalenol and deoxynivalenol by on-line LC-MS and GC-MS. Comparison of methods 318(1985)103
Tietz, A., see Tietz, D. 325(1985)425
Tietz, D. and Tietz, A.
Less hazardous derivatization procedure for GC of plant hormone abscisic acid 325(1985)425
Tiffany, J.M., see Harvey, D.J. 301(1984)173
Timm, D.C., Tian, W.M., Larson, B.J. and Sudduth, R.D.
Thermoset resin microstructure by GPC: an anhydride-cured epoxy 316(1984)343
Timmons, L., Steele, D., Cannon, M., Grese, R., Brown, R., Murrill, E. and Jameson, C.W.
Identification of bromotetrachlorophenol in commercial pentachlorophenol samples 314(1984)476
Tippins, B.L., see Said, H.M. 324(1985)65
Tirronen, H., see Herranen, J. 307(1984)241
Titani, K., see Sasagawa, T. 307(1984)29
Tjaden, U.R., see De Jong, J. 319(1985)23, 322(1985)43
Tjaden, U.R., see Langenberg, J.P. 305(1984)61
Tjioe, T.T., see Stikkelman, R.M. 322(1985)220
To, E.C.A. and Wells, P.G.
Rapid and sensitive assays using HPLC to measure the actitivies of phase II drug metabolising enzymes: glucuronyl transferase and sulfotransferase 301(1984)282
Todd, R.A., see Brazell, R.S. 302(1984)257
Todoriki, H., Hayashi, T. and Naruse, H.
HPLC method for screening disorders of aromatic acid metabolism using a multi-detection system 310(1984)273
Togari, A., see Oka, K. 308(1984)43
Togawa, T., see Toida, T. 308(1984)133
Tohei, S., see Hoshino, T. 316(1984)473
Toida, T., Togawa, T., Tanabe, S. and Imanari, T.
Determination of cyanide and thiocyanate in blood plasma and red cells by HPLC with fluorometric detection 308(1984)133
Tokuda, M., Kamei, M., Yui, S. and Koyama, F.
Rapid resolution of nucleotide sugars by lectin affinity HPLC 323(1985)434
Tokuma, Y., Shiozaki, Y. and Noguchi, H.
Determination of a new orally active cephalosporin in human plasma and urine by HPLC using automated column switching 311(1984)339
Tolan, J.S., see Shiao, S.-Y. 303(1984)13
Toll, P.J.M.M., see Visser, T. 309(1984)81
Tolman, V., Vlasáková, V. and Živný, K.
Simple method for the separation of diastereomers of 4-fluoroglutamic acid by GLC 315(1984)421
Tomás, F., see Barberán, F.A.T. 315(1984)101
Tomkins, B.A., see Maskarinec, M.P. 302(1984)51
Tonelli, D., see Gagliardi, L. 315(1984)465, 325(1985)353
Tong, H.Y., Centen, J.D., Karasek, F.W., Jellum, E. and Helland, P.
Identification of trace organic compounds in dimethyl sulphoxide solution using HPLC and GC-MS 324(1985)373
Tong, H.Y., Sweetman, J.A., Karasek, F.W., Jellum, E. and Thorsrud, A.K.
Quantitative analysis of polycyclic aromatic compounds in diesel exhaust particulate extracts by combined chromatographic techniques 312(1984)183
Tongeren, J.H.M. van, see Van Hogezand, R.A. 305(1984)470
Toppet, S., see Roets, E. 303(1984)117
Torel, J., Cillard, J., Cillard, P. and Vie, M.
Simultaneous analysis of three antimicrobial agents in feed premixes by reversed-phase HPLC 323(1985)447
Toren, P.C., Plakunov, I., Peyton, A.L., Cooper, C.S. and Weinkam, R.J.
Quantitation of methanol formed in cell culture cytotoxicity assays and as a metabolite in microsome suspensions 305(1984)171
Tornheim, P.A., see Allen, K.G. 309(1984)33
Torres, A.R., Dunn, B.E., Edberg, S.C. and Peterson, E.A.
Preparative HPLC of proteins on ion-exchange columns with carboxymethyldextrans as displacers 316(1984)125
Toselli, A., see Bernareggi, A. 309(1984)415
Tóth, K., see Niegreisz, Z. 316(1984)451
Tourville, J.F., see Narang, P.K. 305(1984)135
Toutain, P.L., see Alvinerie, M. 309(1984)385
Touzet, C., see Lartigue-Mattei, C. 310(1984)407
Toyama, A., see Nimura, N. 316(1984)547
Traitler, H., see Kolarovic, L. 314(1984)233
Treffner, E., see Marzo, A. 310(1984)51
Trenk, D., see Bechtold, H. 306(1984)333
Triggs, E.J., see Schneider, J.J. 308(1984)359
Tripathi, D.N., Malhotra, R.C. and Bhattacharya, A.
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b*,*f*]-1:4-oxazepine 315(1984)417

Troitsky, G.V. and Azhitsky, G.Y.
Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285
Trojánek, A.
New, modified flow-through polarographic detector 323(1985)406
Trojanowicz, M., see Haddad, P.R. 315(1984)261, 321(1985)363, 324(1985)319
Trumbore, C.N., Grehlinger, M., Stowe, M. and Kelleher, F.M.
Further experiments on a new, fast method for determining molecular weights of diffusing species in a liquid phase 322(1985)443
Tsao, C.S., Young, M. and Rose, S.M.
Quantitative analysis of ascorbic acid-2-sulfate by HPLC with electrochemical detection 308(1984)306
Tshabalala, M.A., see Parekh, N.J. 314(1984)65
Tsimidou, M., see Robinson, J.L. 324(1985)35
Tsipouras, A., see Pirkle, W.H. 319(1985)392
Tsubota, H., see Hayase, K. 322(1985)358
Tsuchiya, H., Hayashi, T., Sato, M., Tatsumi, M. and Takagi, N.
Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 309(1984)43
Tsuruta, Y., Kohashi, K., Ishida, S. and Ohkura, Y.
Determination of nicotinic acid in serum by HPLC with fluorescence detection 309(1984)309
Tsuruta, Y., see Nakano, Y. 311(1984)390
Tuchek, J.M., see Wilson, T.W. 306(1984)351
Tuchman, M. and Krivit, W.
Capillary GC investigation of plasma lipid alcoholysis during alcohol extraction 307(1984)172
Turánek, J., see Kovář, J. 319(1985)341
Turino, G.M., see Keller, S. 305(1984)461
Tusell, J.M., Artigas, F., Suñol, C., Martínez, E. and Gelpí, E.
Comparison of HPLC and GC-MS for the analysis of indole-3-acetic acid in brain tissue 306(1984)338
Tychopoulos, V., see Tyman, J.H.P. 303(1984)137
Tyman, J.H.P., Tychopoulos, V. and Chan, P.
Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

U

Uchida, H., see Ishida, R. 305(1984)73
Uchida, K., see Imaizumi, N. 308(1984)295
Uchino, K., Isozaki, S., Saitoh, Y., Nakagawa, F., Tamura, Z. and Tanaka, N.
Quantitative determination of furosemide in plasma, plasma water, urine and ascites fluid by HPLC 308(1984)241
Uden, P.C.
Inorganic GC (Review) 313(1984)3
Uden, P.C., see Kirkman, C.M. 317(1984)569
Uden, P.C., see Limentani, G.B. 325(1985)53
Uden, P.C., see Slatkavitz, K.J. 302(1984)277
Udseth, H.R., see Wright, B.W. 314(1984)253
Uebori, M., see Tanaka, M. 301(1984)345
Ueki, H., see Shoji, S. 319(1985)359
Uemura, I., see Kadowaki, H. 308(1984)329
Ujimoto, K. and Kurihara, H.
Evaluation of hydrophobicity of gels by use of some 1-alkanols as pilot solutes 301(1984)57
Ultsch, I., see Begutter, H. 321(1985)475
Unger, K.K., see Kinkel, J.N. 316(1984)193
Unger, T., see Hermann, K. 312(1984)273
Uno, K., see Oka, H. 325(1985)265
Uray, G., see Lindner, W. 316(1984)605
Utamura, T., see Koizumi, K. 321(1985)145
Utoh, M., see Baba, S. 307(1984)1

V

Vacík, J. and Muselasová, I.
Isotachophoretic determination of anionic and cationic species in well and surface waters 320(1985)199
Vacík, J., Zuska, J. and Muselasová, I.
Improvement of the performance of a high-frequency contactless conductivity detector for isotachophoresis 320(1985)233
Vacík, J., see Fidler, V. 320(1985)167
Vacík, J., see Fidler, Z. 320(1985)175
Vacík, J., see Hanibalová, D. 320(1985)185
Vacík, J., see Kašička, V. 320(1985)33
Vacík, J., see Tamchyna, J. 320(1985)241
Vaggi, L., see Forchetti, C. 309(1984)177
Vajta, S., Le Moing, J.P. and Rovei, V.
Reversed-phase HPLC separation of ^{14}C-labelled toloxatone and its metabolites 311(1984)329
Vala, V., see Kovář, J. 319(1985)341
Valencia, R., N'Guyen Công, H. and Bertaux, O.
Separation of naturally occurring or

induced methylated nucleobases of DNA by reversed-phase HPLC 325(1985)207
Valette, J.-P., see Nicot, G. 306(1984)279
Valkenburg, C.A., see Grimsrud. E.P. 302(1984)243
Valkenburg, C.F.M. van, see De Jong, J. 322(1985)43
Valkó, K., Papp, O. and Darvas, F. Selection of GC stationary phase pairs for characterization of the 1-octanol-water partition coefficient 301(1984)355
Van Amsterdam, P.H., see Ruijten, H.M. 314(1984)183
Van Ardenne, R.A.M., see De Raat, W.K. 310(1984)41
Van Balen, H.C.J.G., see Van Hogezand, R.A. 305(1984)470
Van Blitterswijk, H., see Jansen, E.H.J.M. 319(1985)436
Van Brummelen, P., see Kleinbloesem, C.H. 308(1984)209
Van Cutsem, J.-L., see Bontemps, J. 311(1984)59
Van de Waterbeemd, H., see El Tayar, N. 320(1985)293, 320(1985)305
Van den Berg, J.H.M., see Glöckner, G. 317(1984)615
Van den Bossche, W., see De Croo, F. 325(1985)395
Van den Bossche, W., see De Spiegeleer, B. 315(1984)481
Van den Eeckhout, E., De Moerloose, P. and Sinsheimer, J.E. HPTLC assay for epoxide hydrolase activity and the determination of phenoxypropane-1,2-diols 318(1985)343
Van der Greef, J., Tas, A.C., Noever de Brauw, M.C. ten, Höhn, M., Meijerhoff, G. and Rapp, U. Performance of a moving belt LC-MS interface 323(1985)81
Van der Helm, D., see Jalal, M.A.F. 301(1984)247
Van der Kleijn, E., see Oosterbaan, M.J.M. 306(1984)323
Van der Vijgh, W.J.F., Elferink, F., Postma, G.J., Vermorken, J.B. and Pinedo, H.M. Determination of ethylenediamineplatinum(II) malonate in infusion fluids, human plasma and urine by HPLC 310(1984)335
Van der Vijgh, W.J.F., see Elferink, F. 320(1985)379
Van der Vijgh, W.J.F., see Gooijer, C. 312(1984)337
Van der Wiel, J.P., see Stikkelman, R.M. 322(1985)220
Van der Zee, R. and Welling, G.W. Microbore reversed-phase chromatography of proteins with conventional gradient equipment for HPLC 325(1985)187
Van Engelen, D.L., see Adams, A.K. 303(1984)341
Van Ginneken, C.A.M., see Koopmans, P.P. 307(1984)445
Van Harten, J., see Kleinbloesem, C.H. 308(1984)209
Van Hees, P.A.M., see Van Hogezand, R.A. 305(1984)470
Van Hogezand, R.A., Van Balen, H.C.J.G., Van Schaik, A., Tangerman, A., Van Hees, P.A.M., Zwanenburg, B. and Van Tongeren, J.H.M. Determination of sodium azodisalicylate, salazosulphapyridine and their metabolites in serum, urine and faeces by HPLC 305(1984)470
Van Houwelingen, J.C, see De Ligny, C.L. 301(1984)311
Van Lenten, F.J., see Buchta, R.C. 325(1985)456
Van Ooijen, R.D., see Drost, R.H. 310(1984)193
Van Oort, W.J., see Den Hartigh, J. 306(1984)444
Van Rantwijk, F., see Stikkelman, R.M. 322(1985)220
Van Schaik, A., see Van Hogezand, R.A. 305(1984)470
Van Tongeren, J.H.M., see Van Hogezand, R.A. 305(1984)470
Van Valkenburg, C.F.M., see De Jong, J. 322(1985)43
Vanazzi, G., see Beretta, E. 308(1984)341
Vanderhaeghe, H., see De Pourcq, P. 321(1985)441
Vanderhaeghe, H., see Dihuidi, K. 325(1985)413
Vanderhaeghe, H., see Roets, E. 303(1984)117
Vandervennet, R., see Poole, C.F. 302(1984)149
Vandewoude, M., Claeys, M. and De Leeuw, I. Determination of α-tocopherol in human plasma by HPLC with electrochemical detection 311(1984)176
Vanhaelen, M. and Vanhaelen-Fastré, R. Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497
Vanhaelen, M., see Duez, P. 303(1984)391
Vanhaelen, M., see Vanhaelen-Fastré, R. 312(1984)467
Vanhaelen-Fastré, R. and Vanhaelen, M. Qualitative and quantitative determinations of hallucinogenic components of psilocybe mushrooms by reversed-phase HPLC 312(1984)467
Vanhaelen-Fastré, R., see Duez, P. 303(1984)391
Vanhaelen-Fastré, R., see Vanhaelen, M. 312(1984)497
Vapaatalo, H., see Riutta, A. 307(1984)185
Várady, L., see Nagy, J. 317(1984)165
Varhegyi, G. and Alexander, G. Use of histograms in computer-aided comparison of chromatograms 318(1985)247
Vaux, P., see Standard, S.A. 318(1985)433

Veenendaal, J.R. and Meffin, P.J.
Direct analysis of diflunisal ester and ether glucuronides by HPLC 307(1984)432
Vejrosta, J., Mikešová, M. and Novák, J.
Rôle of displacement effects in trapping trace components from gases in sorbent-packed columns 324(1985)269
Veleiro, A.S., see Burton, G. 315(1984)435
Velthorst, N.H., see Gooijer, C. 312(1984)337
Veltkamp, A.C., see Gooijer, C. 312(1984)337
Venard, P., see Abad, P. 318(1985)417
Vercruysse, S.A.R., Delcour, J.A. and Dondeyne, P.
Isolation of quercetin, myricetin, and their respective dihydro-compounds by Sephadex LH-20 chromatography 324(1985)495
Verheggen, T.P.E.M., Everaerts, F.M. and Reijenga, J.C.
UV detection at 206 nm in isotachophoresis 320(1985)99
Verheggen, T.P.E.M., see De Bruijn, C.H.M.M. 320(1985)205
Verheggen, T.P.E.M., see Everaerts, F.M. 320(1985)263
Verheggen, T.P.E.M., see Reijenga, J.C. 320(1985)67, 320(1985)245
Vermeulen, J. dehaas, see Dehaas Vermeulen, J. 306(1984)412
Vermorken, J.B., see Van der Vijgh, W.J.F. 310(1984)335
Veronese, M.E., see Davies, N.W. 310(1984)179
Verzár-Petri, G., see Lemberkovics, É. 318(1985)125
Vessman, J., see Antonsson, A.-M. 308(1984)181
Vickers, S., see Musson, D.G. 308(1984)251
Vidal, J., see Crétin, C. 315(1984)430
Vie, M., see Torel, J. 323(1985)447
Vigh, G. and Leitold, A.
Indirect UV detection of non-absorbing solutes in reversed-phase HPLC with the help of UV-active, non-ionic mobile phase constituents 312(1984)345
Vigh, G., see Bartha, A. 303(1984)29
Vijayalakshmi, M.A., see Amourache, L. 303(1984)285
Vijayalakshmi, M.A., see Rajgopal, S. 315(1984)175
Vijgh, W.J.F. van der, see Elferink, F. 320(1985)379
Vijgh, W.J.F. van der, see Gooijer, C. 312(1984)337
Vijgh, W.J.F. van der, see Van der Vijgh, W.J.F. 310(1984)335
Villar, A., Rios, J.L., Simeón, S. and Zafra-Polo, M.C.
New reagent for the detection of sesquiterpene lactones by TLC 303(1984)306
Vimalasiri, P.A.D.T., Haken, J.K. and Burford, R.P.
Chromatographic analysis of polyether based polyurethanes after alkali fusion 319(1985)121
Vimalasiri, P.A.D.T., see Burford, R.P. 321(1985)295
Vincek, W.C., see Musson, D.G. 308(1984)251
Vincent, A., see Lodge, B.A. 301(1984)477
Vincent, J.E., see Zijlstra, F.J. 311(1984)39
Vinson, J.A. and Patel, A.H.
Detection and quantitation of tetrahydrocannabinol in serum using TLC and fluorometry 307(1984)493
Visconti, M., see Crolla, T. 310(1984)139
Visser, T., Oostelbos, M.C.J.M. and Toll, P.J.M.M.
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81
Vistelle, R., see Lamiable, D. 309(1984)222
Vlasáková, V., see Tolman, V. 315(1984)421
Von Alten, R., see Heizmann, P. 310(1984)119
Vouros, P., see Karger, B.L. 323(1985)13
Vouros, P., see Mohamed, G.B. 314(1984)211
Vouros, P., see Yu, T.J. 301(1984)425
Voyksner, R.D., Bursey, J.T. and Hines, J.W.
Analysis of candidate anticancer drugs by thermospray HPLC-MS 323(1985)383
Voyksner, R.D., Bursey, J.T. and Pellizzari, E.D.
Analysis of selected pesticides by HPLC-MS 312(1984)221
Vrátný, P. and Mudřík, Z.
LC of organic acids in silage extracts using dual detection 322(1985)352
Vree, T.B., see Oosterbaan, M.J.M. 306(1984)323
Vries, J.X. de, see De Vries, J.X. 309(1984)219
Vučković, G., Janjić, T.J., Tešić, Ž.L. and Ćelap, M.B.
Effect of the geometric configuration of square-planar complexes on their R_F values obtained by TLC on silica gel 324(1985)431

W

Wad, N.
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127
Wada, H., Makino, K., Takeuchi, T., Hatano, H. and Noguchi, K.
High-performance gel filtration of water-soluble samples on polyvinyl alcohol columns 320(1985)369
Wagner, J., see Hornsperger, J.-M. 306(1984)364
Wagner, J.G., see Shukla, U.A. 308(1984)189
Wagner, K.R., see Allen, K.G. 309(1984)33
Wagner-Redeker, W., see Levsen, K. 323(1985)135
Wahlund, K.-G.
Polyion ion-pair LC. I. A new method for

the regulation of the retention of ionic compounds in LC 314(1984)127
Wahlund, K.-G., see Arvidsson, T. 317(1984)213
Wainer, I.W., Doyle, T.D., Donn, K.H. and Powell, J.R.
Direct enantiomeric determination of (-)- and (+)-propranolol in human serum by HPLC on a chiral stationary phase 306(1984)405
Wainer, I.W., see Yang, Z.-Y. 324(1985)444
Wainwright, M.S., Nieass, C.S., Haken, J.K. and Chaplin, R.P.
Use of retention plots of *n*-alkyl benzenes for determining dead times in LC and GC 321(1985)287
Wajon, J.E., Alexander, R. and Kagi, R.I.
Determination of trace levels of dimethyl polysulphides by capillary GC 319(1985)187
Walbroehl, Y. and Jorgenson, J.W.
On-column UV absorption detector for open tubular capillary zone electrophoresis 315(1984)135
Walford, J.A., see McDowall, R.D. 317(1984)475
Waliszewski, S.M. and Szymczyński, G.A.
Inexpensive, precise method for the determination of chlorinated pesticide residues in soil 321(1985)480
Walker, C.H., see Rosenfeld, J.M. 325(1985)309
Walker, R.F., Ellwood, P.A., Hardy, H.L. and Goldberg, P.A.
Separation of isocyanate prepolymer components as their urea derivatives by reversed-phase HPLC 301(1984)485
Walle, T., see Wilson, M.J. 310(1984)424
Walle, T., see Wingstrand, K.H. 305(1984)250
Wallwork, M.A., see Gilbert, J. 320(1985)361
Walmsley, L.M., Brodie, R.R. and Chasseaud, L.F.
Determination of carboxybupranolol, the major metabolite of bupranolol, in human plasma by HPLC 311(1984)227
Walsh, J.H., see Reeve, J.R. Jr. 321(1985)421
Walsh, K.A., see Sasagawa, T. 307(1984)29
Walters, R.R., see Moore, R.M. 317(1984)119
Walters, S.M., Westerby, B.C. and Gilvydis, D.M.
Determination of phenylurea pesticides by HPLC with UV and photoconductivity detectors in series 317(1984)533
Walther, H.J., see Hirter, P. 323(1985)89
Wandrey, C., see Hamacher, K. 319(1985)311
Wang, N.S., see De Vries, J.X. 309(1984)219
Wännman, T., see Buijten, J. 301(1984)265
Wännman, T., see Hoffmann, S. 302(1984)95
Wännman, T., see Markides, K. 302(1984)319
Ward, S.A., Edwards, G., Orme, M.L'E. and Breckenridge, A.M.
Determination of primaquine in biological fluids by reversed-phase HPLC 305(1984)239
Warden, B.A. and Giese, R.W.
Soluble antibody affinity chromatography technique investigated with ultratrace [^{125}I]thyroxine 314(1984)295
Wardencki, W. and Staszewski, R.
Nickel(II)-exchanged molecular sieve 13X as a column packing for the GC determination of thiophene in benzene with flame ionization detection 325(1985)430
Warnhoff, M.
Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271
Watabe, K. and Gil-Av, E.
Resolution of α-halogenocarboxylic acids by GC on (*R*)-N-lauroyl-α-(1-naphthyl)ethylamine 318(1985)235
Watanabe, M., see Kawamoto, T. 305(1984)373
Watanabe, N.
Amperometric detection of muramic acid in HPLC with a post-column reaction 316(1984)495
Watanabe, Y. and Imai, K.
Sensitive detection of amino acids in human serum and dried blood disc of 3 mm diameter for diagnosis of inborn errors of metabolism 309(1984)279
Waterbeemd, H. van de, see El Tayar, N. 320(1985)293, 320(1985)305
Weaver, K.M., Luker, R.G. and Neale, M.E.
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288
Weber, D.J., see Shah, J.A. 309(1984)95
Weber, D.V., see Roy, S.K. 303(1984)225
Weber, H., Spahn, H., Mutschler, E. and Möhrke, W.
Activated α-alkyl-α-arylacetic acid enantiomers for stereoselective TLC and HPLC determination of chiral amines 307(1984)145
Weber, H., see Spahn, H. 310(1984)167
Weber, J.D., see Yang, Z.-Y. 324(1985)444
Weber, R., see Tiebach, R. 318(1985)103
Weber, S.G. and Orr, J.D.
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433
Weber, S.G., see Podcasy, J.A. 315(1984)111
Wedlund, P.J., Sweetman, B.J., McAllister, C.B., Branch, R.A. and Wilkinson, G.R.
Direct enantiomeric resolution of mephenytoin and its N-demethylated metabolite in plasma and blood using chiral capillary GC 307(1984)121
Weems, H.B., see Yang, S.K. 316(1984)569
Weesie, H.M., see De Ligny, C.L. 301(1984)311
Weinberger, R.
Solvent and pH effects on peroxyoxalate chemiluminescence detection for LC 314(1984)155
Weinberger, R., see Koziol, T. 317(1984)355
Weiner, S., see Weinstein, S. 303(1984)244
Weinkam, R.J., see Toren, P.C. 305(1984)171

Weinstein, S. and Grinberg, N.
Enantiomeric separation of underivatized α-methyl-α-amino acids by HPLC 318(1985)117
Weinstein, S. and Weiner, S.
Enantiomeric analysis of a mixture of the common protein amino acids as their Dns derivatives. Single-analysis reversed-phase HPLC procedure using a chiral mobile phase additive 303(1984)244
Weinstein, S., see Grinberg, N. 303(1984)251
Weis, L., see Binder, H. 303(1984)375
Weiser, E.L., Salotto, A.W., Flach, S.M. and Snyder, L.R.
Basis of retention in normal-phase HPLC with cyano-propyl columns 303(1984)1
Weisner, S., see Kim, K.R. 325(1985)127
Weitzmann, C.J., see Kerlavage, A.R. 317(1984)201
Welling, G.W., see Van der Zee, R. 325(1985)187
Welling, P., Poppe, H. and Kraak, J.C.
Preparation of efficient and stable reversed-phase microbore columns for HPLC 321(1985)450
Wells, G.
Design and performance of a mas-flow-modulated detector for GC 319(1985)263
Wells, G., see Simon, R. 302(1984)221
Wells, P.G., see To, E.C.A. 301(1984)282
Wen, L. and Reeck, G.R.
Purification of high mobility group nonhistone chromosomal proteins by LC on a column containing immobilized histone H5 314(1984)436
Wendsjö, S., see Persson, B. 321(1985)375
Wendt, G., see Heizmann, P. 310(1984)119
Wenger, T.L., see Luzzi, F.A. 311(1984)291
Wentworth, W.E., see Hernandez-Gil, N. 312(1984)31
West, N., see Ray, G. 317(1984)455
Westerberg, R.B., see Buchta, R.C. 325(1985)456
Westerby, B.C., see Walters, S.M. 317(1984)533
Westholm, F.A., see Short, M.T. 323(1985)418
Wetlaufer, D.B., see Parente, E.S. 314(1984)337
Weyhenmeyer, R., see Backhaus, J. 307(1984)190
Wheals, B.B., see White, P.C. 303(1984)211
Wheeler, W.B., see Bardalaye, P.C. 314(1984)450
Whenham, R.J.
Sensitive assay for amino sugars using capillary GC with nitrogen-selective detection 303(1984)380
White, E. V, see Christensen, R.G. 323(1985)33
White, L.E., see Rege, A.B. 309(1984)397
White, P.C. and Wheals, B.B.
Use of a rotating disc multiwavelength detector operating in the visible region of the spectrum for monitoring ball pen inks separated by HPLC 303(1984)211
Whitney, J.O., see Halloran, B.P. 303(1984)229
Wiatr, R., see Breyer-Pfaff, U. 309(1984)107
Wiberg, K., see Hansén, L. 312(1984)489
Wiegand, U.--W., Chou, R.C., Lanz, E. and Jähnchen, E.
Determination of bupivacaine in human plasma by HPLC 311(1984)218
Wiehle, R.D., Hofmann, G.E., Fuchs, A. and Wittliff, J.L.
High-performance size exclusion chromatography as a rapid method for the separation of steroid hormone receptors 307(1984)39
Wiel, J.P. van der, see Stikkelman, R.M. 322(1985)220
Wielders, J.P.M. and Mink, C.J.K.
Analysis of vanillylmandelic acid, homovanillic acid and 5-hydroxyindoleacetic acid in human urine by HPLC and fluorometry 310(1984)379
Wiesler, D., see Alasandro, M. 308(1984)1
Wiesler, D., see Holland, M.L. 306(1984)23
Wiesler, D., see Raymer, J. 325(1985)13
Wiesmann, W.P., see Nelson, L.D. 324(1985)203
Wilkerson, K., see Annesley, T. 305(1984)199
Wilkes, B.C., see Cody, W.L. 314(1984)313
Wilkinson, G.R., see Wedlund, P.J. 307(1984)121
Williams, D.H., see Roets, E. 303(1984)117
Williams, E.J., see Hutte, R.S. 302(1984)173
Williams, G.H., see Emanuel, R.L. 312(1984)285
Williams, J.M., see Guieze, P. 312(1984)261
Williams, L.A., see Buchta, R.C. 325(1985)456
Williams, M.C., Kelley, J.L. and Kushwaha, R.S.
Detection of an abnormal lipoprotein in a large colony of pedigreed baboons using high-performance gel exclusion chromatography 308(1984)101
Williams, N.E., see Murray, G.R. 308(1984)143
Williams, P.S., see Ali, S.G.A.H. 302(1984)119
Williams, P.S., see Purnell, J.H. 321(1985)249, 323(1985)402, 325(1985)1
Willis, C.R., Greenblatt, D.J., Benjamin, D.M. and Abernethy, D.R.
Simultaneous determination of lidocaine and its deethylated metabolites using GLC with nitrogen-phosphorus detection 307(1984)200
Willis, C.R., see Abernethy, D.R. 307(1984)194
Wills, R.B.H., see Wimalasiri, P. 318(1985)412
Willson, N.J., see Kogan, M.J. 306(1984)441
Wilson, I.D.
TLC of ecdysteroids 318(1985)373
Wilson, I.D., see Lewis, S. 312(1984)133
Wilson, M.J. and Walle, T.
Silica gel HPLC for the simultaneous determination of propanolol and 4-hydroxypropranolol enantiomers after chiral derivatization 310(1984)424
Wilson, T.D.
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39
Wilson, T.W., McCauley, F.A. and Tuchek, J.M.
Determination of 6-keto-prostaglandin $F_{1\alpha}$ and thromboxane B_2 in urine by HPLC and radioimmunoassay 306(1984)351

Wimalasiri, P. and Wills, R.B.H.
Simultaneous analysis of thiamin and riboflavin in foods by HPLC 318(1985)412
Wingstrand, K.H. and Walle, T.
Isolation from urine of 4'-hydroxypropranolol sulphate, a major new propranolol metabolite, by ion-pair extraction 305(1984)250
Winkler, H. and Lemmer, B.
Determination of the beta-adrenoceptor blocking drug bupranolol in plasma and tissues of the rat by HPLC with UV detection 309(1984)193
Winkler, H., see Lemmer, B. 309(1984)187
Winslow, D.P., see Smith, J.A. 305(1984)295
Winter, M. and Herrmann, K.
Analysis of hydroxycinnamic acid esters and their glucosides by reversed-phase HPLC after polyamide separation 315(1984)243
Wise, S.A., see Diamondstone, B.I. 321(1985)319
Wise, S.A., see Sander, L.C. 316(1984)163
Wisneski, H.H., Yates, R.L. and Davis, H.M.
HPLC fluorometric determination of cinnamaldehyde in perfume, cologne and toilet water 317(1984)421
Witkiewicz, Z., Grajek, H. and Szulc, J.
Liquid crystalline stationary phases with a cyclohexyl ring in the molecule 312(1984)141
Witkiewicz, Z., Szulc, J. and Dąbrowski, R.
Disc-like liquid crystalline stationary phases from the triphenylene derivatives group 315(1984)145
Witkiewicz, Z., see Marciniak, W. 324(1985)299, 324(1985)309
Witter, F.R., see Bauza, M.T. 310(1984)61
Wittliff, J.L., see Wiehle, R.D. 307(1984)39
Wittmann-Liebold, B., see Kamp, R.M. 317(1984)181
Wold, J.K., see Haavik, S. 321(1985)199
Womersley, C., Drinkwater, L. and Crowe, J.H.
Separation of tricarboxylic acid cycle acids and other related organic acids in insect haemolymph by HPLC 318(1985)112
Wong, L.T. and Hsia, J.C.
HPLC of proteins: purification of α-fetoprotein from fetal calf serum 310(1984)19
Woodrow, J.E., see Kim, Y.-H. 314(1984)37
Woods, K.L., see Siegel, S.A. 319(1985)351
Woollard, D.C.
New ion-pair reagent for the HPLC separation of B-group vitamins in pharmaceuticals 301(1984)470
Woollard, G.A.
Effect of precipitating agents on the analysis of metronidazole by HPLC 303(1984)222
Woollard, P.M. and Mallet, A.I.
Lipoxygenase products. A novel GC-MS assay for monohydroxy fatty acids 306(1984)1
Wormser, U., see Laufer, R. 301(1984)415
Wozniak, J.M., see Lakings, D.B. 308(1984)261
Wright, B.W., Udseth, H.R., Smith, R.D. and Hazlett, R.N.
Supercritical fluid chromatography and supercritical fluid chromatography-MS of marine diesel fuel 314(1984)253
Wright, G.J., see Krebs, H.A. 310(1984)412
Wroński, M.
Electrophoresis of thiols in cellulose gels. III. Group analysis of urinary thiols 307(1984)416
Wu, C.-G. and Deming, S.N.
Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79
Wu, H.-L., see Funazo, K. 319(1985)143
Wurst, M., see Kysilka, R. 320(1985)414
Wyles, H.F., see Buchta, R.C. 325(1985)456
Wyse, D.G., see Tam, Y.K. 310(1984)438

X

Xie, K.-H., see Driscoll, J.N. 302(1984)43
Xu, G.-Y., see Zhu, A. 314(1984)421
Xu, X.B. and Jin, Z.L.
HPLC studies of environmental carcinogens in China 317(1984)545

Y

Yacobi, A., see Stampfli, H.F. 309(1984)203
Yajima, T., see Suzuki, K.T. 303(1984)131
Yamada, H. and Hattori, T.
Determination of molybdenum and tungsten with tiron by reversed-phase LC 320(1985)403
Yamada, J., Sugimoto, Y. and Horisaka, K.
Simultaneous determination of tryptamine and its metabolites in mouse brain by HPLC with fluorometric detection 311(1984)385
Yamada, Y., see Suemitsu, R. 318(1985)139
Yamagishi, A.
Clay CC for optical resolution. Resolution of aromatic compounds on a Λ-Ru(1,10-phenanthroline)$_3^{2+}$-montmorillonite column 319(1985)299
Yamaguchi, K., see Parker, C.E. 319(1985)273
Yamaguchi, M., see Ishida, J. 305(1984)381
Yamaguchi, M., see Nakano, Y. 311(1984)390
Yamaguchi, Y., see Seki, T. 305(1984)188
Yamaguchi, Y., see Yoshida, A. 307(1984)399

Yamamoto, I., see Imaizumi, N. 308(1984)295
Yamamoto, M., see Miwa, H. 321(1985)165
Yamamoto, S., Iwado, A., Hashimoto, Y., Aoyama, Y. and Makita, M.
GC-MS of polyamines as their N-ethyloxycarbonyl derivatives and identification of *sym*-homospermidine and *sym*-norspermine in mosses and ferns 303(1984)99
Yamamoto, S., see Kataoka, H. 306(1984)61
Yamamura, K. and Yotsuyanagi, T.
HPLC assay for prostaglandin E_1 in various ointment vehicles. Separation and stability testing 303(1984)165
Yamasaki, Y., see Kato, Y. 320(1985)440
Yamauchi, S., see Kawamoto, T. 305(1984)373
Yamazaki, S., see Yoneda, H. 319(1985)35
Yamazaki, Y., see Kato, Y. 318(1985)358
Yan, C., see Sloan, D.L. 316(1984)43
Yanase, T., see Imamura, T. 305(1984)456
Yang, D., see Hjertén, S. 316(1984)301
Yang, S.K., Weems, H.B., Mushtaq, M. and Fu, P.P.
Direct resolution of mono- and diol enantiomers of unsubstituted and methyl-substituted benz[a]anthracene and benzo[a]pyrene by HPLC with a chiral stationary phase 316(1984)569
Yang, T.-H., Zabriskie, T.M. and Poulter, C.D.
Investigation of isoprenoid benzoates and naphthoates by reversed-phase LC. Isocratic elution characteristics of benzoates and naphthoates of C_5-C_{20} terpenoid alcohols 312(1984)121
Yang, Z.-Y., Barkan, S., Brunner, C., Weber, J.D., Doyle, T.D. and Wainer, I.W.
Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444
Yasmin, T. and Qadri, F.
Purification of placental alkaline phosphatase by sulphate-mediated chromatography on Cibacron Blue 3G-A 315(1984)425
Yasuda, H., see Miyazaki, K. 310(1984)219
Yates, R.L., see Wisneski, H.H. 317(1984)421
Yau, M.L., see Tam, Y.K. 310(1984)438
Yau, W.W., see Dilks, C.H. Jr. 315(1984)45
Yeager, R.L., Oldfield, E.H. and Chatterji, D.C.
Quantitation of 1,3-bis(2-chloroethyl)-1-nitrosourea in plasma using HPLC 305(1984)496
Yeh, S.C., see Krol, G.J. 305(1984)105
Yert, L.W., see Cline, R.E. 307(1984)420
Yeung, P.K.F., Hubbard, J.W., Baker, B.W., Looker, M.R. and Midha, K.K.
Isotopic fractionation of N-([^{3}H]methyl)-chlorpromazine and N-([^{3}H]methyl)-7-hydroxychlorpromazine by reversed-phase HPLC 303(1984)412
Yip, M.S., Coates, P.E. and Thiessen, J.J.
HPLC analysis of amiloride in plasma and urine 307(1984)343
Yokota, E., see Imamura, T. 305(1984)456
Yoneda, H.
Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59
Yoneda, H., Nakatani, K. and Yamazaki, S.
Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 319(1985)35
Yoo, B.-T., see Nohta, H. 308(1984)93
Yoshida, A., Yamaguchi, Y. and Yoshioka, M.
Determination of creatinine in human, dog and rat urine by HPLC on a column of hydroxymethylated porous polystyrene 307(1984)399
Yoshida, H., Hida, M. and Taga, M.
Capillary-tube isotachophoretic separation of molybdate VI, Tungstate VI and Vanadate V ions using ion pairing equilibria 325(1985)179
Yoshida, H., see Hirama, Y. 322(1985)139
Yoshida, S., see Iwamoto, M. 310(1984)151
Yoshida, T., see Tanaka, E. 308(1984)393
Yoshioka, M., Nishidate, K., Iizuka, H., Nakamura, A., El-Merzabani, M.M., Tamura, Z. and Miyazaki, T.
Sensitive fluorimetry of adenine-containing compounds with HPLC 309(1984)63
Yoshioka, M., see Yoshida, A. 307(1984)399
Yotsuyanagi, T., see Yamamura, K. 303(1984)165
You, F., see Bidleman, T.F. 301(1984)448
Young, J.D., see Evans, L.J. Jr. 305(1984)163
Young, M., see Tsao, C.S. 308(1984)306
Young, P.M., Boehm, T.M. and Brown, J.E.
Resolution and quantitation of apolipoproteins A-I and A-II from human high-density lipoprotein by size exclusion HPLC 311(1984)79
Yoza, N., Hirano, H., Baba, Y. and Ohashi, S.
Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385
Yoza, N., see Baba, Y. 318(1985)319
Yu, H., see Mourey, T.H. 303(1984)361
Yu, T.J., Schwartz, H.A., Cohen, S.A., Vouros, P. and Karger, B.L.
Sequence analysis of derivatized peptides by HPLC-MS 301(1984)425
Yui, S., see Tokuda, M. 323(1985)434

Z

Zabriskie, T.M., see Yang, T.-H. 312(1984)121

Zadro, S., Haken, J.K. and Pinczewski, W.V.
Analysis of Australian crude oils by high-resolution capillary GC-MS 323(1985)305
Zafra-Polo, M.C., see Villar, A. 303(1984)306
Zaikin, V.G. and Mikaya, A.I.
Reaction GC-MS. VI. Use of deuterium as the carrier and reagent gas in reaction GC-MS 301(1984)77
Zambonin, P.G., see Palmisano, F. 306(1984)205
Zamureenko, V.A., Kluyev, N.A., Dmitriev, L.B. and Grandberg, I.I.
GLC-MS in the analysis of essential oils 303(1984)109
Zee, R. van der, see Van der Zee, R. 325(1985)187
Zelenská, V., see Zelenský, I. 325(1985)161
Zelenský, I., Šimuničová, E., Zelenská, V., Kaniansky, D., Havaši,P. and Chaláni, P.
Factors influencing the reproducibility of isotachophoretic determinations of ions present in wet-process phosphoric acid 325(1985)161
Zeleznikar, R.J. Jr., see Singh, A.K. 324(1985)163
Žemlová, Z., see Pospíchal, J. 320(1985)139
Zhang, T.-Y.
Horizontal flow-through coil planet centrifuge: some practical applications of countercurrent chromatography 315(1984)287
Zhirkov, Y.A., see Piotrovskii, V.K. 309(1984)421
Zhu, A. and Xu, G.-Y.
Derivatization at the liquid-solid interface. Application to the trace analysis of organic acids in aqueous samples 314(1984)421
Ziegler, K., see Blasius, E. 313(1984)161
Zieliński, P., see Paryjczak, T. 301(1984)270
Ziemniak, J.A., see Boppana, V.K. 317(1984)463
Zijlstra, F.J. and Vincent, J.E.
Determination of leukotrienes and prostaglandins in [^{14}C]arachidonic acid labelled human lung tissue by HPLC and radioimmunoassay 311(1984)39
Zijp, J.J., see Meering, P.G. 310(1984)159
Zinapold, K., see Heizmann, P. 310(1984)119, 310(1984)129
Živný, K., see Tolman, V. 315(1984)421
Zlatkis, A., see Kim, K.R. 325(1985)127
Zonta, F. and Stancher, B.
HPLC of retinals, retinols (vitamin A_1) and their dehydro homologues (vitamin A_2): improvements in resolution and spectroscopic characterization of the stereoisomers 301(1984)65
Zonta, F., see Stancher, B. 312(1984)423
Zoontjes, P.W., see Jansen, E.H.J.M. 319(1985)436
Zu-Ben, C., see Kirkman, C.M. 317(1984)569
Žunić, G., Stanimirović, S. and Savić, J.
Rapid ion-exchange method for the determination of 3-methylhistidine in rat urine and skeletal muscle 311(1984)69
Zuska, J., see Tamchyna, J. 320(1985)241
Zuska, J., see Vacík, J. 320(1985)233
Zwanenburg, B., see Van Hogezand, R.A. 305(1984)470

SUBJECT INDEX

VOLUMES 301—325

A

R-α-Methylbenzyl isothiocyanate
R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Abscisic acid
Less hazardous derivatization procedure for GC of plant hormone abscisic acid 325(1985)425

Absorbance-fluorescence detection
Detection of GC eluates by simultaneous absorbance and fluorescence measurements 303(1984)341

π-Acceptors
Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

Acebutolol
Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Separation of acebutolol and diacetolol diastereomers by reversed-phase HPLC 318(1985)393

Acephate
Improved analysis of acephate and methamidophos in biological samples by selective ion monitoring GC-MS 301(1984)465

Acetamidoglutethimide
GLC assay of aminoglutethimide and a HPLC assay for its acetyl metabolite in biological fluids 307(1984)129

Acetate complexes
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11

Acetic acid
Steam GSC: a flexible separation technique 301(1984)11

Acetone
Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

Acetonitrile
Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

N-Acetylaminoglutethimide
Determination of aminoglutethimide and N-acetylaminoglutethimide in human plasma by HPLC 310(1984)431

Acetylcholine
Volatile products from acetylcholine as markers in the rapid urine test using head-space GLC 307(1984)11

Enzymes adsorbed on an ion exchanger as a post-column reactor: application to acetylcholine measurement 317(1984)343

Chromatographic analysis of choline and acetylcholine by UV visualization 319(1985)454

Determination of choline and acetylcholine levels in rat brain regions by LC with electrochemical detection 322(1985)191

Acetylcholinesterase
Liquid-liquid extraction of membranes from calf brain using conventional and centrifugal counter-current distribution techniques 311(1984)277

Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

N-Acetylcysteine
Determination of non-protein-bound N-acetylcysteine in plasma by HPLC 311(1984)170

β-N-Acetylhexosaminidase
Preparative isolation of the polypeptide chains of human liver β-N-acetylhexosaminidase A by

fast protein LC-ion-exchange chromatography 323(1985)439

N-Acetylhistamine
Reversed-phase ion-pair HPLC procedure for determination of histamine and its metabolites in rat urine 312(1984)443

Acetylisoniazid
Determination of isoniazid and acetylisoniazid in plasma or serum by HPLC (in French) 307(1984)137

N-Acetyl-L-cysteine
Determination of N-acetyl-L-cysteine in biological fluids 309(1984)261

N-Acetylneuraminic acid
Automatic amino acid and sugar analysis of glycoproteins 317(1984)35

Acetylsalicylic acid
Rapid and sensitive determination of acetylsalicylic acid and its metabolites using reversed-phase HPLC 310(1984)343

Improved method for the determination of aspirin and its metabolites in biological fluids by HPLC: applications to human and animal studies 311(1984)301

Simultaneous HPLC assay of acetylsalicylic acid and salicylic acid in film-coated aspirin tablets 317(1984)507

N-Acetyl serotonin
Use of impregnated reagents in the pentafluorobenzylation of indoleamine metabolites 325(1985)309

Acetylsulphadoxine
Quantification of antimalarial drugs. I. Simultaneous measurement of sulphadoxine, N_4-acetylsulphadoxine and pyrimethamine in human plasma 305(1984)502

N^4-Acetylsulphadoxine
HPLC assay of pyrimethamine, sulfadoxine and its N^4-acetyl metabolite in serum and urine after ingestion of Suldox 308(1984)217

α_1-Acid glycoprotein
Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Acid strength
Effects of various strengths of acids on the electrophoretic behavior of weak bases in a non-aqueous solvent 314(1984)65

Aconitine
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Acridine
Effect of trialkylamines on the response of the electron-capture detector to acridine 312(1984)49

Acrylate monomers
Chromatography of monomers. II. Glass capillary GC of C_1-C_{18} alkyl esters of acrylic and methacrylic acids 325(1985)367

Actinobacillus actinomycetemcomitans
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Determination of bound cellular fatty acids in *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* by GC and GC-MS 308(1984)282

Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Carbohydrates of whole defatted cells as a basis for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)31

Whole-cell methanolysis as a rapid method for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)347

Activated carbon
Ion-exchange properties of activated carbon filled with hydrous Zr(IV) and Zr(IV)-P(V) oxides 303(1984)13

Activity coefficients
Mixed solvents in GLC. Activity coefficients for chloroform, dichloromethane and ethyl acetate in squalane-dinonyl phthalate mixtures at 303°K 324(1985)231

Adenine
Separation of adenosine and its derivatives by TLC on silica gel 325(1985)317

Adenine nucleotides
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 305(1984)183

Sensitive fluorimetry of adenine-containing compounds with HPLC 309(1984)63

Rapid analysis of adenosine, AMP, ADP, and ATP by anion-exchange CC 311(1984)396

HPLC determination of adenine nucleotides in biological materials. Improvements and adaptations to routine analysis 325(1985)314

Separation of adenosine and its derivatives by TLC on silica gel 325(1985)317

Adenosine
Measurement of adenosine, inosine, and hypoxanthine in human plasma 307(1984)404

Rapid analysis of adenosine, AMP, ADP, and ATP by anion-exchange CC 311(1984)396

Micro column LC-MS using a capillary interface 323(1985)89

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Separation of adenosine and its derivatives by TLC on silica gel 325(1985)317

Adenosine deaminase
Silica-bound polyethyleneglycol as stationary phase for separation of proteins by HPLC 319(1985)396

Adrenaline
Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

HPLC determination of plasma catecholamines during α-methyldopa therapy 308(1984)301

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Dihydrocaffeic acid: a common contaminant in the LC-electrochemical measurement of plasma catecholamines in man 311(1984)148

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

HPLC analysis of epinephrine enantiomers using a UV detector in series with an optical activity detector 319(1985)419

HPLC determination of *d-/l-* epinephrine enantiomer ratio in lidocaine-epinephrine local anesthetics 325(1985)249

Adriamycin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

Continuous extraction of urinary anthracycline antitumor antibiotics with the horizontal flow-through coil planet centrifuge 307(1984)323

Determination of adriamycin, adriamycinol and their 7-deoxyaglycones in human serum by HPLC 311(1984)125

Adriamycinol
Determination of adriamycin, adriamycinol and their 7-deoxyaglycones in human serum by HPLC 311(1984)125

Adsorbents
Steam GSC: a flexible separation technique 301(1984)11

Sample enrichment for GC-MS analysis of polynuclear aromatic hydrocarbons in water and in organic mixtures 302(1984)167

Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333

Adsorption
Effect of activated sorption processes on the results of thermoprogrammed reduction investigations on Fe-Al_2O_3 catalysts 301(1984)270

Characterization of adsorption at kaolinite surfaces by heterogeneous GSC 314(1984)111

Adsorption isotherms
GC determination of binary adsorption isotherms 302(1984)195

Studies in reversed-phase ion-pair chromatography. IV. The role of the chain length of the pairing ion 303(1984)29

Measurement of adsorption isotherms by LC 316(1984)53

Gradient elution micellar LC 316(1984)183

Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267

GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19

Affinity adsorbents
Immobilization of ligands for affinity chromatography. Coupling on a spacer arm gel with N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline as condensation agent: study of coupling conditions by means of a radioimmunologic method 314(1984)445

Introduction of ω-carboxyl spacers onto cross-linked agarose gel beads by O-alkylation for the preparation of affinity adsorbents 315(1984)161

Sepharose derivatives containing citric acid as affinity ligand. Purification of fumarase 319(1985)173

Cibacron Blue 3G-A-substituted cross-linked agar beads: an inexpensive, fast-flow affinity medium for large scale applications 319(1985)218

Affinity partitioning
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Aflatoxins
HPLC studies of environmental carcinogens in China 317(1984)545

Determination of aflatoxins B_1, B_2, G_1, G_2 and M_1 by over-pressurized TLC (in German) 319(1985)105

Design and construction of an interface for direct liquid introduction coupling of a microbore high-performance liquid chromatograph to a quadrupole mass spectrometer 323(1985)121

Agarose
Synthesis of new hydrophobic adsorbents based on homologous series of uncharged alkyl sulphide agarose derivatives 321(1985)305

Quantitative determination of the ligand content and the negative and positive charges on butylamine-substituted Sepharose 4B 321(1985)335

Air particulate matter
Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53

Air samplers
Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448

Albumin
Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

Direct LC separation of enantiomers on immobilized protein stationary phases. IV. Molecular interaction forces and retention behaviour in chromatography on bovine serum albumin as a stationary phase 316(1984)617

Protein separations on reversed-phase HPLC minicolumns 317(1984)119

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Alcohol dehydrogenase
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Rapid chromatographic purification of yeast alcohol dehydrogenase 325(1985)359

Alcohols
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47

Evaluation of hydrophobicity of gels by use of some 1-alkanols as pilot solutes 301(1984)57

Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173

Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167

HPLC of low-molecular-weight aliphatic alcohols using indirect photometric detection 314(1984)488

Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. I. Sorption isotherms of *n*-alcohols on octyl-agarose 322(1985)399

HPLC elution behaviour of oligosaccharides, monosaccharides and sugar degradation products on series-connected ion-exchange resin columns using water as the mobile phase 322(1985)411

Separation and double-bond determination on nanogram quantities of aliphatic monounsaturated alcohols, aldehydes and carboxylic acid methyl esters 322(1985)484

GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

Alcoholysis
Capillary GC investigation of plasma lipid alcoholysis during alcohol extraction 307(1984)172

Aldehydes
Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23

Determination of aldehydes in air as dimethone derivatives by GC with electron-capture detection 315(1984)412

HPLC of carbonyl compounds as 2,4-dinitrophenylhydrazones with electrochemical detection 318(1985)427

Determination of carbonyl compounds in an Australian (Rundle) shale oil 319(1985)230

HPLC elution behaviour of oligosaccharides, monosaccharides and sugar degradation products on series-connected ion-exchange resin columns using water as the mobile phase 322(1985)411

Separation and double-bond determination on nanogram quantities of aliphatic monounsaturated alcohols, aldehydes and carboxylic acid methyl esters 322(1985)484

Alditol acetates
Sensitive assay for amino sugars using capillary GC with nitrogen-selective detection 303(1984)380

Capillary GC of partially methylated alditol acetates on a high-polarity bonded-phase vitreous-silica column 315(1984)373

Aldonic acids
Analysis of uronic and aldonic acids, their lactones, and related compounds by HPLC on cation-exchange resins 319(1985)159

Aldosterone
Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Rapid chromatographic purification of urinary steroid glucuronates for determination of aldosterone secretion rate 317(1984)367

Alginates
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Alimemazine
Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Alkali fusion
Chromatographic analysis of polyester-based polyurethane elastomers after alkali fusion 321(1985)295

Alkaline phosphatase
Purification of placental alkaline phosphatase by sulphate-mediated chromatography on Cibacron Blue 3G-A 315(1984)425

Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

Alkaloids
Determination of glaucine in plasma and urine by HPLC 308(1984)273

Use of electrochemical detection in the HPLC determination of hydroxylated ellipticine derivatives 309(1984)170

New method for the determination of yohimbine in biological fluids by HPLC with amperometric detection 311(1984)449

Qualitative and quantitative determinations of hallucinogenic components of psilocybe mushrooms by reversed-phase HPLC 312(1984)467

Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59

Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Inter-laboratory study on the separation of cinchona alkaloids by HPLC 318(1985)362

Qualitative and quantitative analysis of hyoscyamine-atropine and scopolamine in mother tinctures of Solanaceae by HPLC (in French) 319(1985)412

Partition chromatography of ergot alkaloids on formamide-saturated silica gel columns (in German) 319(1985)440

Determination of quinine in beverages, pharmaceutical preparations and urine by isotachophoresis 320(1985)245

Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Determination of the alkaloid perloline in grasses by HPLC 322(1985)261

Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

Alkanes
Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53

Chromatography with sub- and supercritical eluents: influence of temperature, pressure and flow-rate on the behaviour of lower alkanes 321(1985)273

Reversed-phase separation of alkanes in high-boiling fractions of crude oil 322(1985)347

GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19

Alkanolamines
LC separation of enantiomeric alkanolamines via diastereomeric tartaric acid monoesters 316(1984)605

Alkanols
Steam GSC: a flexible separation technique 301(1984)11

GC of homologous esters. XXVII. Retention increments of C_1-C_{18} primary alkanols and their 2-chloropropanoyl and 3-chloropropanoyl derivatives on SE-30 and OV-351 capillary columns 319(1985)131

Alkenes
Reaction GC-MS. VI. Use of deuterium as the carrier and reagent gas in reaction GC-MS 301(1984)77

Study of GC behaviour of alkenes based on molecular orbital calculations 302(1984)107

Aryl substituted diastereomeric alkenes: GC behavior on a non-polar *versus* a liquid crystal phase 321(1985)127

Alkenesulphonates
Analysis of α-olefinsulphonates by HPLC 325(1985)43

1-O-Alkyl-2-acetyl-*sn*-glycero-3-phosphorylcholines
Rapid isocratic HPLC purification of platelet activating factor (PAF) and *lyso*-PAF from human skin 309(1984)160

Alkylagaroses
Hydrophobic chromatography of cells: adsorption and resolution on homologous series of alkylagaroses 317(1984)103

Alkylalkoxysilanes
GC investigation of organometallic compounds and their carbon analogues. VI. Determination, calculation and correlation of Kováts retention indices for alkylalkoxysilanes 318(1985)41

Alkylamines
Trace determination of aliphatic amines using HPLC with chemiluminescence excitation and photon counting 312(1984)203

Alkylanilines
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

α-Alkyl-α-arylacetic acids
Activated α-alkyl-α-arylacetic acid enantiomers for stereoselective TLC and HPLC determination of chiral amines 307(1984)145

α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167

N-Alkylarylamines
Application of reversed-phase HPLC to in vitro drug metabolism studies with N-alkylarylamines 309(1984)339

Alkylbenzenes
Dependence of the retention of phenols upon Van der Waals volume, π-energy and hydrogen-bonding effects 302(1984)89

Products obtained during combustion of polymers under simulated incinerator conditions. II. Polystyrene 315(1984)201

Products obtained during combustion of polymers under simulated incinerator conditions. III. Polyvinyl chloride 315(1984)211

Computer-assisted retention prediction system for reversed-phase micro HPLC 316(1984)1

Use of retention plots of *n*-alkyl benzenes for determining dead times in LC and GC 321(1985)287

Heats of formation-GC retention relationships for aliphatic hydrocarbons and alkylbenzenes 322(1985)35

High-performance GC on liquid crystal capillaries. VIII. Separation of alkylbenzenes up to C_9 (in German) 323(1985)414

Alkyl benzoates
GLC analyses. XXXV. Capillary column studies of C_1-C_{12} *n*-alkyl esters of benzoic and monochlorobenzoic acids 322(1985)83

Alkylbenzocarbazoles
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

Alkylbenzoquinolines
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

Alkylbenzyldimethylammonium chlorides
Retention behavior of long chain quaternary ammonium homologues and related nitroso-alkylmethylamines 324(1985)209

Alkyl bromides
Evaluation of a calibration marker scheme for open tubular column GC with on-column injection and electron-capture detection 302(1984)289

GC separation and identification of linear and branched-chain alkyl bromides 324(1985)363

Alkylcarbazoles
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

***n*-Alkyldimethylchlorosilanes**
Role of solvent and base in the silanization reaction of silicas for reversed-phase HPLC 316(1984)193

Effect of alkyl chain length on surface silanization of silica 318(1985)173

***n*-Alkyldimethyltrifluoroacetoxysilanes**
Role of solvent and base in the silanization reaction of silicas for reversed-phase HPLC 316(1984)193

Alkyl monochlorobenzoates
GLC analyses. XXXV. Capillary column studies of C_1-C_{12} *n*-alkyl esters of benzoic and monochlorobenzoic acids 322(1985)83

Alkylphenol ethoxylates
On-line LC-MS analysis of non-ionic surfactants 323(1985)135

Alkyl porphyrins
Computerised GC-MS analysis of complex mixtures of alkyl porphyrins 301(1984)107

Alkyl propanoates
GC of homologous esters. XXIX. Propanoyl and monochlorpropanoyl esters of lower saturated brached-chain and unsaturated alcohols 324(1985)343

Alkylpyridines
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

Alkylquinolines
Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

Alkyl radicals
GC identification of alkyl radicals formed in plasma radiofrequency discharges by using iodine as a scavenger 303(1984)61

Alkyl sulphide agarose
Synthesis of new hydrophobic adsorbents based on homologous series of uncharged alkyl sulphide agarose derivatives 321(1985)305

Allergens
Isotachophoresis of allergenic extracts 320(1985)205

Purification of a basic glycoprotein allergen from pollen of timothy by HPLC 321(1985)199

Allopurinol
HPLC with polarographic and voltammetric anodic detection: simultaneous determination of allopurinol, oxipurinol and uric acid in body fluids 306(1984)205

Simultaneous determination of allopurinol, oxipurinol, hypoxanthine and xanthine in biological fluids by HPLC 307(1984)469

Alprenolol
Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Altersolanol
Analysis of naturally occurring reduced anthraquinones obtained from *Alternaria porri* (Ellis) Ciferri by HPLC 318(1985)139

Aluminium chloride
New reagent for the detection of sesquiterpene lactones by TLC 303(1984)306

Aluminium cold rolling oil
Analysis of mechanical lubricant contaminants in aluminium cold rolling oil by reversed-phase LC 319(1985)222

Aluminium trifluoroacetylacetonates
Developments in the HPLC of metallo-organic compounds 317(1984)569

Amidopyrine
Development of a standardized analysis strategy for basic drugs, using ion-pair extraction and HPLC. VI. Drug level determination in saliva 309(1984)329

Amikacin
Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

LC determination of amikacin in serum with spectrophotometric detection 307(1984)224

Amiloride
HPLC analysis of amiloride in plasma and urine 307(1984)343

Amine modifiers
Effects of amine modifiers on retention and peak shape in reversed-phase HPLC 320(1985)313

Amineptine
HPLC method for the determination of amineptine and its main metabolite in human plasma 306(1984)279

Amines
Trace analysis of organic compounds by HPLC with photoionization detection 302(1984)43

Trace analysis of amines and isocyanates using glass capillary GC and selective detection. III. Determination of aliphatic and alicyclic amines as perfluoro fatty acid amides using electron-capture and nitrogen-selective detection 303(1984)89

Activated α-alkyl-α-arylacetic acid enantiomers for stereoselective TLC and HPLC determination of chiral amines 307(1984)145

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Measurement of tyramine in human plasma, utilising ion-pair extraction and HPLC with amperometric detection 310(1984)11

α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167

GC analysis of endogenous catecholamines, phenolic amines and derived isoquinolines using short glass capillary columns and electron-capture detection 311(1984)17

Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141

Trace determination of aliphatic amines using HPLC with chemiluminescence excitation and photon counting 312(1984)203

Recovery of plant amines with carboxylic ion-exchange substrates and their subsequent analysis by TLC 312(1984)357

Determination of aliphatic amines in air by reversed-phase HPLC using 1-naphthyl isocyanate derivatives 312(1984)482

Reversed-phase retention behaviour of Dns-amides as a function of eluent composition and molecular structure in ethanol-water and methanol-water 314(1984)141

R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Retention indices of dimethylformamidines, dimethylacetamidines and tetramethylguanidines on a non-polar column 315(1984)95

Direct analysis of free amines in salt solutions at sub-ppm levels by GLC 315(1984)299

Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

HPLC with amperometric determination of plasma tyramine 317(1984)319

Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Detection of urinary amine metabolites in toluene diisocyanate exposed rats 323(1985)429

Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

New high-performance LC system for the separation of biogenic amines as their Dns derivatives 325(1985)462

Amino acid oxidase
Use of immobilized amino acid oxidase as post-column reactor in the HPLC of amino acids 303(1984)396

Amino acids
Masking effects in "charge-transfer" TLC 303(1984)190

Quantitative GLC of taurine 306(1984)61

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

GC analysis of amino acids in physiological fluids: a critique (Review) 310(1984)223

Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

Reversed-phase high-performance ion-pair chromatography of cobalt(III) coordination complexes (Review) 313(1984)93

Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

Automatic amino acid and sugar analysis of glycoproteins 317(1984)35

Ligand-exchange chromatography on copper(II)-modified silica gel. Improvements and use for screening of protein hydrolyzate and quantitation of dipeptides and amino acid fractions 317(1984)41

Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

HPLC-MS of derivatized and underivatized amino acids 323(1985)67

Development of immobilized metal affinity chromatography. II. Interaction of amino acids with immobilized nickel iminodiacetate 323(1985)255

Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

In-situ preparation of a chemically bonded chiral stationary phase for the separation of aromatic α-amino acid enantiomers 325(1985)103

Amino acids, Dns derivatives
HPLC of amino acids in urine and cerebrospinal fluid 302(1984)21

Sub-picomol chemiluminescence detection of Dns-amino acids separated by HPLC with gradient elution 303(1984)173

Enantiomeric analysis of a mixture of the common protein amino acids as their Dns derivatives. Single-analysis reversed-phase HPLC procedure using a chiral mobile phase additive 303(1984)244

Enantiomeric separation of Dns-amino acids by reversed-phase TLC 303(1984)251

Microbore HPLC and chemiluminescence detection of Dns-amino acids 316(1984)501

Amino acids, heptafluorobutyryl isobutyl esters
Effect of the presence of glucose on the determination of amino acids by GC 309(1984)375

Amino acids, heptafluorobutyryl methyl esters
Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251

Amino acids, NBD derivatives
Sensitive detection of amino acids in human serum and dried blood disc of 3 mm diameter for diagnosis of inborn errors of metabolism 309(1984)279

TLC of hydroxylysine for collagen analysis 311(1984)400

Rapid and sensitive method for the determination of proline by reversed-phase HPLC with automated pre-column fluorescence derivatization 318(1985)132

Amino acids, OPA derivatives
HPLC with automated pre-column derivatization for amino acids 303(1984)238

Rapid and sensitive amino acid analysis of human collagens using HPLC 311(1984)375

Resolution of amino acid enantiomers by HPLC using automated pre-column derivatisation with a chiral reagent 315(1984)279

Application of reversed-phase HPLC using pre-column derivatization with *o*-phthaldialdehyde for the quantitative analysis of amino acids in adult and fetal sheep plasma, animal feeds and tissues 315(1984)321

Complete automatization of peptide maps by reversed-phase LC using *o*-phthalaldehyde pre-column derivatization 323(1985)373

Amino acids, oxazolidinones
Possibilities and limitations in the analysis of amino acid oxazolidinones in the femtomole range by GC with electron-capture detection 305(1984)442

Amino acids, pentane-2,4-dione-formaldehyde condensation
Post-column reaction of amino acids with the pentane-2,4-dione-formaldehyde system for their automated analysis 318(1985)367

Amino acids, PTH derivatives
Identification of phenylthiohydantoin derivatives of substituted amino acids encountered during amino acid sequence analysis by HPLC 314(1984)467

Examination of theoretical principles of gradient elution as applied to reversed-phase HPLC separation of phenylthiohydantoin amino acids 316(1984)359

Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23

HPLC-MS of derivatized and underivatized amino acids 323(1985)67

Amino acids, thiourea derivatives
Optical resolution of amino acid enantiomers by HPLC 316(1984)547

Amino acids, trifluoroacetyl *n*-propyl esters
GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

Amino acids, (N,O)-trifluoroacetyl *n*-propyl esters
Influence of trifluoroacetic anhydride concentration during derivatization and chromatography of N(O,S)-trifluoroacetyl amino acid *n*-propyl esters 318(1985)65

α-Aminoacyl hydroxamate adsorbents
α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81

Amino alcohols
Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Amino aldehydes
Sensitive assay for diamine oxidase activity using HPLC 309(1984)151

1-Aminoalkanephosphonic acids
Chromatographic assignment of the configuration of 1-aminoalkanephosphonic acids 324(1985)455

Amino amides
Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

γ-Aminobutyric acid
Determination of γ-aminobutyric acid in physiological samples by a simple, rapid HPLC method 309(1984)139

Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251

ε-Aminocaproic acid
Microdetermination of two antifibrinolytic agents (ε-aminocaproic acid and tranexamic acid) by LC and fluorimetric detection (in French) 309(1984)183

Amino esters
Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Aminoglutethimide
GLC assay of aminoglutethimide and a HPLC assay for its acetyl metabolite in biological fluids 307(1984)129

HPLC assay for aminoglutethimide and its acetylated metabolite in urine 309(1984)431

Determination of aminoglutethimide and N-acetylaminoglutethimide in human plasma by HPLC 310(1984)431

Aminoglycosides
Use of perfluorinated carboxylic acids in the separation of aminoglycoside antibiotics by ion-pair reversed-phase HPLC 303(1984)277

Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373

LC determination of amikacin in serum with spectrophotometric detection 307(1984)224

Pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene 322(1985)321

LC separation of aminoglycosides with pulsed amperometric detection 324(1985)407

4-Amino-hex-5-enoic acid
Assay of γ-vinyl-γ-aminobutyric acid (4-amino-hex-5-enoic acid) in plasma and urine by automatic amino acid analysis. Application to human pharmacokinetics 306(1984)383

***p*-Aminohippuric acid**
Simple and micro HPLC method for simultaneous determination of *p*-aminohippuric acid and iothalamate in biological fluids 306(1984)89

L-α-Amino-ε-ketopimelate
Use of isotachophoresis to monitor the synthesis and purification of L-α-amino-ε-ketopimelate 323(1985)451

9-Aminophenanthrene
Fluorometric HPLC of 9-aminophenanthrene-derivatized free fatty acids 305(1984)261

Amino phosphonates
Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Aminoquinoline
HPLC method for the analysis of a candidate 8-aminoquinoline antileishmanial drug using oxidative electrochemical detection 311(1984)117

Amino sugars
Sensitive assay for amino sugars using capillary GC with nitrogen-selective detection 303(1984)380

Automatic amino acid and sugar analysis of glycoproteins 317(1984)35

Analysis of uronic and aldonic acids, their lactones, and related compounds by HPLC on cation-exchange resins 319(1985)159

3-Amino-1,2,4-triazole
Determination of 3-amino-1,2,4-triazole (amitrole) in urine by ion-pair HPLC 303(1984)267

Amitraz
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Amitriptyline
GC-MS determination of amitriptyline and its major metabolites in human serum 305(1984)73

Amitriptyline and its basic metabolites determined in plasma by GC 308(1984)165

Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Amitrole
Determination of 3-amino-1,2,4-triazole (amitrole) in urine by ion-pair HPLC 303(1984)267

Ammonia
Determination of urea and ammonia using ion-pair LC with on-line post-column derivatization in an enzymatic solid-phase reactor 325(1985)255

Ammonium
Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Ammonium acetate
Ammonium acetate: a general purpose buffer for clinical applications of HPLC 316(1984)397

Amoxicillin
Isolation and structure elucidation of ampicillin and amoxicillin oligomers 303(1984)117

HPLC determination of amoxicillin in human plasma using a bonded-phase extraction 306(1984)429

Quantitative determination of amoxicillin and its decomposition products by HPLC 321(1985)441

cAMP-dependent protein kinase
Elution of the catalytic subunits of the type I and type II forms of cyclic AMP-dependent protein kinase within the type I chromatographic peak 312(1984)293

Amphetamine
Simple determination of *p*-hydroxyamphetamine by HPLC with electrochemical detection 306(1984)394

Determination of methamphetamine and its metabolites in rat tissues by GC with a nitrogen-phosphorus detector 318(1985)307

Amphetamines
HPLC retention data for 84 basic drugs of forensic interest on a silica column using an aqueous methanol eluent 301(1984)165

HPLC resolution of enantiomers of 1-phenyl-2-aminopropanes (amphetamines) with four chiral reagents 307(1984)335

Ampicillin
Isolation and structure elucidation of ampicillin and amoxicillin oligomers 303(1984)117

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

β-Amylase
Rapid procedure for purification of β-amylase from *Ipomea batata* 324(1985)489

Amylopectin
Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 319(1985)205

Amylose
Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 319(1985)205

Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

***p-tert.*-Amylphenol**
Determination of germicidal phenols in blood by capillary column GC 307(1984)420

Anabolic steroids
Fast HPLC screening method for the presence of trenbolone and its major metabolite in urine of slaughter cattle 319(1985)436

Anacardic acid
Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

Anaesthetic drugs
HPLC systems for the separation of local anaesthetic drugs with applicability to the analysis of illicit cocaine samples 301(1984)155

Androgen receptors
High-performance size exclusion chromatography as a rapid method for the separation of steroid hormone receptors 307(1984)39

Androstenedione
Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Use of Sep-Pak cartridges for the recovery of compounds from scintillation fluids 325(1985)323

Angiotensin
Combined HPLC-radioimmunoassay for the characterization and quantitative measurement of neuropeptides 312(1984)273

Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Separation and purification of diastereomers of angiotensin I by weak anion-exchange HPLC 322(1985)117

Use of benzoin as pre-column fluorescence derivatization reagent for the HPLC of angiotensins 322(1985)473

Anhydrotetracycline
Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303

Anilines
Application of reversed-phase HPLC to in vitro drug metabolism studies with N-alkylarylamines 309(1984)339

Use of cation exchangers for the on-line preconcentration of polar anilines in LC 317(1984)557

Anion exchangers
Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157

Anions
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography; factors that control retention and selectivity 301(1984)25

Methods for the analysis of inorganic anions. IV. Reversed-phase HPLC with aqueous hydrophobic ion pairs as eluents 302(1984)15

Simultaneous determination of anions and metal cations by single-column ion chromatography with ethylenediaminetetraacetate as eluent and conductivity and UV detection 312(1984)327

Single injection ion chromatographic analysis of both anions and cations 312(1984)456

Isotachophoresis in inorganic chemistry (Review) 313(1984)189

Anion chromatography on a conventional reversed-phase column using two UV detectors 314(1984)429

Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

UV visualization of inorganic anions by reversed-phase ion-interaction chromatography: factors that control sensitivity and detection 316(1984)211

Studies on sample preconcentration in ion chromatography. I. Design of an automated, single pump preconcentration system with direct UV absorbance detection 318(1985)279

Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Isotachophoretic determination of anionic and cationic species in well and surface waters 320(1985)199

Ion chromatography of inorganic anions with potentiometric detection using a metallic copper electrode 321(1985)363

Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Elimination of the peak of a major ion in anion chromatography with UV detection 325(1985)327

Anisic acid
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Anisoles
GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

Anserine
Rapid ion-exchange method for the determination of 3-methylhistidine in rat urine and skeletal muscle 311(1984)69

Anthocyanidins
Simultaneous analysis of anthocyanins and anthocyanidins on cellulose thin layers 318(1985)450

Anthocyanins
Simultaneous analysis of anthocyanins and anthocyanidins on cellulose thin layers 318(1985)450

Anthracene
New schemes for the electron-capture sensitization of aromatic hydrocarbons 302(1984)243

Anthracyclines
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

Continuous extraction of urinary anthracycline antitumor antibiotics with the horizontal flow-through coil planet centrifuge 307(1984)323

Anthraquinones
Analysis of naturally occurring reduced anthraquinones obtained from *Alternaria porri* (Ellis) Ciferri by HPLC 318(1985)139

Anthraquinone sulphonic acids
Separation of anthraquinone sulphonic acids by ion-pair LC 319(1985)319

9-Anthroic acid
Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

Antibiotics
Isolation and structure elucidation of ampicillin and amoxicillin oligomers 303(1984)117

Use of perfluorinated carboxylic acids in the separation of aminoglycoside antibiotics by ion-pair reversed-phase HPLC 303(1984)277

Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

HPLC analysis of cefotetan epimers in human plasma and urine 305(1984)363

Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373

HPLC determination of amoxicillin in human plasma using a bonded-phase extraction 306(1984)429

LC determination of amikacin in serum with spectrophotometric detection 307(1984)224

Continuous extraction of urinary anthracycline antitumor antibiotics with the horizontal flow-through coil planet centrifuge 307(1984)323

HPLC methods for the determination of cefpimizole in plasma and urine 308(1984)261

Reversed-phase HPLC method to determine ceftriaxone in biological fluids 309(1984)209

Determination of imipenem (N-formimidoyl thienamycin) in human plasma and urine by HPLC, comparison with microbiological methodology and stability 310(1984)71

Reversed-phase paired-ion HPLC method for the separation and quantification of multiple bleomycin congeners 310(1984)361

New rapid method of analysis of cefoxitin in serum and bone, by HPLC 311(1984)135

Determination of a new orally active cephalosporin in human plasma and urine by HPLC using automated column switching 311(1984)339

Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303

Process scale-up of a β-lactam antibiotic purification by HPLC 316(1984)133

Screening for new microbial products by HPLC using a photodiode array detector 316(1984)487

Determination of penicillin G anmd cloxacillin residues in beef and pork tissue by HPLC 317(1984)311

Studies on residual antibacterials in foods. III. HPLC determination of penicillin G in animal tissues using an on-line pre-column concentration and purification system 318(1985)299

Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213

Quantitative determination of amoxicillin and its decomposition products by HPLC 321(1985)441

Reversed-phase HPLC separations of tetracycline derivatives using volatile mobile phases 321(1985)462

Pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene 322(1985)321

TLC investigation of the antimycin A antibiotic complex 322(1985)386

Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

LC separation of aminoglycosides with pulsed amperometric detection 324(1985)407

Improvement of chemical analysis of antibiotics. VIII. Application of prepacked C_{18} Cartridge for the analysis of tetracycline residues in animal liver 325(1985)265

Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

Antidepressant drugs
Screening procedure for detection of antidepressants and their metabolites in urine using a computerized GC-MS technique 305(1984)309

Antigens
Purification of soluble specific antigens of systemic candidiasis by antibody affinity chromatography 306(1984)117

Antimony *d*-tartrate
Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59

Antimycin A
TLC investigation of the antimycin A antibiotic complex 322(1985)386

Anti-oxidants
Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141

TLC of age resisters for elastomers (phenols and organic phosphites) (in French) 314(1984)349

Micro column LC-MS using a capillary interface 323(1985)89

Antipyrine
Simple reversed-phase HPLC determination of antipyrine in rabbit plasma for pharmacokinetic studies 307(1984)475

Simultaneous determination of antipyrine and metabolites in human plasma and urine by HPLC 309(1984)219

Aphidicolin glycinate
Analysis of candidate anticancer drugs by thermospray HPLC-MS 323(1985)383

Apolipoproteins
Resolution and quantitation of apolipoproteins A-I and A-II from human high-density lipoprotein by size exclusion HPLC 311(1984)79

Aptazapine
Assay for the determination of the tetracyclic antidepressant compound aptazapine in plasma by HPLC 306(1984)412

Arachidic acid
Electrokinetic detection of lauric, myristic, stearic and arachidic acids separated by reversed-phase HPLC 321(1985)433

Arachidonic acid
Rapid, sensitive method for detecting different arachidonic acid metabolites by TLC: the use of autoradiography 307(1984)185

Simple single-step HPLC method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 308(1984)316

Arbaprostil
Silver-modified mobile phase for normal-phase LC determination of prostaglandins and their 5,6-*trans* isomers in prostaglandin bulk drugs and triacetin solutions 321(1985)353

Archive retrieval
Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Area under chromatographic peaks
Analytical solution for the area under chromatographic peaks obtained from the deconvolution method of Vaidya and Hester 303(1984)177

Arenarioside
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Arginine
Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

Aromatic acids
HPLC method for screening disorders of aromatic acid metabolism using a multi-detection system 310(1984)273

Chromatography of aromatic acids on ion exchangers 316(1984)261

Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267

Aromatic compounds
Unmodified and acylated cyclodextrin stationary phases for LC separation of aromatic compounds 301(1984)345

Clay CC for optical resolution. Resolution of aromatic compounds on a Λ-Ru(1,10-phenanthroline)$_3^{2+}$-montmorillonite column 319(1985)299

Determination of laccase activity with various aromatic substrates by HPLC 320(1985)435

Aromatic hydrocarbons
Combination of automated pre-column trace enrichment and on-line evaporation in HPLC 301(1984)335

New schemes for the electron-capture sensitization of aromatic hydrocarbons 302(1984)243

Basis of retention in normal-phase HPLC with cyano-propyl columns 303(1984)1

α-Arylalkylamine-derived chiral stationary phases
α-Arylalkylamine-derived chiral stationary phases. Evaluation of urea linkages 322(1985)295

Aryl β-diketonates
Volatility studies of metal chelates. VI. GC behaviour of aryl β-diketones and their metal chelates 322(1985)1

Volatility studies of metal chelates. VII. GC behaviour of chelates derived from the aromatic β-diketones 1-phenylbutane-1,3-dione and 4,4,4-trifluoro-1-(pentafluorophenyl)butane-1,3-dione 322(1985)23

Aryl β-diketones
Spectroscopic properties and GC behaviour of aryl β-diketones and selected chelates 312(1984)109

Arylnaphthalene lignan lactones
Solid-state fluorodensitometric quantitation of arylnaphthalene lignan lactones of *Cleistanthus collinus* 303(1984)296

Ascorbic acid
Fluorometric determination of total vitamin C in whole blood by HPLC with pre-column derivatization 305(1984)53

Analysis of uronic and aldonic acids, their lactones, and related compounds by HPLC on cation-exchange resins 319(1985)159

Ascorbic acid oxidase
Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Ascorbic acid-2-sulphate
Quantitative analysis of ascorbic acid-2-sulfate by HPLC with electrochemical detection 308(1984)306

Asparagine
GLC assay for asparagine and glutamine 322(1985)228

Aspartame
Isocratic LC method for the simultaneous determination of aspartame and other additives in soft drinks 303(1984)256

Astaxanthins
Direct diastereomeric resolution of carotenoids. I. 3-Hydroxy-4-oxo β-end group 318(1985)122

Astra Blue
Chromatography of Astra Blue dyes with eluents containing potassium thiocyanate in organic solvents 319(1985)400

Astromicin
Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373

Atmospheric-pressure microwave-induced helium plasma emission spectroscopy
Atmospheric-pressure microwave-induced helium plasma spectroscopy for simultaneous multielement GC detection 302(1984)277

Atratone
Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Atrazine
Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Atropine
Qualitative and quantitative analysis of hyoscyamine-atropine and scopolamine in mother tinctures of Solanaceae by HPLC (in French) 319(1985)412

Autofocusing
Autofocusing: a method for isoelectric focusing without carrier ampholytes 320(1985)15

Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213

Avidin
Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Azaarenes
On-line pre-concentration and LC of azaarenes using nitroaromatic sorbents 312(1984)303

Separation of methyl-substituted benz[c]acridines by GLC and HPLC 319(1985)331

Azintamide
GLC determination of azintamide (Ora-gallin) in pharmaceutical formulations 324(1985)475

Azobenzene
Determination of 1,3-diphenyltriazene and azobenzene in D&C Red No. 33 by solvent extraction and reversed-phase HPLC 321(1985)185

Azodisalicylate
Determination of sodium azodisalicylate, salazosulphapyridine and their metabolites in serum, urine and faeces by HPLC 305(1984)470

Azolastone
Analysis of candidate anticancer drugs by thermospray HPLC-MS 323(1985)383

B

***Bacillus* species**
Studies on volatile metabolites of some potentially pathogenic *Bacillus* species, using automated head-space GC 305(1984)178

Baclofen
Improved GLC method for the determination of baclofen in plasma and urine 310(1984)319

Baeocystin
Qualitative and quantitative determinations of hallucinogenic components of psilocybe mushrooms by reversed-phase HPLC 312(1984)467

Balaban indices
Calculation of retention indices by molecular topology. Chlorinated benzenes 319(1985)1

Ball pen inks
Use of a rotating disc multiwavelength detector operating in the visible region of the spectrum for monitoring ball pen inks separated by HPLC 303(1984)211

Band broadening
Speed of temperature increase when using large retention gaps in capillary GC 301(1984)1

Band broadening in space
Band broadening in space in splitless injection 324(1985)251

Band width
Effect of sample volume on peak width in high-performance gel filtration chromatography 324(1985)422

Barbiturates
Separation of underivatised barbiturates by capillary column GC. Preparation of medium polar polymethylsiloxane columns to optimize selectivity 318(1985)255

HPTLC on amino-bonded silica gel: application to barbiturates and steroids 322(1985)149

Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444

Bases
Effects of various strengths of acids on the electrophoretic behavior of weak bases in a non-aqueous solvent 314(1984)65

Basic compounds
Retention-optimization strategy for the HPLC ion-pair separation of samples containing basic compounds 316(1984)241

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. I. Relationship between capacity factors and the methanol concentration in methanol-water eluents 320(1985)293

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. II. Procedure for the determination of a lipophilic index measured by reversed-phase HPLC 320(1985)305

Basic drugs
HPLC retention data for 84 basic drugs of forensic interest on a silica column using an aqueous methanol eluent 301(1984)165

Development of a standardized analysis strategy for basic drugs, using ion-pair extraction and HPLC. VI. Drug level determination in saliva 309(1984)329

HPLC analysis of basic drugs on silica columns using non-aqueous ionic eluents. I. Factors influencing retention, peak shape and detector response 323(1985)173

HPLC analysis of basic drugs on silica columns using non-aqueus ionic eluents. II. Application of UV, fluorescence and electrochemical oxidation detection 323(1985)191

Bence-Jones proteins
Comparison of chromatographic characteristics of a series of homologous Bence-Jones proteins during size-exclusion chromatography by HPLC and by Sephadex 323(1985)418

Benomyl
Determination of benomyl and its metabolites by cation-exchange HPLC 317(1984)527

Benoxaprofen
Activated α-alkyl-α-arylacetic acid enantiomers for stereoselective TLC and HPLC determination of chiral amines 307(1984)145

α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167

Benperidol
Determination of benperidol in human plasma by HPLC 311(1984)109

Benz[c]acridines
Separation of methyl-substituted benz[c]acridines by GLC and HPLC 319(1985)331

Benzaldehyde
Simple HPLC assay for benzylamine oxidation products in cell suspensions 305(1984)256

Benzamidine
New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188

Benz[a]anthracene
Direct resolution of mono- and diol enantiomers of unsubstituted and methyl-substituted benz[a]anthracene and benzo[a]pyrene by HPLC with a chiral stationary phase 316(1984)569

Benzene
Use of LC with dual-electrode electrochemical detection in the investigation of glutathione oxidation during benzene metabolism 317(1984)579

Nickel(II)-exchanged molecular sieve 13X as a column packing for the GC determination of thiophene in benzene with flame ionization detection 325(1985)430

Benzene derivatives
Indirect UV detection of non-absorbing solutes in reversed-phase HPLC with the help of UV-active, non-ionic mobile phase constituents 312(1984)345

Preparation and retention behaviour of chemically bonded methylated-cyclodextrin stationary phases for LC 314(1984)193

Computer-assisted retention prediction system for reversed-phase micro HPLC 316(1984)1

Benzils
GLC of substituted benzils and benzophenones 324(1985)450

Benzoates
Simple HPLC assay for benzylamine oxidation products in cell suspensions 305(1984)256

GLC analyses. XXXVII. Capillary column studies of benzoyl and monochlorobenzoyl esters of lower saturated branched-chain alcohols 323(1985)331

GLC Analyses. XXXVIII. Capillary column studies of unsaturated esters of benzoic and monochlorobenzoic acids 324(1985)113

GLC analyses. XLI. Ethyl and ω-chloroethyl esters of benzoic and monochlorobenzoic acids on SE-30 and OV-351 capillary columns 325(1985)433

1,4-Benzodiazepine glucuronides
Separation, isolation and identification of optical isomers of 1,4-benzodiazepine glucuronides from biological fluids by reversed-phase HPLC 306(1984)231

Benzodiazepines
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Benzoic acid
Determination of preservatives in cosmetic products by reversed-phase HPLC 315(1984)465

Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

GLC analyses. XXXV. Capillary column studies of C_1-C_{12} *n*-alkyl esters of benzoic and monochlorobenzoic acids 322(1985)83

Benzoin
HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281

HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257

Use of benzoin as pre-column fluorescence derivatization reagent for the HPLC of angiotensins 322(1985)473

Benzo[*ghi*]perylene
Products obtained during combustion of polymers under simulated incinerator conditions. II. Polystyrene 315(1984)201

Benzophenones
GLC of substituted benzils and benzophenones 324(1985)450

Benzo[*a*]pyrene
Direct resolution of mono- and diol enantiomers of unsubstituted and methyl-substituted benz[*a*]anthracene and benzo[*a*]pyrene by HPLC with a chiral stationary phase 316(1984)569

Benzoquinoline
Effects of various strengths of acids on the electrophoretic behavior of weak bases in a non-aqueous solvent 314(1984)65

***p*-Benzoquinones**
HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. I. Interplay of on-column redox reaction and the chromatographic retention process 316(1984)151

Benzyl alcohol
Simple HPLC assay for benzylamine oxidation products in cell suspensions 305(1984)256

Benzylamine
Simple HPLC assay for benzylamine oxidation products in cell suspensions 305(1984)256

***o*-Benzyl-*p*-chlorophenol**
Determination of germicidal phenols in blood by capillary column GC 307(1984)420

Bepridil
Determination of bepridil in biological fluids by HPLC 309(1984)125

Bestatin
Determination of bestatin in serum by HPLC with fluorescence detection 305(1984)381

Betaxolol
Simple method for routine determination of betaxolol in blood and urine by automated HPLC with fluorimetric detection 311(1984)199

Bevantolol
GC determination of bevantolol in plasma 308(1984)345

Bile acids
Selective LC isolation and GC-MS analysis of ketonic bile acids in faeces 302(1984)1

Laser fluorimetry for capillary column LC: high-sensitivity detection of derivatized biological compounds 317(1984)443

Binary adsorption isotherms
GC determination of binary adsorption isotherms 302(1984)195

Biotin
Use of LC-MS for the quantitation of dethiobiotin and biotin in biological samples 303(1984)272

Biphenyl polysiloxane
Polarizable polysiloxane stationary phases for capillary column GC 302(1984)303

Bipolar pulsed electron-capture detector
Bipolar pulsed microvolume electron-capture detector for capillary GC 302(1984)221

[1,1-Bis(aminomethyl)cyclohexane]platinum(II) complexes
Analysis of antitumour [1,1-bis(aminomethyl)cyclohexane]platinum(II) complexes derived from spiroplatin by HPLC with differential pulse amperometric detection 320(1985)379

1,3-Bis(2-chloroethyl)-1-nitrosourea
Quantitation of 1,3-bis(2-chloroethyl)-1-nitrosourea in plasma using HPLC 305(1984)496

Bis(malonato)ethylenediaminechromate(III)
HPLC in the study of some reactions of chromium(III) complexes 314(1984)387

Bis(2,4-pentanedionato)[2-(methylseleno)ethylamine]cobalt(III)
Chromatographic separation of the diastereomers of the bis(2,4-pentanedionato)[2-(methylseleno)ethylamine]cobalt(III) ion on a column of sulphoethyl-Toyopearl and determination of the rate of inversion at the selenium atom 301(1984)241

Bleomycin
Reversed-phase paired-ion HPLC method for the separation and quantification of multiple bleomycin congeners 310(1984)361

Blood plasma
Procedures for direct injections of untreated blood plasma into LC columns with emphasis on a pre-column venting technique 317(1984)213

Bonded phases
Comparison of chromatographic behaviour of silica and chemically bonded phases in supercritical fluid chromatography 315(1984)31

Borates
Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

Boric acid
Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

Boric acid complexes
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

B proteins
Separation and quantitative assay of three pathogenesis-related (b)proteins from tobacco mosaic virus hypersensitive *Nicotiana tabacum* by reversed-phase HPLC 318(1985)417

Bradykinin
Detection and quantitation of fluorescamine-labeled bradykinin, its analogues and metabolites using HPLC 306(1984)109

Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Branched-chain amino acid transaminase
Affinity chromatography of B_6-vitamin-dependent enzymes: purification of pig-heart branched-chain amino acid transaminase 318(1985)333

Brodifacoum
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Bromadiolone
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Bromazepam
Determination of bromazepam in plasma and of its main metabolites in urine by reversed-phase HPLC 310(1984)129

Bromide
Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Bromine
Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Bromoacetaldehyde
Sensitive fluorimetry of adenine-containing compounds with HPLC 309(1984)63

Bromopride
Determination of bromopride in human plasma and urine by HPLC 310(1984)353

Bromotetrachlorophenol
Identification of bromotetrachlorophenol in commercial pentachlorophenol samples 314(1984)476

Brompheniramine
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Bubble flow meter
Diffusion in the bubble flow meter in inverse GC experiments 301(1984)261

Bufetonin
HPLC determination of hallucinogenic indoleamines with simultaneous UV photometric and voltammetric detection 320(1985)414

Buffers
Ammonium acetate: a general purpose buffer for clinical applications of HPLC 316(1984)397

Bunitrolol
Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

Bupivacaine
HPLC determination of bupivacaine in human serum 309(1984)369

Determination of bupivacaine in human plasma by HPLC 311(1984)218

Bupranolol
Determination of the beta-adrenoceptor blocking drug bupranolol in plasma and tissues of the rat by HPLC with UV detection 309(1984)193

Determination of carboxybupranolol, the major metabolite of bupranolol, in human plasma by HPLC 311(1984)227

Butane
Characterization of adsorption at kaolinite surfaces by heterogeneous GSC 314(1984)111

Butanoates
GC of homologous esters. XXVIII. Retention increments of aliphatic C_1-C_{18} *n*-alkyl esters of butanoic acid and its monochloro derivatives on SE-30 and OV-351 capillary columns 320(1985)325

GC of homologous esters. XXXI. Butanoyl and monochlorobutanoyl esters of lower saturated branched chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 325(1985)61

Butanol
HPLC of low-molecular-weight aliphatic alcohols using indirect photometric detection 314(1984)488

***tert.*-Butanol**
Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.*-butanol 319(1985)247

Butene
Characterization of adsorption at kaolinite surfaces by heterogeneous GSC 314(1984)111

Butylamine
Determination of butylamine in air samples by isotachophoresis 312(1984)489

Butyl-Sepharose 4B
Quantitative determination of the ligand content and the negative and positive charges on butylamine-substituted Sepharose 4B 321(1985)335

2-*sec.*-Butylthiazoline
Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23

C

Cadaverine
Sensitive assay for diamine oxidase activity using HPLC 309(1984)151

Cadralazine
HPLC separation of cadralazine from its potential metabolites and degradation products. Quantitation of the drug in human plasma and urine 310(1984)139

Caeruloplasmin
Purification of glycopeptides of human plasma proteins by HPLC 317(1984)11

Caesium
Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Caffeine
Simultaneous determination of caffeine and its N-demethylated metabolites in umbilical cord plasma using HPLC 306(1984)191

Isothermal GC analysis of diphenhydramine after direct injection onto a fused-silica capillary column 307(1984)295

Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107

Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Determination of caffeine in tea by HPLC and a modified digestion procedure 317(1984)403

Simultaneous HPLC assay of acetylsalicylic acid and salicylic acid in film-coated aspirin tablets 317(1984)507

Use of microbore HPLC for the determination of caffeine, theobromine and theophylline in cocoa 318(1985)408

Quantitation of caffeine in doping analysis using an automated extractor/concentrator 325(1985)304

Caffeoylquinic acid
Chromatographic separation and identification of naturally occurring chlorogenic acids by ^1H nuclear magnetic resonance spectroscopy and MS 315(1984)253

Calcium
Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Calibration
Experimental evaluation of the errors in high-performance GPC measurements 325(1985)37

Calibration marker scheme
Evaluation of a calibration marker scheme for open tubular column GC with on-column injection and electron-capture detection 302(1984)289

Candidiasis antigens
Purification of soluble specific antigens of systemic candidiasis by antibody affinity chromatography 306(1984)117

Capillary column gas chromatography-field ionization mass spectrometry
Thick-film capillary column GC-field ionization MS. A complementary technique for the rapid analysis of volatiles 322(1985)200

Capillary columns
see also Columns and microbore columns

Method for testing retention gaps and data on the GC retention power of differently deactivated capillary tubes 323(1985)237

Capillary interfaces
Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Capsaicin
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288

Capsaicinoids
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288

Captopril
Combined GC-MS procedure for the measurement of captopril and sulfur-conjugated metabolites of captopril in plasma and urine 305(1984)83

Carbamates
Analysis of selected pesticides by HPLC-MS 312(1984)221

Application of electrochemical detection to the determination of residues of some carbamate insecticides by HPLC (in French) 325(1985)239

Carbamazepine
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Development of a standardized analysis strategy for basic drugs, using ion-pair extraction and HPLC. VI. Drug level determination in saliva 309(1984)329

Automated analysis of antiepileptic drugs in serum by column-switching HPLC 309(1984)357

Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Carbapenem antibiotics
Determination of imipenem (N-formimidoyl thienamycin) in human plasma and urine by HPLC, comparison with microbiological methodology and stability 310(1984)71

Carbohydrates
see also Sugars and saccharides

Analysis of carbohydrates in lens, erythrocytes, and plasma by HPLC of nitrobenzoate derivatives 307(1984)180

Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Carbohydrates of whole defatted cells as a basis for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)31

HPLC of disaccharides on amine-bonded silica columns 319(1985)51

Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 319(1985)205

Retention of carbohydrates on silica and amine-bonded silica stationary phases: application of the hydration model 325(1985)287

β-Carbolines
Analysis of β-carbolines by reversed-phase ion-pair partition chromatography with fluorometric detection 319(1985)23

Carbon analysis
Carbon analysis for the study of bonded phases in LC 321(1985)319

Carbon black
GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19

Carbonic anhydrase
Affinity chromatography of membrane fragments. Separation of carbonic anhydrase-enriched and -depleted brain membranes 312(1984)435

Rapid ion-exchange chromatography for preparative separation of proteins. Application to porcine and bovine carbonic anhydrases 319(1985)59

Carbonyl compounds
HPLC of carbonyl compounds as 2,4-dinitrophenylhydrazones with electrochemical detection 318(1985)427

Determination of carbonyl compounds in an Australian (Rundle) shale oil 319(1985)230

Carbonyl sulphide
GC study of the analysis and elution mechanism of hydrogen sulphide, carbonyl sulphide and light mercaptans in petroleum gases 321(1985)59

Carboplatin
Analysis of platinum complexes by LC with quenched phosphorescence detection 312(1984)337

Carboprost
HPLC determination of acid-catalyzed degradation products of methyl carboprost in a polymeric controlled-release device 314(1984)369

Carboxybupranolol
Determination of carboxybupranolol, the major metabolite of bupranolol, in human plasma by HPLC 311(1984)227

γ-Carboxyglutamic acid
Improved method for the determination of γ-carboxyglutamic acid in human urine by HPLC 308(1984)398

Analysis of γ-carboxyglutamic acid by GLC 319(1985)404

Carboxylates
Separation of sulfonate and carboxylate mixtures by ion-exchange HPLC 302(1984)65

Carboxylic acid methyl esters
Separation and double-bond determination on nanogram quantities of aliphatic monounsaturated alcohols, aldehydes and carboxylic acid methyl esters 322(1985)484

Carboxylic acids
see also Organic acids

Analysis of selected pesticides by HPLC-MS 312(1984)221

Chromatographic separation of enantiomers of acids with quinine as chiral counter ion 316(1984)553

Derivatization, identification and separation of carboxylic acids in wines and beverages by HPLC 322(1985)177

ω-Carboxyl spacers
Introduction of ω-carboxyl spacers onto cross-linked agarose gel beads by O-alkylation for the preparation of affinity adsorbents 315(1984)161

Carboxymethylcellulose
Recovery of plant amines with carboxylic ion-exchange substrates and their subsequent analysis by TLC 312(1984)357

Carboxymethyldextrans
Preparative HPLC of proteins on ion-exchange columns with carboxymethyldextrans as displacers 316(1984)125

Cardanol
Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

Cardiac glycosides
Investigation of conditions of the preparative LC separation of cardiac glycosides present in *Digitalis lanata* 303(1984)179

Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

Cardol
Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

Carminomycin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

Carnosine
Rapid ion-exchange method for the determination of 3-methylhistidine in rat urine and skeletal muscle 311(1984)69

Carob
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

β-Carotene
Use of Sep-Pak cartridges for the recovery of compounds from scintillation fluids 325(1985)323

Carotenoids
Computer-optimized normal-phase HPLC separation of *Corynebacterium poinsettiae* carotenoids 310(1984)372

Direct diastereomeric resolution of carotenoids. I. 3-Hydroxy-4-oxo β-end group 318(1985)122

Carrageenan
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Carrier gas compressibility
Approximate approach to correction for carrier compressibility effects in coupled packed GC columns 323(1985)402

Carrier gases
Reaction GC-MS. VI. Use of deuterium as the carrier and reagent gas in reaction GC-MS 301(1984)77

Cashew nut-shell liquid
Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

Catalysts
Effect of activated sorption processes on the results of thermoprogrammed reduction investigations on Fe-Al_2O_3 catalysts 301(1984)270

Catecholamines
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

HPLC determination of plasma catecholamines during α-methyldopa therapy 308(1984)301

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Non-radiochemical procedure for the measurement of O-methylation of the stereoisomers of isoprenaline 310(1984)283

GC analysis of endogenous catecholamines, phenolic amines and derived isoquinolines

using short glass capillary columns and electron-capture detection 311(1984)17

Dihydrocaffeic acid: a common contaminant in the LC-electrochemical measurement of plasma catecholamines in man 311(1984)148

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

HPLC separations on dihydroxyboryl-agarose 316(1984)301

Selective retention of catecholamines and their derivatives in reversed-phase ion-pair partition chromatography 322(1985)43

Catechol-O-methyltransferase
Assay for catechol-O-methyltransferase in erythrocytes using a new fluorogenic substrate, 2-(3,4-dihydroxyphenyl)naphtho[1,2-*d*]thiazole 308(1984)93

Catechols
Studies of cyclic di-*tert.* -butylsilylene derivatives of diols and hydroxy acids by GC-MS 315(1984)119

Cation exchangers
Use of cation exchangers for the on-line preconcentration of polar anilines in LC 317(1984)557

Cations
Single injection ion chromatographic analysis of both anions and cations 312(1984)456

Isotachophoretic determination of anionic and cationic species in well and surface waters 320(1985)199

Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Application of indirect potentiometric detection with a metallic copper electrode to ion chromatography of transition metal ions 324(1985)319

Cefotetan
HPLC analysis of cefotetan epimers in human plasma and urine 305(1984)363

Cefoxitin
New rapid method of analysis of cefoxitin in serum and bone, by HPLC 311(1984)135

Cefpimizole
HPLC methods for the determination of cefpimizole in plasma and urine 308(1984)261

Ceftriaxone
Reversed-phase HPLC method to determine ceftriaxone in biological fluids 309(1984)209

Cellooligomers
Structural heterogeneity of cellooligomers homogeneous according to high-resolution size-exclusion chromatography 319(1985)311

Cells
Hydrophobic chromatography of cells: adsorption and resolution on homologous series of alkylagaroses 317(1984)103

Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213

Cellulose
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Cephalexin
LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Cephalosphorins
HPLC analysis of cefotetan epimers in human plasma and urine 305(1984)363

Cephalosporins
HPLC methods for the determination of cefpimizole in plasma and urine 308(1984)261

Reversed-phase HPLC method to determine ceftriaxone in biological fluids 309(1984)209

New rapid method of analysis of cefoxitin in serum and bone, by HPLC 311(1984)135

Determination of a new orally active cephalosporin in human plasma and urine by HPLC using automated column switching 311(1984)339

Process scale-up of a β-lactam antibiotic purification by HPLC 316(1984)133

Cetrimide
Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267

Cetyltrimethylammonium bromide
Factors influencing the reproducibility of isotachophoretic determinations of ions present in wet-process phosphoric acid 325(1985)161

Channels for sedimentation field flow fractionation
Sampling and flow effects in sedimentation field flow fractionation 315(1984)45

Charge
Determination of ionic charge by liquid-liquid partition 322(1985)425

Charge-transfer thin-layer chromatography
Masking effects in "charge-transfer" TLC 303(1984)190

Chemiluminescence detection
Solvent and pH effects on peroxyoxalate chemiluminescence detection for LC 314(1984)155

Chimassorb 944
Determination of polymeric hindered-amine light stabilisers in plastics by pyrolysis-GC 325(1985)87

Chiral additives
Enantiomeric resolution of an optically active guanine derivative by HPLC with phenylalanine-Cu(II) in the mobile phase. 303(1984)217

Enantiomeric analysis of a mixture of the common protein amino acids as their Dns derivatives. Single-analysis reversed-phase HPLC procedure using a chiral mobile phase additive 303(1984)244

Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Optical resolution of amino acid enantiomers by HPLC 316(1984)547

Chromatographic separation of enantiomers of acids with quinine as chiral counter ion 316(1984)553

Enantiomeric separation of underivatized α-methyl-α-amino acids by HPLC 318(1985)117

Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 319(1985)35

Chiral stationary phases
see also Stationary phases

Chiral stationary phases, GC
Direct enantiomeric resolution of mephenytoin and its N-demethylated metabolite in plasma and blood using chiral capillary GC 307(1984)121

Monitoring of *S*- and *R*-tocainide in human plasma after heptafluorobutyrylation, separation on Chirasil-Val and electron-capture detection 308(1984)181

Resolution of α-halogenocarboxylic acids by GC on (*R*)-N-lauroyl-α-(1-naphthyl)ethylamine 318(1985)235

GC separation of enantiomeric sulphur compounds on Chirasil-Val 320(1985)393

GC resolution of substituted glutamic acid enantiomers 325(1985)444

Chiral stationary phases, LC
Direct enantiomeric determination of (-)- and (+)-propranolol in human serum by HPLC on a chiral stationary phase 306(1984)405

Direct resolution of mono- and diol enantiomers of unsubstituted and methyl-substituted benz[*a*]anthracene and benzo[*a*]pyrene by HPLC with a chiral stationary phase 316(1984)569

Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

Direct LC separation of enantiomers on immobilized protein stationary phases. IV. Molecular interaction forces and retention behaviour in chromatography on bovine serum albumin as a stationary phase 316(1984)617

Separation of diastereomeric amides by preparative HPLC and analysis of enantiomers on a chiral support (in French) 318(1985)398

Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 319(1985)35

Clay CC for optical resolution. Resolution of aromatic compounds on a Λ-Ru(1,10-phenanthroline)$_3^{2+}$-montmorillonite column 319(1985)299

Preparative separation of enantiomers by flash chromatography 319(1985)392

Reversed-phase chromatographic resolution of N-(3,5-dinitrobenzoyl)-α-amino acids on chiral stationary phases 322(1985)287

α-Arylalkylamine-derived chiral stationary phases. Evaluation of urea linkages 322(1985)295

Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444

In-situ preparation of a chemically bonded chiral stationary phase for the separation of aromatic α-amino acid enantiomers 325(1985)103

Chloral hydrate
Chloral hydrate is compatible with isoelectric focusing 325(1985)333

Chloramphenicol
LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Chloranil
Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

Chlordiazepoxide
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Chlorhexidine
Improved method for the determination of chlorhexidine in urine 306(1984)303

Simultaneous determination of chlorhexidine, tetracaine and their degradation products by ion-pair LC 315(1984)457

Chloride
Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Chloride-attachment negative chemical ionization
LC-chloride-attachment negative chemical ionization MS 319(1985)273

Chlorinated carboxylic acids
Analysis of selected pesticides by HPLC-MS 312(1984)221

Chlorinated compounds
The mass spectrometer as a chlorine-selective chromatographic detector. I. Description and evaluation of the technique 314(1984)83

The mass spectrometer as a chlorine-selective chromatographic detector. II. Applications to chemical systems 314(1984)93

Chlorinated dibenzo-*p*-dioxins
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351

Characteristics of an extraction and purification procedure for chlorinated dibenzo-*p*-dioxins and dibenzofurans in soil and liver 312(1984)165

Chlorinated diphenyl ethers
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351

Chlorinated hydrocarbons
Determinations of chlorinated hydrocarbons by GC using response factor calibration 314(1984)243

Chlorinated 2-phenoxyphenols
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351

Chlorine
Production of mutagenic artifacts by the action of residual chlorine on XAD-4 resin 325(1985)95

Chloroacetanilides
Determination of chloroacetanilides and chloroanilines in sulphanilamides and waste water by HPLC 319(1985)99

ω-Chloroacetophenone
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b*,*f*]-1:4-oxazepine 315(1984)417

N-Chloroacetyl-6-aminohexanoic acid
Introduction of ω-carboxyl spacers onto cross-linked agarose gel beads by O-alkylation for the preparation of affinity adsorbents 315(1984)161

Chloroacetylglycylglycine
Introduction of ω-carboxyl spacers onto cross-linked agarose gel beads by O-alkylation for the preparation of affinity adsorbents 315(1984)161

Chloroalkanes
Calculation of retention indices by molecular topology: chlorinated alkanes 314(1984)1

Chloroanilines
Determination of chloroacetanilides and chloroanilines in sulphanilamides and waste water by HPLC 319(1985)99

Chloroanisoles
GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

GLC analyses. XXXIII. Retention increments for *o*-chloro and *o*-methoxy substitution in isomeric chloroanisoles 321(1985)467

Chlorobenzenes
Comparison of a new electrochemical detector for GC analysis with the electrolytic conductivity detector 302(1984)269

Calculation of retention indices by molecular topology. Chlorinated benzenes 319(1985)1

GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

Chlorobenzoates
GLC analyses. XLI. Ethyl and ω-chloroethyl esters of benzoic and monochlorobenzoic acids on SE-30 and OV-351 capillary columns 325(1985)433

***o*-Chlorobenzylidenemalononitrile**
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b*,*f*]-1:4-oxazepine 315(1984)417

Chlorobutanoates
GC of homologous esters. XXVIII. Retention increments of aliphatic C_1-C_{18} *n*-alkyl esters of butanoic acid and its monochloro derivatives on SE-30 and OV-351 capillary columns 320(1985)325

Chloro complexes
Separation of anionic chloro complexes by capillary tube isotachophoresis in non-aqueous systems 322(1985)139

Chlorodibenzofurans
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351

Characteristics of an extraction and purification procedure for chlorinated dibenzo-*p*-dioxins and dibenzofurans in soil and liver 312(1984)165

D-(-)-6-Chloro-1,5-dihydro-3-methylimidazo(2,1-*b*)quinazolin-2(3H)-one
Determination of the imidazo quinazoline derivative Ro 13-6438 in biological fluids by HPLC 305(1984)153

ω-Chloroethanols
GLC analyses. XXXIX. ω-Chloroethanols on low-polarity (SE-30) and polar (OV-351) capillary columns 324(1985)181

Chloroethyl alkanoates
GLC analyses. XXXVI. Separation of ethyl, 2-chloroethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns with temperature programming 322(1985)97

Chloroform
Mixed solvents in GLC. Activity coefficients for chloroform, dichloromethane and ethyl acetate in squalane-dinonyl phthalate mixtures at 303°K 324(1985)231

Chlorogenic acids
Chromatographic separation and identification of naturally occurring chlorogenic acids by ^1H nuclear magnetic resonance spectroscopy and MS 315(1984)253

4-Chloro-3-methylphenol
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Chlorophacinone
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Chlorophenols
GLC analyses. XXIX. Separation of free chlorophenol isomers on non-polar and polar quartz capillary columns 303(1984)197

Identification of bromotetrachlorophenol in commercial pentachlorophenol samples 314(1984)476

GLC analyses. XXXI. Retention increments of isomeric chlorophenols on low-polarity (SE-30) and polar (FFAP) capillary columns 315(1984)185

GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

GLC analyses. XL. Incremental effects of a formyl group introduced into the *ortho*- and *para*-positions of isomeric chlorophenols 324(1985)192

Chlorophenyl acetates
GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

Chlorophyll
HPLC study of the chlorophyll allomerization reaction 316(1984)25

Chloropropanoates
GC of homologous esters. XXVII. Retention increments of C_1-C_{18} primary alkanols and their 2-chloropropanoyl and 3-chloropropanoyl derivatives on SE-30 and OV-351 capillary columns 319(1985)131

Chloroquine
Improved GC method for the simultaneous determination of chloroquine and two metabolites using capillary columns 306(1984)147

Separation and identification of a plasma and urinary mono-acetylated conjugate of chloroquine in man by ion-pair HPLC 309(1984)426

Chlorphenesin
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Chlorpheniramine
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Chlorpromazine
Isotopic fractionation of N-([^{3}H]methyl)-chlorpromazine and N-([^{3}H]methyl)-7-hydroxychlorpromazine by reversed-phase HPLC 303(1984)412

Chlorpyrifos
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Chlortetracycline
Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303

Improvement of chemical analysis of antibiotics. VIII. Application of prepacked C_{18} Cartridge for the analysis of tetracycline residues in animal liver 325(1985)265

Cholecystokinins
Purification of a cholecystokinin fragment peptide in the horizontal flow-through coil planet centrifuge 301(1984)277

Reversed-phase HPLC and separation of peptide hormones in human tissues and plasma with reference to cholecystokinins 305(1984)43

Cholesterol
Simultaneous determination by HPLC of tocopherol isomers, α-tocopheryl quinone, and cholesterol in red blood cells and plasma 306(1984)371

Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65

Faecal lipid chromatography. I. Quantitative determination with Chromarods 310(1984)297

Liquid-liquid extraction of membranes from calf brain using conventional and centrifugal counter-current distribution techniques 311(1984)277

Cholesterol-*p*-chlorocinnamate
Aryl substituted diastereomeric alkenes: GC behavior on a non-polar *versus* a liquid crystal phase 321(1985)127

Cholesteryl esters
Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65

Choline
Chromatographic analysis of choline and acetylcholine by UV visualization 319(1985)454

Determination of choline and acetylcholine levels in rat brain regions by LC with electrochemical detection 322(1985)191

Choline kinase
Simultaneous assay of choline kinase and choline oxidase in tissue by high-performance cation-exchange chromatography and continuous radioactive detection 324(1985)203

Choline oxidase
Simultaneous assay of choline kinase and choline oxidase in tissue by high-performance cation-exchange chromatography and continuous radioactive detection 324(1985)203

Chromarods
Enhanced response of more highly unsaturated lipids on TLC-Iatroscan flame ionization detection after exposing the developed Chromarods to iodine vapour 319(1985)215

Silica gel particle size and coating thicknesses of Chromarods-S and -SII 323(1985)410

Chromatogram evaluation
Use of histograms in computer-aided comparison of chromatograms 318(1985)247

Chromatotron
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Chromium
GC studies on group VIB organometallic complexes. The analytical and preparative separation of chromium, molybdenum and tungsten trifluorophosphine carbonyl complexes 301(1984)253

Chromium complexes
HPLC in the study of some reactions of chromium(III) complexes 314(1984)387

Chrysanthemates
GC and GC-MS analysis of diastereomeric acyclic isoprenoid esters using fused-silica capillary columns 312(1984)395

Chymosin
Affinity chromatography of kid chymosin on histidyl-Sepharose 303(1984)285

Chymotrypsin
Purification of chymotrypsin and trypsin by CC on agar gel particles 303(1984)283

Chymotrypsinogen
Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

α-Chymotrypsinogen-A
Effects of urea-thermal denaturation on the high-performance cation-exchange chromatography of α-chymotrypsinogen-A 314(1984)337

Ciglitazone
Isomodal column switching HPLC technique for the analysis of ciglitazone and its metabolites in human serum 307(1984)155

Cimetidine
HPLC determination of cimetidine and related compounds (raw material and tablets) 319(1985)235

Cinchona alkaloids
Inter-laboratory study on the separation of cinchona alkaloids by HPLC 318(1985)362

Cinnamaldehyde
HPLC fluorometric determination of cinnamaldehyde in perfume, cologne and toilet water 317(1984)421

Ciprostene
HPLC method for the separation of isomers of 9β-methylcarbacyclin (ciprostene) and the analysis of pharmaceutical samples 321(1985)209

Cisplatin
Analysis of platinum complexes by LC with quenched phosphorescence detection 312(1984)337

Quantitative analysis of *cis*-dichlorodiammineplatinum(II) by high-performance TLC 315(1984)481

Citalopram
HPLC determination of citalopram and four of its metabolites in plasma and urine samples from psychiatric patients 308(1984)199

Class N solvents
Influence of water in LSC systems on retention data. II. Fixation of monolocalized solutes with class N solvents and silica gels 316(1984)373

Cleistanthin
Solid-state fluorodensitometric quantitation of arylnaphthalene lignan lactones of *Cleistanthus collinus* 303(1984)296

Clomipramine
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Clonazepam
Quantitation of clonazepam and its 7-amino and 7-acetamido metabolites in plasma by HPLC 306(1984)241

Cloxacillin
Determination of penicillin G anmd cloxacillin residues in beef and pork tissue by HPLC 317(1984)311

CN
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b,f*]-1:4-oxazepine 315(1984)417

Coating thickness
Silica gel particle size and coating thicknesses of Chromarods-S and -SII 323(1985)410

Cobalt complexes
Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59

Reversed-phase high-performance ion-pair chromatography of cobalt(III) coordination complexes (Review) 313(1984)93

Paper electrophoretic study of ion pair formation. XV. Resolution of optically active cobalt(III)-cyclohexanediamine complexes 318(1985)440

Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 319(1985)35

Cobalt(III)-iminodiacetate
Developments in the HPLC of metallo-organic compounds 317(1984)569

Cocaine
HPLC systems for the separation of local anaesthetic drugs with applicability to the analysis of illicit cocaine samples 301(1984)155

Determination of cocaine in plasma by automated GC 306(1984)173

Codeine
Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Colchiceine
Separation of colchicine and related hydrolysis and photodecomposition products by HPLC, using copper ion complexation 315(1984)233

Colchicine
Separation of colchicine and related hydrolysis and photodecomposition products by HPLC, using copper ion complexation 315(1984)233

Collagen
Separation of collagen α-chains by reversed-phase HPLC. Comparison of column alkyl stationary phases and temperature effects 308(1984)111

Rapid and sensitive amino acid analysis of human collagens using HPLC 311(1984)375

TLC of hydroxylysine for collagen analysis 311(1984)400

Collection efficiency
Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448

Collinusin
Solid-state fluorodensitometric quantitation of arylnaphthalene lignan lactones of *Cleistanthus collinus* 303(1984)296

Colour additives
Determination of synthetic organic colours in lipsticks by TLC and HPLC 318(1985)149

Column bleeding
GC-MS analysis of compounds generated upon thermal degradation of some stationary phases in capillary GC 302(1984)95

Column calibration
Semi-micro size-exclusion chromatography of polymers. Calibration of columns and comparison with conventional columns 320(1985)343

Column connector
Miniature connector with a low dead volume for fused-silica capillary columns 301(1984)454

Column coupling
General theory for correction for compressibility effects in binary coupled GC columns and the procedure for window diagram optimisation of relative lengths 321(1985)249

Column dead time
Use of retention plots of *n*-alkyl benzenes for determining dead times in LC and GC 321(1985)287

Column efficiency
Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448

Role of extra-column volume in the efficiency of high-speed LC 321(1985)69

Column life-time
Column lifetime in fast protein LC 314(1984)396

Columns
see also Capillary columns and microbore columns

Protein separations on reversed-phase HPLC minicolumns 317(1984)119

Preparation of efficient and stable reversed-phase microbore columns for HPLC 321(1985)450

Column switching
Isomodal column switching HPLC technique for the analysis of ciglitazone and its metabolites in human serum 307(1984)155

Automated analysis of antiepileptic drugs in serum by column-switching HPLC 309(1984)357

Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Role of column switching in semi-preparative LC. Isolation of the sweetener stevioside 316(1984)105

Influence of solute polarity in column-switching chromatography for the assay of drugs in plasma and urine 317(1984)493

Single-valve, single-pump, HPLC column-switching analysis of hydrolyzed wood components 318(1985)269

Column temperature
Separation of collagen α-chains by reversed-phase HPLC. Comparison of column alkyl stationary phases and temperature effects 308(1984)111

Mobile phase effects in reversed-phase chromatography. VII. Dependence of retention on mobile phase composition and column temperature 318(1985)1

Complex formation
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11

Complex-forming compounds
Uranyl salts as fluorescence reagents for the TLC detection of complex-forming compounds (in German) 322(1985)272

Compressibility effects
General theory for correction for compressibility effects in binary coupled GC columns and the procedure for window diagram optimisation of relative lengths 321(1985)249

Approximate approach to correction for carrier compressibility effects in coupled packed GC columns 323(1985)402

Computer-assisted chromatography
Versatile microcomputer-controlled, automated gradient analytical HPLC system 316(1984)407

Use of histograms in computer-aided comparison of chromatograms 318(1985)247

Computer programs
Quantitative densitometric method for the rapid separation and quantitation of the major lipids of tissues and lipoproteins by HPTLC. II. Reduction of the densitometric data 307(1984)81

The mass spectrometer as a chlorine-selective chromatographic detector. I. Description and evaluation of the technique 314(1984)83

Computer simulation
Dynamics of isotachophoretic separation. I. Computer simulation 320(1985)167

Dynamics of isotachophoretic separation. II. Impurities in the electrolytes and sample injection 320(1985)175

Concentration effects
Characterization of extracolumn and concentration-dependent distortion of chromatographic peaks by Edgeworth-Cramér series 315(1984)67

Concentration-pulse chromatography
GC determination of binary adsorption isotherms 302(1984)195

Concentration techniques
Development of an on-line high-pressure pre-column concentration method for micro HPLC and its application to the rapid determination of dibutyl phthalate in water 321(1985)159

Concentration traps
Headspace gas analysis: the role and the design of concentration traps specifically suitable for capillary GC 321(1985)45

Connectivity indices
Calculation of retention indices by molecular topology: chlorinated alkanes 314(1984)1

Reversed-phase retention behaviour of Dns-amides as a function of eluent composition and molecular structure in ethanol-water and methanol-water 314(1984)141

Calculation of retention indices by molecular topology. Chlorinated benzenes 319(1985)1

Continuous flow carbon-13 nuclear magnetic resonance spectroscopy
Continuous-flow carbon-13 nuclear magnetic resonance spectroscopy 312(1984)91

Contraceptives
Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313

General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470

Coordination complexes
Isotachophoresis in inorganic chemistry (Review) 313(1984)189

Copolymers
Two-dimensional analysis of copolymers by size-exclusion chromatography and gradient-elution reversed-phase precipitation chromatography 317(1984)615

Copper
Enantiomeric resolution of an optically active guanine derivative by HPLC with phenylalanine-Cu(II) in the mobile phase 303(1984)217

Copper chelates
Volatility and GC of some derivatives of Schiff base chelates 318(1985)350

Copper phthalocyanine
Determination of polychlorinated biphenyls in copper phthalocyanine pigments 325(1985)456

Coprogen
Separation of ferrichromes and other hydroxamate siderophores of fungal origin by reversed-phase chromatography 301(1984)247

Cortexolone
Simultaneous analysis of cortexolone and cortisol by HPLC for use in the metyrapone test 306(1984)345

Corticosteroids
Analysis of flumethasone pivalate formulations by HPLC 301(1984)477

Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188

Application of a high-performance gel permeation LC procedure to the determination of binding of prednisolone to high-affinity binding sites in human serum 305(1984)194

Simultaneous analysis of cortexolone and cortisol by HPLC for use in the metyrapone test 306(1984)345

Application of reversed-phase HPLC for radioimmunoassay of plasma 18-hydroxycorticosterone 308(1984)295

Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Qualitative detection of corticosteroids in equine biological fluids and the comparison of relative dexamethasone metabolite/dexamethasone concentration in equine urine by micro-LC-MS 315(1984)359

Determination of trace levels of steroids in blood plasma by LC with peroxyoxalate chemiluminescence detection 317(1984)355

Clinical application of the rapid HPLC determination of serum cortisol 317(1984)377

Corticosterone
Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Corticotropin
HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99

Cortisol
Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

Analysis of cortisol, methylprednisolone, and methylprednisolone hemisuccinate. Absence of effects of troleandomycin on ester hydrolysis 305(1984)271

Simultaneous analysis of cortexolone and cortisol by HPLC for use in the metyrapone test 306(1984)345

Clinical application of the rapid HPLC determination of serum cortisol 317(1984)377

Cortisone
Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Cotinine
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291

Coumaphos
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Coumarins
Separations of some coumarins of higher plants by LC 314(1984)323

HPLC analysis of coumarins and flavonoids from section Tridentatae of *Artemisia* 322(1985)236

Coumatetralyl
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Counter-current chromatography
Purification of a cholecystokinin fragment peptide in the horizontal flow-through coil planet centrifuge 301(1984)277

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. I. Hydrodynamic distribution of two solvent phases in a helical column subjected to two types of synchronous planetary motion 301(1984)377

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. II. Phase distribution diagrams for helical and spiral columns 301(1984)387

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

Continuous extraction of urinary anthracycline antitumor antibiotics with the horizontal flow-through coil planet centrifuge 307(1984)323

Horizontal flow-through coil planet centrifuge: some practical applications of countercurrent chromatography 315(1984)287

Fractionation of condensed tannins by counter-current chromatography 318(1985)85

Coupled columns
Theory of the efficiency of serially connected GC columns and the effect on optimized speed of analysis 325(1985)1

CR
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b*,*f*]-1:4-oxazepine 315(1984)417

Creatine
HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281

Determination of creatine in body fluids and muscle 305(1984)450

Creatine phosphate
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 305(1984)183

Creatinine
HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281

Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Determination of creatinine in human, dog and rat urine by HPLC on a column of hydroxymethylated porous polystyrene 307(1984)399

Cromoglycate
Determination of sodium cromoglycate in human urine by HPLC on an anion-exchange column 305(1984)228

Crotoxyphos
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Crowded aliphatic esters
GC of homologous esters. XXX. Unexpected predominancy of polar effects over steric effects in crowded aliphatic esters: a GLC appraisal of specific interactions 324(1985)355

Crude oils
Fingerprinting of crude oil spills 322(1985)107

Reversed-phase separation of alkanes in high-boiling fractions of crude oil 322(1985)347

Analysis of Australian crude oils by high-resolution capillary GC-MS 323(1985)305

Crushing device
GC analysis of volatiles in fluid and gas inclusions 302(1984)181

CS
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b*,*f*]-1:4-oxazepine 315(1984)417

Curdlan
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Cyanide
Determination of cyanide and thiocyanate in blood plasma and red cells by HPLC with fluorometric detection 308(1984)133

4-Cyano4'-*n*-alkoxyformyl-oxyazobenzene
Phase transitions of 4-cyano-4'-*n*-alkoxyformyloxyazobenzenes on silanized supports 324(1985)309

Cyanogenic glycosides
Electrochemical detection of cyanogenic glycosides after enzymatic post-column cleavage 303(1984)67

Cyanogenic glycosides and cyanohydrins in plant tissues. Qualitative and quantitative

determination by enzymatic post-column cleavage and electrochemical detection, after separation by HPLC 303(1984)77

4-Cyano-4'-*n*-heptyloxyformyl-oxyazobenzene
Distribution of the liquid crystal stationary phase on the surface of diatomite supports 324(1985)299

Cyanohydrins
Cyanogenic glycosides and cyanohydrins in plant tissues. Qualitative and quantitative determination by enzymatic post-column cleavage and electrochemical detection, after separation by HPLC 303(1984)77

Cyanopropyl-silica
Basis of retention in normal-phase HPLC with cyano-propyl columns 303(1984)1

Cyanosilicones
Cyanosilicones as stationary phases in GC. III. Synthesis, characterization and evaluation 302(1984)319

Cyclazocine
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Cyclodextrins
Unmodified and acylated cyclodextrin stationary phases for LC separation of aromatic compounds 301(1984)345

Preparation and retention behaviour of chemically bonded methylated-cyclodextrin stationary phases for LC 314(1984)193

Cyclohexenes
Structure assignment by retention index in gas-liquid radiochromatography of substituted cyclohexenes 312(1984)75

Cyclophosphamide
Plasma cyclophosphamide assay by selective ion monitoring 310(1984)407

Cyclopropanes
Reaction GC-MS. VI. Use of deuterium as the carrier and reagent gas in reaction GC-MS 301(1984)77

Cyclosophoraose
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Cyclosporine
Semi-automated HPLC method for the determination of cyclosporine in plasma and blood using column switching 305(1984)353

Simple and rapid HPLC analysis of cyclosporine in human blood and serum 309(1984)441

Cystathionine
Quantitative analysis of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using isotachophoresis 311(1984)183

Cysteine
Analysis of hepatic reduced glutathione, cysteine and homocysteine by cation-exchange HPLC with electrochemical detection 308(1984)83

5-S-Cysteinyl-DOPA
Determination of DOPA, dopamine, and 5-S-cysteinyl-DOPA in plasma, urine, and tissue samples by HPLC with electrochemical detection 311(1984)154

Cysteinylglycine
Cysteinylglycine in urine determined by HPLC 308(1984)75

Cytochrome *c*
Protein separations on reversed-phase HPLC minicolumns 317(1984)119

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Cytochrome P-448
High resolution GC analysis of cytochrome P-448 inducing polychlorinated biphenyl congeners in hazardous waste 314(1984)482

Cytokinins
Assay of enzymes involved in cytokinin metabolism by means of reversed-phase HPLC 317(1984)263

Trace enrichment of cytokinins using neutral polystyrene resins 321(1985)175

Cytosine
Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201

GC-MS characteristics of methylated perfluoroacyl derivatives of cytosine and 5-methylcytosine 314(1984)211

Evidence for surface effects in an electron-capture detector 314(1984)219

D

2,4-D
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Dactylariol
Analysis of naturally occurring reduced anthraquinones obtained from *Alternaria porri* (Ellis) Ciferri by HPLC 318(1985)139

Dapsone
Quantification of antimalarial drugs. II. Simultaneous measurement of dapsone, monoacetyldapsone and pyrimethamine in human plasma 307(1984)426

Daunorubicin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

Continuous extraction of urinary anthracycline antitumor antibiotics with the horizontal flow-through coil planet centrifuge 307(1984)323

D&C Red No. 33
Purification of D&C Red No. 33 by preparative HPLC 314(1984)379

Determination of 1,3-diphenyltriazene and azobenzene in D&C Red No. 33 by solvent extraction and reversed-phase HPLC 321(1985)185

***o,p'*-DDD**
Mitotane (*o,p'*-DDD) emulsion and tablet analysis by HPLC 319(1985)467

3-Deazaguanine
Determination of 3-deazaguanine in mice plasma by HPLC 309(1984)403

***n*-Decane**
Indirect determination of specific retention volumes in GC. Use of retentions of the McReynolds probes relative to *n*-decane 312(1984)69

Deconvolution methods
Analytical solution for the area under chromatographic peaks obtained from the deconvolution method of Vaidya and Hester 303(1984)177

Dehydroepiandrosterone
Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Dehydroretinals
HPLC of retinals, retinols (vitamin A_1) and their dehydro homologues (vitamin A_2): improvements in resolution and spectroscopic characterization of the stereoisomers 301(1984)65

3-Dehydroretinol
Quantitative HPLC method for determining the isomer distribution of retinols (vitamin A_1) and 3-dehydroretinol (vitamin A_2) in fish oils 312(1984)423

Dehydroretinols
HPLC of retinals, retinols (vitamin A_1) and their dehydro homologues (vitamin A_2): improvements in resolution and spectroscopic characterization of the stereoisomers 301(1984)65

Delsoline
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Demecarium bromide
Determination of demecarium bromide and related compounds by HPLC 321(1985)401

4-Demethoxy-adriamycin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

4-Demethoxy-daunorubicin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

Denaturation
Effects of urea-thermal denaturation on the high-performance cation-exchange chromatography of α-chymotrypsinogen-A 314(1984)337

Effects of HPLC solvents and hydrophobic matrices on the secondary and quaternary matrices on the secondary and quaternary structure of a model protein. Reversed-phase and size-exclusion HPLC 317(1984)129

Kinetics of unfolding of proteins on hydrophobic surfaces in reversed-phase LC 317(1984)227

Reversed-phase chromatography of proteins with diphenyl-silica column and hydro-organic eluents containing two organic solvents 318(1985)11

Density
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

4'-Deoxy-adriamycin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

Deoxynivalenol
Confirmation of nivalenol and deoxynivalenol by on-line LC-MS and GC-MS. Comparison of methods 318(1985)103

Deoxynucleosides
Mobile phase optimization for multi-functional solutes in LSC. Resolution of synthetic

intermediates of oligodeoxyribonucleotides 323(1985)227

2'-Deoxynucleoside 5'-triphosphates
HPLC purification of deoxynucleoside 5'-triphosphates and their use in a sensitive electrophoretic assay of misincorporation during DNA synthesis 317(1984)283

3-Deoxy-D-*manno*-oct-2-ulosonic acid
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Reversed-phase HPLC of hydrazones of 3-deoxy-D-*manno*-oct-2-ulosonic acid and neuraminic acids 315(1984)451

Derivatization, GC
Computerised GC-MS analysis of complex mixtures of alkyl porphyrins 301(1984)107

Phosgene: a versatile reagent for enantiomer separation by capillary GC 301(1984)129

Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173

Use of steroid profiling in the resolution of pregnenolone metabolites from porcine testicular preparations 301(1984)189

GLC separation of hydroxy monocarboxylic acids and dicarboxylic acids on a fused-silica capillary column 301(1984)273

Selective LC isolation and GC-MS analysis of ketonic bile acids in faeces 302(1984)1

Trace analysis of amines and isocyanates using glass capillary GC and selective detection. III. Determination of aliphatic and alicyclic amines as perfluoro fatty acid amides using electron-capture and nitrogen-selective detection 303(1984)89

GC-MS of polyamines as their N-ethyloxycarbonyl derivatives and identification of *sym*-homospermidine and *sym*-norspermine in mosses and ferns 303(1984)99

Determination of lysinoalanine as the heptafluorobutyryl isobutyl ester derivative by GLC 303(1984)234

Sensitive assay for amino sugars using capillary GC with nitrogen-selective detection 303(1984)380

Preparation of pentafluorobenzyl esters of arachidonic acid lipoxygenase metabolites. Analysis by GC and negative-ion chemical ionization MS 305(1984)3

Possibilities and limitations in the analysis of amino acid oxazolidinones in the femtomole range by GC with electron-capture detection 305(1984)442

Lipoxygenase products. A novel GC-MS assay for monohydroxy fatty acids 306(1984)1

Quantitative GLC of taurine 306(1984)61

Revised method for the quantitative determination of 5-hydroxytryptamine in human plasma by GC-MS-selected ion monitoring 307(1984)1

Determination of urochloralic acid, the glucuronic acid conjugate of trichloroethanol, by GC with electron-capture detection and its application to urine, plasma and liver 307(1984)111

GC determination of gemfibrozil and its metabolites in plasma and urine 307(1984)210

Quantitative analysis of prostaglandins in cell culture medium by high-resolution GC with electron-capture detection 307(1984)251

Simultaneous quantitation of disopyramide and its mono-dealkylated metabolite in human plasma by fused-silica capillary GC using nitrogen-phosphorus specific detection 307(1984)305

Determination of germicidal phenols in blood by capillary column GC 307(1984)420

Monitoring of *S*- and *R*-tocainide in human plasma after heptafluorobutyrylation, separation on Chirasil-Val and electron-capture detection 308(1984)181

Sensitive GLC-electron-capture detection method for determination of sobrerol in biological fluids 308(1984)189

Sensitive method for the determination of diclofenac in human plasma by GC-MS 308(1984)329

Determination of morphine in biological samples by GC-MS. Evidence for persistent tissue binding in rats twenty-two days post-withdrawal 309(1984)73

GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

Hydroxycarboxylic and oxocarboxylic acids in urine: products from branched-chain amino acid degradation and from ketogenesis 309(1984)225

Characterization of artefacts produced by treatment of organic acids with diazomethane 309(1984)243

Determination of N-acetyl-L-cysteine in biological fluids 309(1984)261

Sensitive detection of amino acids in human serum and dried blood disc of 3 mm diameter for diagnosis of inborn errors of metabolism 309(1984)279

Simultaneous determination of metoprolol and metabolites in urine by capillary column GC as oxazolidineone and trimethylsilyl derivatives 309(1984)317

Branched-chain α-keto acid analysis in biological fluids: preparative clean-up by anion-exchange and analysis by capillary GC 310(1984)1

GLC evaluation of lofemizole in biological samples for pharmacokinetic investigations. Comparison of two analytical methods 310(1984)51

Determination of plasma phenytoin by capillary GC with nitrogen-phosphorus detection and with selective ion monitoring 310(1984)199

GC analysis of amino acids in physiological fluids: a critique (Review) 310(1984)223

GC-MS profiling with negative-ion chemical ionization detection of prostaglandins and their 15-keto and 15-keto-13,14-dihydro catabolites in rat blood 310(1984)233

Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251

GC quantitation of methoxyphenamine and three of its metabolites in plasma 310(1984)307

Improved GLC method for the determination of baclofen in plasma and urine 310(1984)319

Plasma cyclophosphamide assay by selective ion monitoring 310(1984)407

Silica gel HPLC for the simultaneous determination of propanolol and 4-hydroxypropranolol enantiomers after chiral derivatization 310(1984)424

GC analysis of endogenous catecholamines, phenolic amines and derived isoquinolines using short glass capillary columns and electron-capture detection 311(1984)17

Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141

GLC analysis with electron-capture detection of diclofensine in human plasma following derivatization 311(1984)189

Spectroscopic properties and GC behaviour of aryl β-diketones and selected chelates 312(1984)109

GC and GC-MS analysis of diastereomeric acyclic isoprenoid esters using fused-silica capillary columns 312(1984)395

Inorganic GC (Review) 313(1984)3

Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201

GC-MS characteristics of methylated perfluoroacyl derivatives of cytosine and 5-methylcytosine 314(1984)211

Derivatization at the liquid-solid interface. Application to the trace analysis of organic acids in aqueous samples 314(1984)421

Separation and analysis of ethylenediamines as their *tert.* -butyldimethylsilyl derivatives by GLC and MS 314(1984)455

Studies of cyclic di-*tert.* -butylsilylene derivatives of diols and hydroxy acids by GC-MS 315(1984)119

Determination of aldehydes in air as dimethone derivatives by GC with electron-capture detection 315(1984)412

Simple method for the separation of diastereomers of 4-fluoroglutamic acid by GLC 315(1984)421

Resolution of α-halogenocarboxylic acids by GC on (*R*)-*N*-lauroyl-α-(1-naphthyl)ethylamine 318(1985)235

Determination of methamphetamine and its metabolites in rat tissues by GC with a nitrogen-phosphorus detector 318(1985)307

Separation of sequence isomeric dipeptides by high-resolution GC 318(1985)384

Separation of acebutolol and diacetolol diastereomers by reversed-phase HPLC 318(1985)393

Chromatographic analysis of polyether based polyurethanes after alkali fusion 319(1985)121

Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Determination of carbonyl compounds in an Australian (Rundle) shale oil 319(1985)230

Optimisation of conditions for the trimethylsilylation of trichothecene mycotoxins 319(1985)376

Alkylation with dialkylsulfate and diisopropylethylamine 319(1985)382

Analysis of γ-carboxyglutamic acid by GLC 319(1985)404

N-Permethylation of polyamines at trace levels for GC analysis 322(1985)169

GLC assay for asparagine and glutamine 322(1985)228

Improved elution of metal β-diketonates on fused-silica wall-coated open-tubular columns 324(1985)75

Less hazardous derivatization procedure for GC of plant hormone abscisic acid 325(1985)425

GC resolution of substituted glutamic acid enantiomers 325(1985)444

Derivatization, GC-MS
Use of impregnated reagents in the pentafluorobenzylation of indoleamine metabolites 325(1985)309

Derivatization, LC
Sequence analysis of derivatized peptides by HPLC-MS 301(1984)425

Separation of isocyanate prepolymer components as their urea derivatives by reversed-phase HPLC 301(1984)485

Selective LC isolation and GC-MS analysis of ketonic bile acids in faeces 302(1984)1

Sub-picomol chemiluminescence detection of Dns-amino acids separated by HPLC with gradient elution 303(1984)173

HPLC with automated pre-column derivatization for amino acids 303(1984)238

Use of immobilized amino acid oxidase as post-column reactor in the HPLC of amino acids 303(1984)396

Separation of (*R*)- and (*S*)-proxyphylline as diastereoisomeric camphanates by reversed-phase LC 303(1984)433

Fluorometric determination of total vitamin C in whole blood by HPLC with pre-column derivatization 305(1984)53

Determination of (endogenous) vitamin K_1 in human plasma by reversed-phase HPLC using fluorometric detection after post-column electrochemical reduction. Comparison with UV, single and dual electrochemical detection 305(1984)61

New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188

Fluorometric HPLC of 9-aminophenanthrene-derivatized free fatty acids 305(1984)261

HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281

Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Determination of bestatin in serum by HPLC with fluorescence detection 305(1984)381

Determination of free trimethyllysine in plasma and tissue specimens by HPLC 306(1984)79

Detection and quantitation of fluorescamine-labeled bradykinin, its analogues and metabolites using HPLC 306(1984)109

Simultaneous determination of the enantiomers of tocainide in blood plasma using GLC with electron-capture detection 306(1984)155

Direct enantiomeric determination of (-)- and (+)-propranolol in human serum by HPLC on a chiral stationary phase 306(1984)405

Studies on steroids. CC. Determination of 17-ketosteroid sulphates in serum by HPLC with electrochemical detection using pre-column derivatization 307(1984)23

Determination of isoniazid and acetylisoniazid in plasma or serum by HPLC (in French) 307(1984)137

Activated α-alkyl-α-arylacetic acid enantiomers for stereoselective TLC and HPLC determination of chiral amines 307(1984)145

Analysis of carbohydrates in lens, erythrocytes, and plasma by HPLC of nitrobenzoate derivatives 307(1984)180

Resolution of the enantiomers of ephedrine, norephedrine and pseudoephedrine by HPLC 307(1984)220

LC determination of amikacin in serum with spectrophotometric detection 307(1984)224

Determination of thiamine and thiamine phosphates in excitable tissues as thiochrome derivatives by reversed-phase HPLC on octadecyl silica 307(1984)283

HPLC resolution of enantiomers of 1-phenyl-2-aminopropanes (amphetamines) with four chiral reagents 307(1984)335

LC assay of heptaminol in serum and its oral pharmacokinetics in the dog 307(1984)351

Separation and determination of diacylglycerols as their naphthylurethanes by HPLC 307(1984)387

Automated anion-exchange chromatographic method for the quantitation of lactulose in physiological fluids and processed milks 307(1984)393

Cysteinylglycine in urine determined by HPLC 308(1984)75

Determination of cyanide and thiocyanate in blood plasma and red cells by HPLC with fluorometric detection 308(1984)133

Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 309(1984)43

Sensitive fluorimetry of adenine-containing compounds with HPLC 309(1984)63

HPLC assay of pirlimycin in human serum and urine using 9-fluorenylmethylchloroformate 309(1984)95

Measurement of maprotiline and oxaprotiline in plasma by HPLC of fluorescent derivatives 309(1984)107

Determination of γ-aminobutyric acid in physiological samples by a simple, rapid HPLC method 309(1984)139

Microdetermination of two antifibrinolytic agents (ε-aminocaproic acid and tranexamic acid) by LC and fluorimetric detection (in French) 309(1984)183

Determination of nicotinic acid in serum by HPLC with fluorescence detection 309(1984)309

Digoxin and metabolites in urine: a derivatization-HPLC method capable of quantitating individual epimers of dihydrodigoxin 310(1984)85

α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167

Histamine assay in tears by fluorescamine derivatization and HPLC 310(1984)390

Improved procedure for the HPLC determination of valproic acid in serum as its phenacyl ester 310(1984)450

Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59

Determination of non-protein-bound N-acetylcysteine in plasma by HPLC 311(1984)170

Determination of sodium 2-mercaptoethanesulphonate by HPLC using post-column reaction colorimetry or electrochemical detection 311(1984)234

HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257

Rapid and sensitive amino acid analysis of human collagens using HPLC 311(1984)375

Investigation of isoprenoid benzoates and naphthoates by reversed-phase LC. Isocratic elution characteristics of benzoates and naphthoates of C_5-C_{20} terpenoid alcohols 312(1984)121

Trace determination of aliphatic amines using HPLC with chemiluminescence excitation and photon counting 312(1984)203

Determination of fatty acids by HPLC of Dns-ethanolamine derivatives 312(1984)461

Determination of aliphatic amines in air by reversed-phase HPLC using 1-naphthyl isocyanate derivatives 312(1984)482

Reversed-phase retention behaviour of Dns-amides as a function of eluent composition and molecular structure in ethanol-water and methanol-water 314(1984)141

R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

HPLC resolution of enantiomers of γ-vinyl-γ-aminobutyric acid 314(1984)495

Resolution of amino acid enantiomers by HPLC using automated pre-column derivatisation with a chiral reagent 315(1984)279

Application of reversed-phase HPLC using pre-column derivatization with *o*-phthaldialdehyde for the quantitative analysis of amino acids in adult and fetal sheep plasma, animal feeds and tissues 315(1984)321

Examination of theoretical principles of gradient elution as applied to reversed-phase HPLC separation of phenylthiohydantoin amino acids 316(1984)359

Microbore HPLC and chemiluminescence detection of Dns-amino acids 316(1984)501

Optical resolution of amino acid enantiomers by HPLC 316(1984)547

LC separation of enantiomeric alkanolamines via diastereomeric tartaric acid monoesters 316(1984)605

Determination of trace levels of steroids in blood plasma by LC with peroxyoxalate chemiluminescence detection 317(1984)355

HPLC fluorometric determination of cinnamaldehyde in perfume, cologne and toilet water 317(1984)421

Laser fluorimetry for capillary column LC: high-sensitivity detection of derivatized biological compounds 317(1984)443

Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23

Influence of trifluoroacetic anhydride concentration during derivatization and chromatography of N(O,S)-trifluoroacetyl amino acid *n*-propyl esters 318(1985)65

Enantiomeric separation of underivatized α-methyl-α-amino acids by HPLC 318(1985)117

Rapid and sensitive method for the determination of proline by reversed-phase HPLC with automated pre-column fluorescence derivatization 318(1985)132

Post-column reaction of amino acids with the pentane-2,4-dione-formaldehyde system for their automated analysis 318(1985)367

HPLC of carbonyl compounds as 2,4-dinitrophenylhydrazones with electrochemical detection 318(1985)427

Formation and separation of fluorescent derivatives of phosphatidic acid 320(1985)421

HPLC of short- and long-chain fatty acids as 2-nitrophenylhydrazides 321(1985)165

Pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene 322(1985)321

Use of benzoin as pre-column fluorescence derivatization reagent for the HPLC of angiotensins 322(1985)473

Complete automatization of peptide maps by reversed-phase LC using *o*-phthalaldehyde pre-column derivatization 323(1985)373

HPLC determination of *d*-/*l*-epinephrine enantiomer ratio in lidocaine-epinephrine local anesthetics 325(1985)249

Determination of urea and ammonia using ion-pair LC with on-line post-column derivatization in an enzymatic solid-phase reactor 325(1985)255

New high-performance LC system for the separation of biogenic amines as their Dns derivatives 325(1985)462

Derivatization, TLC
α-Alkyl-α-arylacetic acid derivatives as fluorescence markers for TLC and HPLC assay of amines and alcohols 310(1984)167

TLC of hydroxylysine for collagen analysis 311(1984)400

Fluorometric evaluation of sarcosine in urine and serum 311(1984)405

Desipramine
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

N-Desmethylmethsuximide
Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Desmethylmisonidazole
Determination of misonidazole and desmethylmisonidazole in plasma by HPLC with reductive electrochemical detection 310(1984)159

Desmosine
Separation of the cross-linking amino acids of elastin on thin-layer plates 305(1984)461

Desorption isotachophoresis
Desorption isotachophoresis: quantitative characterization of sorption and desorption conditions 320(1985)75

Detection, LC
Indirect UV detection of non-absorbing solutes in reversed-phase HPLC with the help of UV-active, non-ionic mobile phase constituents 312(1984)345

Detector response
HPLC analysis of basic drugs on silica columns using non-aqueous ionic eluents. I. Factors influencing retention, peak shape and detector response 323(1985)173

HPLC analysis of basic drugs on silica columns using non-aqueus ionic eluents. II. Application of UV, fluorescence and electrochemical oxidation detection 323(1985)191

Detectors, electrophoresis
On-column UV absorption detector for open tubular capillary zone electrophoresis 315(1984)135

Detectors, GC
Bipolar pulsed microvolume electron-capture detector for capillary GC 302(1984)221

New design for helium ionization detection 302(1984)257

Comparison of a new electrochemical detector for GC analysis with the electrolytic conductivity detector 302(1984)269

Atmospheric-pressure microwave-induced helium plasma spectroscopy for simultaneous multielement GC detection 302(1984)277

Ionization quenching of a hydrogen atmosphere flame as a detection method for GC 303(1984)331

Detection of GC eluates by simultaneous absorbance and fluorescence measurements 303(1984)341

Use of electron-capture data at short pulse intervals for study of electron-capture mechanisms 312(1984)31

Effect of trialkylamines on the response of the electron-capture detector to acridine 312(1984)49

Inorganic GC (Review) 313(1984)3

The mass spectrometer as a chlorine-selective chromatographic detector. I. Description and evaluation of the technique 314(1984)83

The mass spectrometer as a chlorine-selective chromatographic detector. II. Applications to chemical systems 314(1984)93

Evidence for surface effects in an electron-capture detector 314(1984)219

Determinations of chlorinated hydrocarbons by GC using response factor calibration 314(1984)243

Design and performance of a mas-flow-modulated detector for GC 319(1985)263

Detectors, isotachophoresis
Aspects of detection and identification in isotachophoresis 320(1985)67

UV detection at 206 nm in isotachophoresis 320(1985)99

Spectrophotometric detection in capillary isotachophoresis: determination of the composition of mixed zones 320(1985)185

Improvement of the performance of a high-frequency contactless conductivity detector for isotachophoresis 320(1985)233

The performance of the UVM-4 UV analyser in isotachophoresis 320(1985)241

Simple UV absorption detector for capillary isotachophoresis 325(1985)137

Detectors, LC
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Trace analysis of organic compounds by HPLC with photoionization detection 302(1984)43

Use of a rotating disc multiwavelength detector operating in the visible region of the spectrum for monitoring ball pen inks separated by HPLC 303(1984)211

Comparison of detection systems for the HPLC analysis of complex triglyceride mixtures 303(1984)386

New thermionic detector utilizing orthogonal nebulization for capillary column LC 314(1984)103

HPLC of organic acids with potentiometric detection using a metallic copper electrode 315(1984)261

UV visualization of inorganic anions by reversed-phase ion-interaction chromatography: factors that control sensitivity and detection 316(1984)211

Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Multi-channel detectors for micro HPLC: examination of flow cell structures 316(1984)441

Modifications of the wall-jet electrochemical detector for LC and flow analysis 316(1984)451

Analysis of steroids in bulk pharmaceuticals by LC with light-scattering detection 316(1984)461

Application of photodiode array UV detector to unresolved peak analysis 316(1984)473

Screening for new microbial products by HPLC using a photodiode array detector 316(1984)487

Limitation of assessing HPLC peak purity with photodiode array detectors 319(1985)387

Electrochemiluminescence as a detection technique for reversed-phase HPLC. I. Preliminary experiments 321(1985)227

Ion chromatography of inorganic anions with potentiometric detection using a metallic copper electrode 321(1985)363

Ten years of LC-MS 323(1985)3

Chromatographic perspective of HPLC-MS 323(1985)13

Determination of dibenzothiophene in oils by LC-tandem MS 323(1985)33

Open-tubular LC-MS using direct liquid introduction 323(1985)37

Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

HPLC-MS of derivatized and underivatized amino acids 323(1985)67

Performance of a moving belt LC-MS interface 323(1985)81

Micro column LC-MS using a capillary interface 323(1985)89

Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

On-line trace enrichment for improved sensitivity in LC with direct liquid introduction MS detection 323(1985)113

Design and construction of an interface for direct liquid introduction coupling of a microbore high-performance liquid chromatograph to a quadrupole mass spectrometer 323(1985)121

Studies with a laboratory-constructed thermospray liquid chromatograph-mass spectrometer interface 323(1985)127

On-line LC-MS analysis of non-ionic surfactants 323(1985)135

Qualitative and quantitative analysis of ranitidine and its metabolites by HPLC-MS 323(1985)143

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Applications of thermospray ionisation LC-MS to compounds of pharmaceutical interest 323(1985)163

HPLC analysis of basic drugs on silica columns using non-aqueus ionic eluents. II. Application of UV, fluorescence and electrochemical oxidation detection 323(1985)191

Examination of the concentration response of evaporative light-scattering mass detectors 323(1985)297

New, modified flow-through polarographic detector 323(1985)406

Evaluation of the mass detector for quantitative detection of triglycerides and fatty acid methyl esters 324(1985)35

Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Construction of a simple wall-jet electrochemical flow-cell for HPLC 325(1985)294

Dethiobiotin
Use of LC-MS for the quantitation of dethiobiotin and biotin in biological samples 303(1984)272

Deuterated solvents
HPLC of phospholipids using deuterated solvents for IR detection 307(1984)261

Deuterium
Reaction GC-MS. VI. Use of deuterium as the carrier and reagent gas in reaction GC-MS 301(1984)77

Dexamethasone
Qualitative detection of corticosteroids in equine biological fluids and the comparison of relative dexamethasone metabolite/dexamethasone concentration in equine urine by micro-LC-MS 315(1984)359

Dextran
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Dextran sulphate
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Dextromethorphan
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Diacetolol
Separation of acebutolol and diacetolol diastereomers by reversed-phase HPLC 318(1985)393

Diacylglycerols
Separation and determination of diacylglycerols as their naphthylurethanes by HPLC 307(1984)387

***gem*-Dialkyl effect**
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

Dialkylphosphates
Post-column reaction detection system for the determination of organophosphorus compounds by LC 324(1985)53

Dialkyl phthalates
Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Supercritical fluid chromatography of various samples with different polarities 315(1984)39

Dialkylsulphate
Alkylation with dialkylsulfate and diisopropylethylamine 319(1985)382

Dialkyltin
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341

Speciation of alkyltin compounds by HPLC with a cyanopropyl-bonded silica column 315(1984)351

Diamine oxidase
Sensitive assay for diamine oxidase activity using HPLC 309(1984)151

Diamines
Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59

Chromatographic analysis of polyether based polyurethanes after alkali fusion 319(1985)121

3,4-Diaminopyridine
HPLC determination of 3,4-diaminopyridine in human plasma 309(1984)222

Diamorphine
see Heroin

Diastereomer separation
Aryl substituted diastereomeric alkenes: GC behavior on a non-polar *versus* a liquid crystal phase 321(1985)127

Diazepam
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Diazomethane
Convenient apparatus for the generation of small amounts of diazomethane 303(1984)193

Characterization of artefacts produced by treatment of organic acids with diazomethane 309(1984)243

Dibekacin
Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Dibenzofurans
Characteristics of an extraction and purification procedure for chlorinated dibenzo-*p*-dioxins and dibenzofurans in soil and liver 312(1984)165

Dibenzothiophene
Determination of dibenzothiophene in oils by LC-tandem MS 323(1985)33

Dibenzothiophene-5-oxide
New injection syringe for GC 322(1985)55

Dibenz[*b,f*]-1:4-oxazepine
GC-MS identification of ω-chloroacetophenone, *o*-chlorobenzylidenemalononitrile and dibenz[*b,f*]-1:4-oxazepine 315(1984)417

3,3'-Dibromo-5,5'-dichloro-2,2'-dihydroxydiphenylmethane
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

2,6-Di-*tert.*-butyl-4-methylphenol
Determination of 2,6-di-*tert.*-butyl-4-methylphenol in aviation turbine fuel by LC with electrochemical detection 322(1985)376

Dibutyl phthalate
Development of an on-line high-pressure pre-column concentration method for micro HPLC and its application to the rapid determination of dibutyl phthalate in water 321(1985)159

Dibutyltin
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341

Dicaffeoylquinic acid
Chromatographic separation and identification of naturally occurring chlorogenic acids by ^1H nuclear magnetic resonance spectroscopy and MS 315(1984)253

Dicarboxylic acids
Detection of volatile fatty and dicarboxylic acids on alkaline pherograms with Mills' reagent 318(1985)446

Dicarboxylic porphyrins
HPLC of dicarboxylic porphyrins and metalloporphyrins: retention behaviour and biomedical applications 317(1984)333

Dichlorobenzenes
Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

2,5-Dichloro-p-benzoquinone
Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

***cis*-Dichlorodiammineplatinum(II)**
Quantitative analysis of *cis*-dichlorodiammineplatinum(II) by high-performance TLC 315(1984)481

Dichloroethyl alkanoates
GLC analyses. XXXVI. Separation of ethyl, 2-chloroethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns with temperature programming 322(1985)97

Dichloromethane
Mixed solvents in GLC. Activity coefficients for chloroform, dichloromethane and ethyl acetate in squalane-dinonyl phthalate mixtures at 303°K 324(1985)231

Dichlorophene
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

4,4'-Dichloro-3-trifluoromethylcarbanilide
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Dichlorprop
Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Dichotomous filters
Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53

Diclofenac
Sensitive method for the determination of diclofenac in human plasma by GC-MS 308(1984)329

Diclofensine
GLC analysis with electron-capture detection of diclofensine in human plasma following derivatization 311(1984)189

Dicyclomine
Determination of nanogram amounts of dicyclomine with GC and nitrogen-selective detection 308(1984)341

Dielectric constants
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433

Diesel exhaust particulate
Quantitative analysis of polycyclic aromatic compounds in diesel exhaust particulate

extracts by combined chromatographic techniques 312(1984)183

Determination of polycyclic aromatic hydrocarbons in diesel exhaust particulate matter and diesel fuel oil 312(1984)247

Diesel fuel
HPLC procedure for direct determination of the compound class composition of diesel and kerosine fuels using refractive index detection 312(1984)237

Determination of polycyclic aromatic hydrocarbons in diesel exhaust particulate matter and diesel fuel oil 312(1984)247

Supercritical fluid chromatography and supercritical fluid chromatography-MS of marine diesel fuel 314(1984)253

Diethylcarbamazine
Determination of diethylcarbamazine in blood using GC with alkali flame ionization detection 308(1984)334

Diethylene glycol
Identification of diethylene glycol in sera from egyptian children by frequency-pulsed electron-capture GLC 309(1984)269

Diethylenetriamine
Separation and analysis of ethylenediamines as their *tert.* -butyldimethylsilyl derivatives by GLC and MS 314(1984)455

Diethyl ether
Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

Di-(2-ethylhexyl)phthalate
HPLC procedure for the determination of di-(2-ethylhexyl)phthalate in human blood specimens. Problems of variable-extraction yield and the use of standard addition for calibration 311(1984)101

Difenacoum
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Diffusion
Diffusion in the bubble flow meter in inverse GC experiments 301(1984)261

Diflunisal
Direct analysis of diflunisal ester and ether glucuronides by HPLC 307(1984)432

Digoxin
Digoxin and metabolites in urine: a derivatization-HPLC method capable of quantitating individual epimers of dihydrodigoxin 310(1984)85

Dihydrocaffeic acid
Dihydrocaffeic acid: a common contaminant in the LC-electrochemical measurement of plasma catecholamines in man 311(1984)148

Dihydrocapsaicin
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288

Dihydrodigoxin
Digoxin and metabolites in urine: a derivatization-HPLC method capable of quantitating individual epimers of dihydrodigoxin 310(1984)85

1,3-Dihydro-4-methyl-5-[4-(methylthio)benzoyl-]-2H-imidazol-2-one
Simultaneous analysis of a new cardiotonic agent, MDL 17,043, and its major sulfoxide metabolite in plasma by HPLC 306(1984)249

Dihydromyricetin
Isolation of quercetin, myricetin, and their respective dihydro-compounds by Sephadex LH-20 chromatography 324(1985)495

Dihydroquercetin
Isolation of quercetin, myricetin, and their respective dihydro-compounds by Sephadex LH-20 chromatography 324(1985)495

Dihydroquinidine
Inter-laboratory study on the separation of cinchona alkaloids by HPLC 318(1985)362

Dihydroxybenzoic acid
Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133

Dihydroxybenzoic acids
Capillary GC of dihydroxybenzoic, -phenylacetic and -phenylpropionic acids 324(1985)141

Dihydroxyboryl-agarose
HPLC separations on dihydroxyboryl-agarose 316(1984)301

9-(3,4-Dihydroxybutyl)guanine
Enantiomeric resolution of an optically active guanine derivative by HPLC with phenylalanine-Cu(II) in the mobile phase 303(1984)217

3,4-Dihydroxyphenylacetic acid
Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

Rapid and simple method for the simultaneous determination of 3,4-

dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by HPLC with electrochemical detection 308(1984)55

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Selective retention of catecholamines and their derivatives in reversed-phase ion-pair partition chromatography 322(1985)43

Dihydroxyphenylacetic acids
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Capillary GC of dihydroxybenzoic, -phenylacetic and -phenylpropionic acids 324(1985)141

2-(3,4-Dihydroxyphenyl)naphtho[1,2-*d*]-thiazole
Assay for catechol-O-methyltransferase in erythrocytes using a new fluorogenic substrate, 2-(3,4-dihydroxyphenyl)naphtho[1,2-*d*]thiazole 308(1984)93

Dihydroxyphenylpropionic acids
Capillary GC of dihydroxybenzoic, -phenylacetic and -phenylpropionic acids 324(1985)141

***trans*-2,3-*cis*-3,4-Dihydroxy-L-proline**
Chromatographic separation of preparative quantities of the stereoisomers of *trans*-2,3-*cis*-3,4-dihydroxy-L-proline 315(1984)447

Diisocyanates
Improved chromatographic procedure for determination of 9-(N-methylaminomethyl)anthracene isocyanate derivatives by HPLC 301(1984)481

Diisopropylethylamine
Alkylation with dialkylsulfate and diisopropylethylamine 319(1985)382

2,5-Diketo-D-gluconic acid
Determination of D-gluconic, 5-keto-D-gluconic, 2-keto-D-gluconic and 2,5-diketo-D-gluconic acids by HPLC 312(1984)211

β-Diketonates
Volatility studies of metal chelates. VI. GC behaviour of aryl β-diketones and their metal chelates 322(1985)1

Volatility studies of metal chelates. VII. GC behaviour of chelates derived from the aromatic β-diketones 1-phenylbutane-1,3-dione and 4,4,4-trifluoro-1-(pentafluorophenyl)butane-1,3-dione 322(1985)23

Improved elution of metal β-diketonates on fused-silica wall-coated open-tubular columns 324(1985)75

Dilevalol
HPLC assay for labetalol in human plasma using a PRP-1 column and fluorometric detection 311(1984)319

Dimedone
Determination of aldehydes in air as dimethone derivatives by GC with electron-capture detection 315(1984)412

1-(2,5-Dimethoxy-4-methylphenyl)-2-aminopropane
R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Dimethylacetamidines
Retention indices of dimethylformamidines, dimethylacetamidines and tetramethylguanidines on a non-polar column 315(1984)95

Dimethylarginine
GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

7,12-Dimethylbenz[*a*]anthracene
Identification of phenols of 7,12-dimethylbenz[*a*]anthracene by conversion into their methyl ethers 319(1985)153

3,5-Dimethyl-4-chlorophenol
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Dimethyl ether
Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Dimethylformamidines
Retention indices of dimethylformamidines, dimethylacetamidines and tetramethylguanidines on a non-polar column 315(1984)95

Dimethyllysine
GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

Dimethylnicotinium
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291

5,5-Dimethyl-2,4-oxazolidinedione
Simultaneous determination of trimethadione and its metabolite in rat and human serum by HPLC 308(1984)393

Dimethyl polysulphides
Determination of trace levels of dimethyl polysulphides by capillary GC 319(1985)187

Dimethyl sulphoxide
Identification of trace organic compounds in dimethyl sulphoxide solution using HPLC and GC-MS 324(1985)373

N,N-Dimethyltryptamine
HPLC determination of hallucinogenic indoleamines with simultaneous UV photometric and voltammetric detection 320(1985)414

1,3-Dimethyluric acid
Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107

3-(2,4-Dinitroanilino)propyl-silica
Chromatographic selectivity in LC of polycondensed aromatic hydrocarbons on 3-(2,4-Dinitroanilino)propyl silica 324(1985)261

3,5-Dinitrobenzamides
Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Dinitrobenzene
Unmodified and acylated cyclodextrin stationary phases for LC separation of aromatic compounds 301(1984)345

N-(3,5-Dinitrobenzoyl)-α-amino acids
Reversed-phase chromatographic resolution of N-(3,5-dinitrobenzoyl)-α-amino acids on chiral stationary phases 322(1985)287

Dioctylsulphosuccinate
New ion-pair reagent for the HPLC separation of B-group vitamins in pharmaceuticals 301(1984)470

Diode-array detectors
Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Application of photodiode array UV detector to unresolved peak analysis 316(1984)473

Screening for new microbial products by HPLC using a photodiode array detector 316(1984)487

Limitation of assessing HPLC peak purity with photodiode array detectors 319(1985)387

Diols
Studies of cyclic di-*tert.*-butylsilylene derivatives of diols and hydroxy acids by GC-MS 315(1984)119

Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

Di(oxalato)diaquochromate(III)
HPLC in the study of some reactions of chromium(III) complexes 314(1984)387

Dioxins
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351

Characteristics of an extraction and purification procedure for chlorinated dibenzo-*p*-dioxins and dibenzofurans in soil and liver 312(1984)165

Dipeptides
Reversed-phase high-performance ion-pair chromatography of cobalt(III) coordination complexes (Review) 313(1984)93

Separation of sequence isomeric dipeptides by high-resolution GC 318(1985)384

Selection of optimum conditions for the preparative-scale separation of a dipeptide diastereomer mixture on the basis of measurements made in an analytical column 320(1985)397

Development of immobilized metal affinity chromatography. III. Interaction of oligopeptides with immobilized nickel iminodiacetate 323(1985)265

Prediction of retention data of protected dipeptides in normal phase HPLC 325(1985)282

Diphacinone
HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Diphenhydramine
Isothermal GC analysis of diphenhydramine after direct injection onto a fused-silica capillary column 307(1984)295

Diphenylphosphine-octadecylsilane
Synthesis, solid-state nuclear magnetic resonance characterization and chromatographic evaluation of a diphenylphosphine-octadecylsilane hybrid bonded phase 323(1985)281

1,3-Diphenyltriazene
Determination of 1,3-diphenyltriazene and azobenzene in D&C Red No. 33 by solvent extraction and reversed-phase HPLC 321(1985)185

Diphyllin
Solid-state fluorodensitometric quantitation of arylnaphthalene lignan lactones of *Cleistanthus collinus* 303(1984)296

Dipyrone
Simultaneous determination of dipyrone metabolites in plasma by HPLC 305(1984)477

Diquat
Anomalies in the HPLC determination of diquat in water samples 303(1984)206

Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Direct liquid introduction interface
Ten years of LC-MS 323(1985)3

Open-tubular LC-MS using direct liquid introduction 323(1985)37

Micro column LC-MS using a capillary interface 323(1985)89

Micro column LC-MS using a capillary interface 323(1985)89

Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Design and construction of an interface for direct liquid introduction coupling of a microbore high-performance liquid chromatograph to a quadrupole mass spectrometer 323(1985)121

Disaccharides
HPLC of disaccharides on amine-bonded silica columns 319(1985)51

Disc-like liquid crystals
Disc-like liquid crystalline stationary phases from the triphenylene derivatives group 315(1984)145

Disopyramide
Simultaneous quantitation of disopyramide and its mono-dealkylated metabolite in human plasma by fused-silica capillary GC using nitrogen-phosphorus specific detection 307(1984)305

Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Dispersion solubility parameter
Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1

Displacement effects
Rôle of displacement effects in trapping trace components from gases in sorbent-packed columns 324(1985)269

Displacement thin-layer chromatography
Conditions and parameters dominating displacement TLC 316(1984)95

Dissociation constants
Isotachophoretic determination of mobility and pK_a by means of computer simulation. III. Evaluation of mobility and pK_a of fifteen nucleotides and seven phosphorus oxoacids and their isotachophoretic separation 318(1985)195

Determination of dissociation constants of weak electrolytes by capillary isotachophoresis 320(1985)33

Determination of absolute mobilities and pK values by isotachophoresis 320(1985)147

Dissolved organic matter
Application of reversed-phase LC to dissolved organic matter in estuarine seawater 322(1985)358

Distribution coefficients
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47

Partitioning of oligoglucans in Sephadex G-15 in relation to their conformational structure 318(1985)57

Disulphonates
Analysis of α-olefinsulphonates by HPLC 325(1985)43

Diterpenoid alkaloids
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Dityrosine
Determination of dityrosine in protein fractions isolated from merino wool using reversed-phase HPLC 303(1984)429

Diuretics
HPLC behaviour of some pharmaceutically important thiazide, loop and potassium-sparing diuretics 325(1985)395

DNA
Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 308(1984)121

HPLC purification of deoxynucleoside 5'-triphosphates and their use in a sensitive electrophoretic assay of misincorporation during DNA synthesis 317(1984)283

Separation of large DNA restriction fragments by high-performance gel filtration on TSKgel DNA-PW 320(1985)440

DOPA
Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Determination of DOPA, dopamine, and 5-S-cysteinyl-DOPA in plasma, urine, and tissue samples by HPLC with electrochemical detection 311(1984)154

Determination of urinary and plasma dihydroxyphenylalanine by coupled-column HPLC with C_8 and C_{18} stationary phases 317(1984)27

In-situ preparation of a chemically bonded chiral stationary phase for the separation of aromatic α-amino acid enantiomers 325(1985)103

Dopamine
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Rapid and simple method for the simultaneous determination of 3,4-dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by HPLC with electrochemical detection 308(1984)55

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Dihydrocaffeic acid: a common contaminant in the LC-electrochemical measurement of plasma catecholamines in man 311(1984)148

Determination of DOPA, dopamine, and 5-S-cysteinyl-DOPA in plasma, urine, and tissue samples by HPLC with electrochemical detection 311(1984)154

Automated LC of urinary dopamine (in French) 311(1984)160

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

α-Doradexanthin
Direct diastereomeric resolution of carotenoids. I. 3-Hydroxy-4-oxo β-end group 318(1985)122

Double-bond determination
Separation and double-bond determination on nanogram quantities of aliphatic monounsaturated alcohols, aldehydes and carboxylic acid methyl esters 322(1985)484

Doxepin
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Doxorubicin
see Adriamycin

Doxycycline
Improvement of chemical analysis of antibiotics. VIII. Application of prepacked C_{18} Cartridge for the analysis of tetracycline residues in animal liver 325(1985)265

Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

Drugs
Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213

Dyes
Purification of D&C Red No. 33 by preparative HPLC 314(1984)379

Determination of synthetic organic colours in lipsticks by TLC and HPLC 318(1985)149

Chromatography of Astra Blue dyes with eluents containing potassium thiocyanate in organic solvents 319(1985)400

TLC of acid dyes on polyamide layers (in German) 322(1985)267

E

Ecdysteroids
HPLC determination of ecdysteroid titers in the house fly 317(1984)413

TLC of ecdysteroids 318(1985)373

Analytical droplet counter-current chromatography isolation of 20-hydroxyecdysone from *Vitex thyrsiflora* (Verbenaceae) 321(1985)246

Echinacoside
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Edgeworth-Cramér series
Characterization of extracolumn and concentration-dependent distortion of chromatographic peaks by Edgeworth-Cramér series 315(1984)67

Effluent fractionation
High-sensitivity radioassay in chromatographic effluents 302(1984)31

Eicosanoids
Picogram detection of eicosanoids by UV absorbance after narrow-bore HPLC. Comparison with conventional-bore column 308(1984)31

Eicosyl acetate
Microanalytical methods in the structure elucidation of sex-specific components in the

large pine weevil, *Hylobius abietis* L. (*Coleoptera, Curculionidae*) 323(1985)343

Elastin
Separation of the cross-linking amino acids of elastin on thin-layer plates 305(1984)461

Elastomers
TLC of age resisters for elastomers (phenols and organic phosphites) (in French) 314(1984)349

Chromatographic analysis of polyester-based polyurethane elastomers after alkali fusion 321(1985)295

Electric field gradients
Computer aided analysis of electric field gradients within isotachophoretic boundaries between weak electrolytes 320(1985)23

Electrochemical detector
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Comparison of a new electrochemical detector for GC analysis with the electrolytic conductivity detector 302(1984)269

HPLC of organic acids with potentiometric detection using a metallic copper electrode 315(1984)261

Modifications of the wall-jet electrochemical detector for LC and flow analysis 316(1984)451

HPLC analysis of basic drugs on silica columns using non-aqueus ionic eluents. II. Application of UV, fluorescence and electrochemical oxidation detection 323(1985)191

New, modified flow-through polarographic detector 323(1985)406

Construction of a simple wall-jet electrochemical flow-cell for HPLC 325(1985)294

Electrochemiluminescence
Electrochemiluminescence as a detection technique for reversed-phase HPLC. I. Preliminary experiments 321(1985)227

Electrodesorption
Micropreparative isotachophoretic electrodesorption of monoclonal antibodies from an affinity adsorbent 320(1985)81

Electrolyte cascading
Enhancing isotachophoresis sensitivity by low-concentration electrolyte cascading 320(1985)45

Electron-capture detector
Bipolar pulsed microvolume electron-capture detector for capillary GC 302(1984)221

Use of electron-capture data at short pulse intervals for study of electron-capture mechanisms 312(1984)31

Effect of trialkylamines on the response of the electron-capture detector to acridine 312(1984)49

Evidence for surface effects in an electron-capture detector 314(1984)219

Electron-capture mechanisms
Use of electron-capture data at short pulse intervals for study of electron-capture mechanisms 312(1984)31

Electrospray interface
Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Eleutherococcus senticosus
Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497

Eleutheroside E
Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497

Ellipticines
Use of electrochemical detection in the HPLC determination of hydroxylated ellipticine derivatives 309(1984)170

Eluent concentration
Computer-assisted optimization of eluent concentration and pH in ion chromatography 303(1984)321

Emepronium bromide
Determination of emepronium bromide in tablets by reversed-phase HPLC 322(1985)246

Emergence temperatures
Emergence temperatures as physical constants for measuring analyte retention in programmed temperature GC 312(1984)59

Enamino ketones
TLC of enamino ketones. II 312(1984)387

Enantiomer separation
Phosgene: a versatile reagent for enantiomer separation by capillary GC 301(1984)129

Quantitation of the enantiomers of β-hydroxyamino acids by GC resolution on an optically active packed column 301(1984)461

Enantiomeric resolution of an optically active guanine derivative by HPLC with phenylalanine-Cu(II) in the mobile phase 303(1984)217

Enantiomeric separation of Dns-amino acids by reversed-phase TLC 303(1984)251

Separation of (*R*)- and (*S*)-proxyphylline as diastereoisomeric camphanates by reversed-phase LC 303(1984)433

Simultaneous determination of the enantiomers of tocainide in blood plasma using GLC with electron-capture detection 306(1984)155

Separation, isolation and identification of optical isomers of 1,4-benzodiazepine glucuronides from biological fluids by reversed-phase HPLC 306(1984)231

Direct enantiomeric determination of (-)- and (+)-propranolol in human serum by HPLC on a chiral stationary phase 306(1984)405

Direct enantiomeric resolution of mephenytoin and its N-demethylated metabolite in plasma and blood using chiral capillary GC 307(1984)121

Activated α-alkyl-α-arylacetic acid enantiomers for stereoselective TLC and HPLC determination of chiral amines 307(1984)145

HPLC resolution of enantiomers of 1-phenyl-2-aminopropanes (amphetamines) with four chiral reagents 307(1984)335

Monitoring of *S*- and *R*-tocainide in human plasma after heptafluorobutyrylation, separation on Chirasil-Val and electron-capture detection 308(1984)181

Silica gel HPLC for the simultaneous determination of propanolol and 4-hydroxypropranolol enantiomers after chiral derivatization 310(1984)424

Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59

R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

HPLC resolution of enantiomers of γ-vinyl-γ-aminobutyric acid 314(1984)495

Resolution of amino acid enantiomers by HPLC using automated pre-column derivatisation with a chiral reagent 315(1984)279

Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Optical resolution of amino acid enantiomers by HPLC 316(1984)547

Chromatographic separation of enantiomers of acids with quinine as chiral counter ion 316(1984)553

Direct resolution of mono- and diol enantiomers of unsubstituted and methyl-substituted benz[*a*]anthracene and benzo[*a*]pyrene by HPLC with a chiral stationary phase 316(1984)569

Rational approach to the design of highly-effective chiral stationary phases 316(1984)585

LC separation of enantiomeric alkanolamines via diastereomeric tartaric acid monoesters 316(1984)605

Direct LC separation of enantiomers on immobilized protein stationary phases. IV. Molecular interaction forces and retention behaviour in chromatography on bovine serum albumin as a stationary phase 316(1984)617

Enantiomeric separation of underivatized α-methyl-α-amino acids by HPLC 318(1985)117

Resolution of α-halogenocarboxylic acids by GC on (*R*)-N-lauroyl-α-(1-naphthyl)ethylamine 318(1985)235

Separation of acebutolol and diacetolol diastereomers by reversed-phase HPLC 318(1985)393

Separation of diastereomeric amides by preparative HPLC and analysis of enantiomers on a chiral support (in French) 318(1985)398

Paper electrophoretic study of ion pair formation. XV. Resolution of optically active cobalt(III)-cyclohexanediamine complexes 318(1985)440

Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 319(1985)35

Clay CC for optical resolution. Resolution of aromatic compounds on a Λ-Ru(1,10-phenanthroline)$_3^{2+}$-montmorillonite column 319(1985)299

Preparative separation of enantiomers by flash chromatography 319(1985)392

HPLC analysis of epinephrine enantiomers using a UV detector in series with an optical activity detector 319(1985)419

GC separation of enantiomeric sulphur compounds on Chirasil-Val 320(1985)393

Reversed-phase chromatographic resolution of N-(3,5-dinitrobenzoyl)-α-amino acids on chiral stationary phases 322(1985)287

α-Arylalkylamine-derived chiral stationary phases. Evaluation of urea linkages 322(1985)295

Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444

In-situ preparation of a chemically bonded chiral stationary phase for the separation of aromatic α-amino acid enantiomers 325(1985)103

Determination of epimeric peptides for assessing enantiomeric purity of starting materials and studying racemization in peptide synthesis using HPLC 325(1985)111

HPLC determination of *d-/l-* epinephrine enantiomer ratio in lidocaine-epinephrine local anesthetics 325(1985)249

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

GC resolution of substituted glutamic acid enantiomers 325(1985)444

π-Energy effect
Dependence of the retention of phenols upon Van der Waals volume, π-energy and hydrogen-bonding effects 302(1984)89

Enkephalins
Separation of the enkephalins from proteins in an aqueous medium by chromatofocusing 314(1984)499

Separation of enkephalins from their metabolites and precursor by a single pump gradient HPLC method 325(1985)467

Enzymes
see also Proteins

Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

Rapid and sensitive assays using HPLC to measure the actitivies of phase II drug metabolising enzymes: glucuronyl transferase and sulfotransferase 301(1984)282

Ion-exchange chromatographic assay of peptidases acting on the C-terminal hexapeptide sequence of substance P 301(1984)415

Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Purification of chymotrypsin and trypsin by CC on agar gel particles 303(1984)283

Affinity chromatography of kid chymosin on histidyl-Sepharose 303(1984)285

Assay for catechol-O-methyltransferase in erythrocytes using a new fluorogenic substrate, 2-(3,4-dihydroxyphenyl)naphtho[1,2-*d*]thiazole 308(1984)93

Sensitive assay for diamine oxidase activity using HPLC 309(1984)151

Determination of monoamine oxidase B activity by HPLC 309(1984)156

Separation of haem compounds by reversed-phase ion-pair HPLC and its application in the assay of ferrochelatase activity 310(1984)31

Measurement of prolidase activity in erythrocytes using isotachophoresis 310(1984)401

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Assay for histidine decarboxylase in rat stomach and brain by HPLC with fluorescence detection 311(1984)390

Elution of the catalytic subunits of the type I and type II forms of cyclic AMP-dependent protein kinase within the type I chromatographic peak 312(1984)293

Product analysis of the action of *Trichoderma reesei* polygalacturonase by TLC 312(1984)492

Metal-ion mediated interaction of luciferase with tetraiodofluorescein 315(1984)175

HPLC of proteins. Purification of plant enzymes by ion-exchange chromatography 315(1984)333

Purification of placental alkaline phosphatase by sulphate-mediated chromatography on Cibacron Blue 3G-A 315(1984)425

Isolation of plant phosphoenolpyruvate carboxylase by high-performance size-exclusion chromatography 315(1984)430

Isolation of trypsin by column chromatography on tea particles 315(1984)478

Enzymatic assay procedures that employ HPLC: competition between phosphoribosyltransferases for a common substrate 316(1984)43

Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

Enzymatic coupled assay procedures that employ HPLC: the synthesis of orotidylate from ribose 317(1984)245

Assay of enzymes involved in cytokinin metabolism by means of reversed-phase HPLC 317(1984)263

Enzymes adsorbed on an ion exchanger as a post-column reactor: application to acetylcholine measurement 317(1984)343

Affinity chromatography of B_6-vitamin-dependent enzymes: purification of pig-heart branched-chain amino acid transaminase 318(1985)333

HPTLC assay for epoxide hydrolase activity and the determination of phenoxypropane-1,2-diols 318(1985)343

Comparison of high-performance ion-exchange chromatography and gel electrophoresis in protein separations 318(1985)358

Determination of folylpoly-γ-glutamate carboxypeptidase (folacin conjugase) activity using reversed-phase HPLC 318(1985)387

Sepharose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 319(1985)43

Rapid ion-exchange chromatography for preparative separation of proteins. Application to porcine and bovine carbonic anhydrases 319(1985)59

Sepharose derivatives containing citric acid as affinity ligand. Purification of fumarase 319(1985)173

Cibacron Blue 3G-A-substituted cross-linked agar beads: an inexpensive, fast-flow affinity medium for large scale applications 319(1985)218

Pyrimidine nucleoside phosphorylase assay by automated HPLC 319(1985)351

Fractionation of peptic glycopeptides from chicken ovalbumin by reversed-phase HPLC 320(1985)426

Determination of laccase activity with various aromatic substrates by HPLC 320(1985)435

Fast protein LC for the isolation of *Clostridium perfringens* type A α-toxin 321(1985)235

Headspace GC determination of β-galactosidase activity using electron-capture detection 321(1985)385

Simultaneous assay of choline kinase and choline oxidase in tissue by high-performance cation-exchange chromatography and continuous radioactive detection 324(1985)203

Rapid procedure for purification of β-amylase from *Ipomea batata* 324(1985)489

Rapid chromatographic purification of yeast alcohol dehydrogenase 325(1985)359

Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Ephedrine
Resolution of the enantiomers of ephedrine, norephedrine and pseudoephedrine by HPLC 307(1984)220

R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

4'-Epi-adriamycin
Rapid quantitative determination of seven anthracyclines and their hydroxy metabolites in body fluids 306(1984)323

4-Epianhydrotetracycline
Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303

Epidermal growth factor
Use of HPLC in the isolation and characterisation of mouse and rat epidermal growth factors and examination of apparent heterogeneity 305(1984)295

4,6-Epidoxycycline
Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

4-Epidoxycycline
Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

6-Epidoxycycline
Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

Epinephrine
see Adrenaline

4-Epitetracycline
Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303

Epoxide hydrolase
HPTLC assay for epoxide hydrolase activity and the determination of phenoxypropane-1,2-diols 318(1985)343

Epoxy novolac resins
Characterization of epoxy novolac resins using phenyl-bonded phase HPLC 321(1985)325

Epoxy resins
Thermoset resin microstructure by GPC: an anhydride-cured epoxy 316(1984)343

Fingerprinting epoxy resin systems using two-dimensional reversed-phase TLC 318(1985)75

Ergot alkaloids
Partition chromatography of ergot alkaloids on formamide-saturated silica gel columns (in German) 319(1985)440

Erythrocytes
Hydrophobic chromatography of cells: adsorption and resolution on homologous series of alkylagaroses 317(1984)103

Erythropoietin
HPLC separation of serum erythrotropin and erythropoietin 310(1984)396

Erythrotropin
HPLC separation of serum erythrotropin and erythropoietin 310(1984)396

Esmolol
HPLC assay for the major blood metabolite of esmolol: an ultra short acting beta blocker 309(1984)203

Essential oils
see also Volatile oils

GLC-MS in the analysis of essential oils 303(1984)109

GC characterization of frequently occuring sesquiterpenes in essential oils 318(1985)125

Thick-film capillary column GC-field ionization MS. A complementary technique for the rapid analysis of volatiles 322(1985)200

Esters
GC of homologous esters. XXVII. Retention increments of C_1-C_{18} primary alkanols and their 2-chloropropanoyl and 3-chloropropanoyl derivatives on SE-30 and OV-351 capillary columns 319(1985)131

GC of homologous esters. XXVIII. Retention increments of aliphatic C_1-C_{18} *n*-alkyl esters of butanoic acid and its monochloro derivatives on SE-30 and OV-351 capillary columns 320(1985)325

GLC analyses. XXXV. Capillary column studies of C_1-C_{12} *n*-alkyl esters of benzoic and monochlorobenzoic acids 322(1985)83

GLC analyses. XXXVI. Separation of ethyl, 2-chloroethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns with temperature programming 322(1985)97

GLC analyses. XXXVII. Capillary column studies of benzoyl and monochlorobenzoyl esters of lower saturated branched-chain alcohols 323(1985)331

GLC Analyses. XXXVIII. Capillary column studies of unsaturated esters of benzoic and monochlorobenzoic acids 324(1985)113

GC of homologous esters. XXIX. Propanoyl and monochlorpropanoyl esters of lower saturated brached-chain and unsaturated alcohols 324(1985)343

GC of homologous esters. XXX. Unexpected predominancy of polar effects over steric effects in crowded aliphatic esters: a GLC appraisal of specific interactions 324(1985)355

GC of homologous esters. XXXI. Butanoyl and monochlorobutanoyl esters of lower saturated branched chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 325(1985)61

Ethanol
Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.*-butanol 319(1985)247

Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

Ether-bonded silica phases
Wide-pore silica-based ether-bonded phases for separation of proteins by high-performance hydrophobic-interaction and size-exclusion chromatography 316(1984)519

Ethers
GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

Ethinyloestradiol
Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Ethisterone
Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Ethosuximide
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

N-Ethoxycarbonyl-2-ethoxy-1,2-dihydro-quinoline
Immobilization of ligands for affinity chromatography. Coupling on a spacer arm gel with N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline as condensation agent: study of coupling conditions by means of a radioimmunologic method 314(1984)445

Ethoxyquin
Chromatographic studies of the autoxidation products of ethoxyquin and its photochemical conversion 324(1985)462

Ethyl acetate
Volatile products from acetylcholine as markers in the rapid urine test using head-space GLC 307(1984)11

Mixed solvents in GLC. Activity coefficients for chloroform, dichloromethane and ethyl acetate in squalane-dinonyl phthalate mixtures at 303°K 324(1985)231

Ethyl alkanoates
GLC analyses. XXXVI. Separation of ethyl, 2-chloroethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns with temperature programming 322(1985)97

Ethylamine
Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

N,N'-Ethylenebis(acetylacetoneimine) chelates
Developments in the HPLC of metallo-organic compounds 317(1984)569

Ethylenediamineplatinum(II) malonate
Determination of ethylenediamineplatinum(II) malonate in infusion fluids, human plasma and urine by HPLC 310(1984)335

Ethylenediamines
Separation and analysis of ethylenediamines as their *tert.*-butyldimethylsilyl derivatives by GLC and MS 314(1984)455

Ethylenediaminetetraacetate
Simultaneous determination of anions and metal cations by single-column ion chromatography with ethylenediaminetetraacetate as eluent and conductivity and UV detection 312(1984)327

Ethylene dibromide
Separation and use of the perdeuterated analogue as an internal standard for the analysis of ethylene dibromide 312(1984)450

N-Ethylephedrine
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

2-Ethyl-2-phenylmalonediamide
Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Evaporative light-scattering mass detector
Examination of the concentration response of evaporative light-scattering mass detectors 323(1985)297

Exchange capacity
Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

Extra-column effects
Characterization of extracolumn and concentration-dependent distortion of chromatographic peaks by Edgeworth-Cramér series 315(1984)67

Extra-column volume
Role of extra-column volume in the efficiency of high-speed LC 321(1985)69

Extraction solvents
GC analysis of the extraction solvent used in nuclear fuel reprocessing plants 325(1985)75

F

F 2207 VII
Separation of diastereomeric amides by preparative HPLC and analysis of enantiomers on a chiral support (in French) 318(1985)398

Factor analysis
Three-mode factor analysis of data on retention in normal-phase HPLC 301(1984)311

Fats
On-line GC evaluation of iodine values in edible fats and oils 314(1984)233

Fatty acid alcohol esters
Capillary GC investigation of plasma lipid alcoholysis during alcohol extraction 307(1984)172

Fatty acid methyl esters
On-line GC evaluation of iodine values in edible fats and oils 314(1984)233

Evaluation of the mass detector for quantitative detection of triglycerides and fatty acid methyl esters 324(1985)35

Fatty acids
Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173

Studies on volatile metabolites of some potentially pathogenic *Bacillus* species, using automated head-space GC 305(1984)178

Fluorometric HPLC of 9-aminophenanthrene-derivatized free fatty acids 305(1984)261

Lipoxygenase products. A novel GC-MS assay for monohydroxy fatty acids 306(1984)1

Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Determination of bound cellular fatty acids in *Actinobacillus actinomycetemcomitans* and

Haemophilus aphrophilus by GC and GC-MS 308(1984)282

New procedure to analyze free fatty acids. Application to 20-mg brain tissue samples 309(1984)33

Simultaneous separation and sensitive determination of free fatty acids in blood plasma by HPLC 309(1984)43

Reliable measurement of non-esterified long-chain fatty acid pattern in blood plasma 311(1984)9

Whole-cell methanolysis as a rapid method for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)347

Determination of fatty acids by HPLC of Dns-ethanolamine derivatives 312(1984)461

Analysis of phospholipids in cow's milk by high-temperature injection GC and HPLC 315(1984)223

Detection of volatile fatty and dicarboxylic acids on alkaline pherograms with Mills' reagent 318(1985)446

HPLC of short- and long-chain fatty acids as 2-nitrophenylhydrazides 321(1985)165

Electrokinetic detection of lauric, myristic, stearic and arachidic acids separated by reversed-phase HPLC 321(1985)433

Analysis of saturated normal fatty acids in hydrocarbon matrices by capillary isotachophoresis 325(1985)151

Fenethazine
HPLC of haloperidol in serum at the concentrations achieved during chronic therapy 308(1984)387

Fenfluramine
Determination of fenfluramine and norfenfluramine in plasma using a nitrogen-sensitive detector 310(1984)412

Fenoldopam
Determination of fenoldopam (SK&F 82526) and its metabolites in human plasma and urine by HPLC with electrochemical detection 317(1984)463

Fenretinide
Stability-indicating reversed-phase HPLC assay for fenretinide drug substances 322(1985)380

Ferrichromes
Separation of ferrichromes and other hydroxamate siderophores of fungal origin by reversed-phase chromatography 301(1984)247

Ferritin
Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157

Ferrochelatase
Separation of haem compounds by reversed-phase ion-pair HPLC and its application in the assay of ferrochelatase activity 310(1984)31

Feruloylquinic acid
Chromatographic separation and identification of naturally occurring chlorogenic acids by ^1H nuclear magnetic resonance spectroscopy and MS 315(1984)253

Fetoprotein
HPLC of proteins: purification of α-fetoprotein from fetal calf serum 310(1984)19

Field flow fractionation
Sampling and flow effects in sedimentation field flow fractionation 315(1984)45

Field ionization mass spectrometry
Thick-film capillary column GC-field ionization MS. A complementary technique for the rapid analysis of volatiles 322(1985)200

Fish species
Identification of fish species by HPLC 324(1985)199

FK-027
Determination of a new orally active cephalosporin in human plasma and urine by HPLC using automated column switching 311(1984)339

Flavonoids
TLC behaviour and chemical structure of 6- and 8-methoxy-5-hydroxyflavones 315(1984)101

HPLC analysis of coumarins and flavonoids from section Tridentatae of *Artemisia* 322(1985)236

Flecainide
HPLC method for the measurement of mexiletine and flecainide in blood plasma or serum 307(1984)439

Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Flordipine
HPLC analysis of flordipine in human plasma 308(1984)382

Flow characteristics
Sampling and flow effects in sedimentation field flow fractionation 315(1984)45

Flow-rate
Optimizing flow-rates in capillary GC-MS 325(1985)23

Fluid inclusions
GC analysis of volatiles in fluid and gas inclusions 302(1984)181

Flumethasone
Analysis of flumethasone pivalate formulations by HPLC 301(1984)477

Flunitrazepam
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Fluocortin butyl
Determination of trace levels of steroids in blood plasma by LC with peroxyoxalate chemiluminescence detection 317(1984)355

Fluoranil
Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

9-Fluorenylmethylchloroformate
HPLC assay of pirlimycin in human serum and urine using 9-fluorenylmethylchloroformate 309(1984)95

Fluorescamine
Rapid fluorescamine based protein assay usable in the presence of interfering substances 319(1985)446

Fluorescence reagents
Uranyl salts as fluorescence reagents for the TLC detection of complex-forming compounds (in German) 322(1985)272

Use of benzoin as pre-column fluorescence derivatization reagent for the HPLC of angiotensins 322(1985)473

Fluorescent whitening agents
HPTLC of fluorescent whitening agents and their identification in detergents 322(1985)363

1-Fluoro-2,4-dinitrobenzene
Pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene 322(1985)321

4-Fluoroglutamic acid
Simple method for the separation of diastereomers of 4-fluoroglutamic acid by GLC 315(1984)421

5-Fluorouracil
Separation of 5-fluorouracil and uracil by ion-pair reversed-phase HPLC on a column with porous polymeric packing 307(1984)464

Fluorescence determination of 5-fluorouracil and 1-(tetrahydro-2-furanyl)-5-fluorouracil in blood serum by HPLC 310(1984)151

Flupirtine
Quantitation of flupirtine and its active acetylated metabolite by reversed-phase HPLC using fluorometric detection 305(1984)135

Flurazepam
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Folacin conjugase
Determination of folylpoly-γ-glutamate carboxypeptidase (folacin conjugase) activity using reversed-phase HPLC 318(1985)387

Folylpoly-γ-glutamate carboxypeptidase
Determination of folylpoly-γ-glutamate carboxypeptidase (folacin conjugase) activity using reversed-phase HPLC 318(1985)387

Food thickeners
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Formaldehyde
Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 308(1984)121

Formamide
Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Formyl substitution
GLC analyses. XL. Incremental effects of a formyl group introduced into the *ortho-* and *para-*positions of isomeric chlorophenols 324(1985)192

Fostedil
HPLC determination of a new calcium antagonist, fostedil, in plasma and urine using fluorescence detection 305(1984)233

Fragilin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Fritschiellaxanthin
Direct diastereomeric resolution of carotenoids. I. 3-Hydroxy-4-oxo β-end group 318(1985)122

Frontal chromatography
Rôle of displacement effects in trapping trace components from gases in sorbent-packed columns 324(1985)269

Fructan oligomers
Determination of fructan oligomers of degree of polymerization 2-30 by HPLC 303(1984)417

Fructose
HPLC separation of glucose-1-phosphate, fructose, sucrose and inorganic orthophosphate 322(1985)220

Ftorafur
Fluorescence determination of 5-fluorouracil and 1-(tetrahydro-2-furanyl)-5-fluorouracil in blood serum by HPLC 310(1984)151

Fucose
Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus*

aphrophilus based on carbohydrates in lipopolysaccharide 310(1984)261

Fulvic acids
Isolation, identification and determination of 3-(4-hydroxyphenyl)propionic acid by HPLC of the phenolic compounds of *Loiseleuria procumbens* Desf. (Ericaceae) (in French) 322(1985)257

Fumarase
Sepharose derivatives containing citric acid as affinity ligand. Purification of fumarase 319(1985)173

Functional group structure
Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

Furans
Study of chlorinated diphenyl ethers and chlorinated 2-phenoxyphenols as interferences in the determination of chlorinated dibenzo-*p*-dioxins and chlorinated dibenzofurans in biological samples 303(1984)351

Furcellaran
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Furfural
Single-valve, single-pump, HPLC column-switching analysis of hydrolyzed wood components 318(1985)269

Furosemide
Quantitative determination of furosemide in plasma, plasma water, urine and ascites fluid by HPLC 308(1984)241

Fused-silica capillary tubing
Preliminary investigation of the effects of gamma-radiation on the surface of raw fused-silica tubing 302(1984)341

G

Galactosamine
Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Galactose
Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

β-Galactosidase
Headspace GC determination of β-galactosidase activity using electron-capture detection 321(1985)385

Galactosyl hydroxylysine
Measurement of hydroxylysine glycosides in urine and its application to spinal cord injury 308(1984)65

Gamma-radiation
Preliminary investigation of the effects of gamma-radiation on the surface of raw fused-silica tubing 302(1984)341

Gangliosides
Quantitation of gangliosides in the picomolar range 305(1984)512

Separation of ganglioside molecular species, with homogeneous long-chain base composition, by reversed-phase TLC 315(1984)395

Separation of gangliosides by anion-exchange chromatography on Mono Q™ 322(1985)465

Garryine
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Gases
Polymeric Schiff base complexes as solid phases in GC 322(1985)63

Gas inclusions
GC analysis of volatiles in fluid and gas inclusions 302(1984)181

Gasoline
Quantitative analysis of hydrocarbons in gasolines by capillary GLC. I. Isothermal analysis 303(1984)151

Gel electrophoresis
Comparison of high-performance ion-exchange chromatography and gel electrophoresis in protein separations 318(1985)358

Gels
Evaluation of hydrophobicity of gels by use of some 1-alkanols as pilot solutes 301(1984)57

Gemfibrozil
GC determination of gemfibrozil and its metabolites in plasma and urine 307(1984)210

Gentamicin
Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Gentisic acid
Rapid and sensitive determination of acetylsalicylic acid and its metabolites using reversed-phase HPLC 310(1984)343

Improved method for the determination of aspirin and its metabolites in biological fluids by HPLC: applications to human and animal studies 311(1984)301

Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133

Ginger
HPLC separation of pungency components of ginger 319(1985)471

Gingerols
HPLC separation of pungency components of ginger 319(1985)471

Ginsenosides
Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497

Glaucine
Determination of glaucine in plasma and urine by HPLC 308(1984)273

Globin
Electrophoretic studies of blood globin preparations 324(1985)395

Globins
Analysis of the globins from fast human haemoglobins by isoelectrofocusing on polyacrylamide gel rods 307(1984)91

D-Gluconic acid
Determination of D-gluconic, 5-keto-D-gluconic, 2-keto-D-gluconic and 2,5-diketo-D-gluconic acids by HPLC 312(1984)211

D-Gluco-oligosaccharides
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Glucose
Effect of the presence of glucose on the determination of amino acids by GC 309(1984)375

Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Glucose 1-phosphate
HPLC separation of glucose-1-phosphate, fructose, sucrose and inorganic orthophosphate 322(1985)220

Glucose 6-phosphate dehydrogenase
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Glucosylgalactosyl hydroxylysine
Measurement of hydroxylysine glycosides in urine and its application to spinal cord injury 308(1984)65

Glucuronides
Separation, isolation and identification of optical isomers of 1,4-benzodiazepine glucuronides from biological fluids by reversed-phase HPLC 306(1984)231

Glucuronyl transferase
Rapid and sensitive assays using HPLC to measure the actitivies of phase II drug metabolising enzymes: glucuronyl transferase and sulfotransferase 301(1984)282

Glutamic acid
Analysis of glutamine, glutamic acid and pyroglutamic acid in protein hydrolysates by HPLC 322(1985)248

GC resolution of substituted glutamic acid enantiomers 325(1985)444

Glutamine
GLC assay for asparagine and glutamine 322(1985)228

Analysis of glutamine, glutamic acid and pyroglutamic acid in protein hydrolysates by HPLC 322(1985)248

Glutathione
Analysis of hepatic reduced glutathione, cysteine and homocysteine by cation-exchange HPLC with electrochemical detection 308(1984)83

Use of LC with dual-electrode electrochemical detection in the investigation of glutathione oxidation during benzene metabolism 317(1984)579

Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Glutethimide
Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444

Glyceraldehyde phosphate dehydrogenase
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

D-Glycero-D-mannoheptose
Carbohydrates of whole defatted cells as a basis for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)31

L-Glycero-D-mannoheptose
Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Glycine
Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251

Application of reversed-phase HPLC using pre-column derivatization with *o*-phthaldialdehyde for the quantitative analysis of amino acids in adult and fetal sheep plasma, animal feeds and tissues 315(1984)321

Glycinehydroxamate-Sepharose 6B
Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Glycopeptides
Purification of glycopeptides of human plasma proteins by HPLC 317(1984)11

Fractionation of peptic glycopeptides from chicken ovalbumin by reversed-phase HPLC 320(1985)426

Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173

Glycoproteins
Isolation and purification of HLA-DR antigens from tumour cells by affinity chromatography and chromatofocusing 308(1984)322

Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Automatic amino acid and sugar analysis of glycoproteins 317(1984)35

Molecular weight measurement of human satietin 317(1984)165

Isotachophoresis of allergenic extracts 320(1985)205

Purification of a basic glycoprotein allergen from pollen of timothy by HPLC 321(1985)199

Glycopyrronium
Quantitative capillary column GC method for the determination of glycopyrronium in human plasma 308(1984)143

Glycosides
Electrochemical detection of cyanogenic glycosides after enzymatic post-column cleavage 303(1984)67

Cyanogenic glycosides and cyanohydrins in plant tissues. Qualitative and quantitative determination by enzymatic post-column cleavage and electrochemical detection, after separation by HPLC 303(1984)77

Investigation of conditions of the preparative LC separation of cardiac glycosides present in *Digitalis lanata* 303(1984)179

Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Glycosphingolipids
Separation of ganglioside molecular species, with homogeneous long-chain base composition, by reversed-phase TLC 315(1984)395

Glycylxylidide
Determination of lidocaine and active metabolites in blood serum by LC with electrochemical detection 306(1984)269

Simultaneous determination of lidocaine and its deethylated metabolites using GLC with nitrogen-phosphorus detection 307(1984)200

Simultaneous determination of lidocaine and its metabolites in plasma and myocardium 311(1984)291

Glycyrrhetinic acid
Separation and quantitative analysis of 18α-glycyrrhetinic acid and 18β-glycyrrhetinic acid in Glycyrrhizae Radix by GLC 320(1985)430

Gossypol
HPLC determination of gossypol in plasma 307(1984)235

Gradient elution
Gradient elution micellar LC 316(1984)183

Examination of theoretical principles of gradient elution as applied to reversed-phase HPLC separation of phenylthiohydantoin amino acids 316(1984)359

Versatile microcomputer-controlled, automated gradient analytical HPLC system 316(1984)407

Simple device for sequential linear gradient elution with a single, conventional, one-piston reciprocating pump 316(1984)415

Grandidentatin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Graphite
GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19

Graphitized carbon black
Sample enrichment for GC-MS analysis of polynuclear aromatic hydrocarbons in water and in organic mixtures 302(1984)167

Growth factors
Use of HPLC in the isolation and characterisation of mouse and rat epidermal growth factors and examination of apparent heterogeneity 305(1984)295

Growth hormone
Comparison by gel filtration chromatography and reversed-phase HPLC of the immunoreactive growth hormone composition of a human pituitary extract 324(1985)385

Growth inhibitory glycopeptide
Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173

Guaifenesin
Selection of mixed ion-pair modifiers for HPLC mobile phases 314(1984)402

Guanidine hydrochloride
Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395

Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173

Guanidinium chloride
Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Guanidinium sulphate
Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Guanidinobutyric acid
HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257

Guanidino compounds
HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281

Guanidinosuccinic acid
HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257

Guar
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Gum arabic
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

H

Haemagglutination enhancing factor
Characterization of a heterogeneous chicken plasma protein, HEF, by analytical isotachophoresis in agarose gel 320(1985)119

Haemoglobin
Fast protein liquid chromatofocusing of lyophilized native haemoglobin and its chemically modified derivatives 303(1984)422

Separation of glyoxylated hemoglobin with varying oxygen affinity by polyanion fast protein LC 303(1984)425

Analytical peptide mapping by ion-exchange HPLC: application to haemoglobin variants 305(1984)456

Analysis of the globins from fast human haemoglobins by isoelectrofocusing on polyacrylamide gel rods 307(1984)91

Determination of glycosylated haemoglobin by isoelectric focusing in non-linear pH gradients 307(1984)103

HPLC separations on dihydroxyboryl-agarose 316(1984)301

Separation of hemoglobin variants by ion-exchange chromatography on monobead resins 318(1985)325

Reversed-phase HPLC of human haemoglobin chains 321(1985)407

Haemopexin
Fractionation of human serum proteins by immobilized metal affinity chromatography 315(1984)167

Purification of glycopeptides of human plasma proteins by HPLC 317(1984)11

Haemophilus aphrophilus
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Determination of bound cellular fatty acids in *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* by GC and GC-MS 308(1984)282

Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Carbohydrates of whole defatted cells as a basis for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)31

Whole-cell methanolysis as a rapid method for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)347

Haems
Separation of haem compounds by reversed-phase ion-pair HPLC and its application in the assay of ferrochelatase activity 310(1984)31

Hall electrolytic conductivity detector
Determinations of chlorinated hydrocarbons by GC using response factor calibration 314(1984)243

Halocarbons
GC determination of C_2-C_8 hydrocarbons and halocarbons in ambient air by simultaneous use of three detectors 323(1985)323

α-Halogenocarboxylic acids
Resolution of α-halogenocarboxylic acids by GC on (*R*)-N-lauroyl-α-(1-naphthyl)ethylamine 318(1985)235

Haloperidol
Haloperidol determination in serum and cerebrospinal fluid using GLC with nitrogen-phosphorus detection: application to pharmacokinetic studies 307(1984)194

HPLC of haloperidol in serum at the concentrations achieved during chronic therapy 308(1984)387

Halothane
Analytical isotachophoresis utilizing computer simulation. II. Assessment of optimum separation conditions for urinary trifluoroacetic acid metabolized from anaesthetic halothane 305(1984)429

Headspace analysis
Headspace gas analysis: the role and the design of concentration traps specifically suitable for capillary GC 321(1985)45

Headspace GC determination of β-galactosidase activity using electron-capture detection 321(1985)385

Heat capacity
Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

Heats of formation
Heats of formation-GC retention relationships for aliphatic hydrocarbons and alkylbenzenes 322(1985)35

Heavy metals
Determination of heavy metals by isotachophoresis 320(1985)263

α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81

Heliotric acid
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

Helium ionization detector
New design for helium ionization detection 302(1984)257

Heptafluorobutyric acid amides
Trace analysis of amines and isocyanates using glass capillary GC and selective detection. III. Determination of aliphatic and alicyclic amines as perfluoro fatty acid amides using electron-capture and nitrogen-selective detection 303(1984)89

Heptaminol
LC assay of heptaminol in serum and its oral pharmacokinetics in the dog 307(1984)351

Hexachlorocyclohexane
Selected ion monitoring MS method for the determination of hexachlorocyclohexane isomers in soils 312(1984)403

Hexachlorophene
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Hexadecyl acetate
Microanalytical methods in the structure elucidation of sex-specific components in the large pine weevil, *Hylobius abietis* L. (*Coleoptera, Curculionidae*) 323(1985)343

2H-Hexafluoropropanol-2
Determination of 2H-hexafluoropropanol-2 in air 319(1985)407

***n*-Hexanol**
Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79

HI-6-
Stability study of HI-6 dichloride in various anticholinergic formulations 315(1984)389

High-frequency contactless detector
Improvement of the performance of a high-frequency contactless conductivity detector for isotachophoresis 320(1985)233

High mobility group nonhistone chromosomal proteins
Purification of high mobility group nonhistone chromosomal proteins by LC on a column containing immobilized histone H5 314(1984)436

High-performance ion-exchange chromatography
Comparison of high-performance ion-exchange chromatography and gel electrophoresis in protein separations 318(1985)358

Hindered-amine light stabilisers
Determination of polymeric hindered-amine light stabilisers in plastics by pyrolysis-GC 325(1985)87

Hippurate
Simple HPLC assay for benzylamine oxidation products in cell suspensions 305(1984)256

Histamine
Sensitive GLC procedure for urinary Nτ-methylimidazole acetic acid, an index of histamine turnover 306(1984)69

Histamine assay in tears by fluorescamine derivatization and HPLC 310(1984)390

Reversed-phase ion-pair HPLC procedure for determination of histamine and its metabolites in rat urine 312(1984)443

Histidine
Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

Histidine decarboxylase
Assay for histidine decarboxylase in rat stomach and brain by HPLC with fluorescence detection 311(1984)390

Histograms
Use of histograms in computer-aided comparison of chromatograms 318(1985)247

HLA-DR antigens
Isolation and purification of HLA-DR antigens from tumour cells by affinity chromatography and chromatofocusing 308(1984)322

Homocysteine
Analysis of hepatic reduced glutathione, cysteine and homocysteine by cation-exchange HPLC with electrochemical detection 308(1984)83

Homologous series
Reversed-phase LC of homologous series. A general method for prediction of retention 314(1984)13

Homospermidine
GC-MS of polyamines as their N-ethyloxycarbonyl derivatives and identification of *sym*-homospermidine and *sym*-norspermine in mosses and ferns 303(1984)99

Homovanillic acid
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Measurement of homovanillic acid in human plasma by HPLC with electrochemical detection 309(1984)379

Analysis of vanillylmandelic acid, homovanillic acid and 5-hydroxyindoleacetic acid in human urine by HPLC and fluorometry 310(1984)379

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Direct determination of urinary vanillylmandelic acid and homovanillic acid by HPLC on an anion-exchange column 311(1984)369

Selective retention of catecholamines and their derivatives in reversed-phase ion-pair partition chromatography 322(1985)43

Horizontal flow-through coil planet centrifuge
Purification of a cholecystokinin fragment peptide in the horizontal flow-through coil planet centrifuge 301(1984)277

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. I. Hydrodynamic distribution of two solvent phases in a helical column subjected to two types of synchronous planetary motion 301(1984)377

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. II. Phase distribution diagrams for helical and spiral columns 301(1984)387

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

Horizontal flow-through coil planet centrifuge: some practical applications of countercurrent chromatography 315(1984)287

Hormones
see also Plant hormones and steroids

Reversed-phase HPLC and separation of peptide hormones in human tissues and plasma with reference to cholecystokinins 305(1984)43

Semi-micro quantitative analysis of complex urinary steroid mixtures in healthy and diseased states 306(1984)39

Rapid measurement of oestradiol and oestriol by HPLC after automatic pretreatment 311(1984)249

HPLC determination of bovine and porcine insulins in a commercial injection 312(1984)476

Reversed-phase HPLC studies of α-MSH fragments 314(1984)313

Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313

General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470

Laser fluorimetry for capillary column LC: high-sensitivity detection of derivatized biological compounds 317(1984)443

TLC of ecdysteroids 318(1985)373

Subunits of bovine lutropin. Hormonal and immunological activity after separation with reversed-phase HPLC 319(1985)195

Convenient method for the determination of picomole amounts of juvenile hormone 323(1985)355

Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395

Comparison by gel filtration chromatography and reversed-phase HPLC of the immunoreactive growth hormone composition of a human pituitary extract 324(1985)385

Humic acids
Isolation, identification and determination of 3-(4-hydroxyphenyl)propionic acid by HPLC of the phenolic compounds of *Loiseleuria procumbens* Desf. (Ericaceae) (in French) 322(1985)257

Hyaluronate oligosaccharides
Comparison of gel permeation and ion-exchange chromatographic procedures for the separation of hyaluronate oligosaccharides 320(1985)351

Hydantoin-based chiral stationary phases
Preparation and use of hydantoin-based chiral stationary phases 322(1985)309

Hydration model
Retention of carbohydrates on silica and amine-bonded silica stationary phases: application of the hydration model 325(1985)287

Hydrocarbons
Trace analysis of organic compounds by HPLC with photoionization detection 302(1984)43

New schemes for the electron-capture sensitization of aromatic hydrocarbons 302(1984)243

Quantitative analysis of hydrocarbons in gasolines by capillary GLC. I. Isothermal analysis 303(1984)151

HPLC procedure for direct determination of the compound class composition of diesel and kerosine fuels using refractive index detection 312(1984)237

Heats of formation-GC retention relationships for aliphatic hydrocarbons and alkylbenzenes 322(1985)35

Analysis of Australian crude oils by high-resolution capillary GC-MS 323(1985)305

GC determination of C_2-C_8 hydrocarbons and halocarbons in ambient air by simultaneous use of three detectors 323(1985)323

GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

Hydrochlorothiazide
HPLC determination of hydrochlorothiazide in plasma and urine 307(1984)445

Hydrodynamic distribution
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. I. Hydrodynamic distribution of two solvent phases in a helical column subjected to two types of synchronous planetary motion 301(1984)377

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. II. Phase distribution diagrams for helical and spiral columns 301(1984)387

Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

Hydrodynamic volume
Effect of solute size and shape on the elution volume in size-exclusion chromatography 319(1985)115

Hydrogen-atmosphere flame ionization detector
Ionization quenching of a hydrogen-atmosphere flame as a detection method for GC 303(1984)331

Hydrogen bonding
Dependence of the retention of phenols upon Van der Waals volume, π-energy and hydrogen-bonding effects 302(1984)89

Hydrogen sulphide
GC study of the analysis and elution mechanism of hydrogen sulphide, carbonyl sulphide and light mercaptans in petroleum gases 321(1985)59

Hydromorphone
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Hydrophobic-interaction chromatography
Hydrophobic interaction HPLC of proteins 317(1984)55

Salt-mediated retention of proteins in hydrophobic interaction chromatography. Application of solvophobic theory 317(1984)67

Comparison of hydrophobic-interaction and reversed-phase chromatography of proteins 317(1984)141

Hydrophobic interactions
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47

Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Hydrophobicity
see also Lipophilicity

Evaluation of hydrophobicity of gels by use of some 1-alkanols as pilot solutes 301(1984)57

Comparison of phenyl- and octadecyl-modified silica gel as stationary phase for the prediction of *n*-octanol-water partition coefficients by HPLC 320(1985)271

R_m values, retention times and π values of a series of potentially mutagenic nitroimidazo[2,1-*b*]thiazoles 320(1985)281

Hydrophobic substituent constant
R_m values, retention times and π values of a series of potentially mutagenic nitroimidazo[2,1-*b*]thiazoles 320(1985)281

Hydroquinone
Use of LC with dual-electrode electrochemical detection in the investigation of glutathione oxidation during benzene metabolism 317(1984)579

***p*-Hydroquinones**
HPLC of substituted *p*-benzoquinones and *p*-hydroquinones. I. Interplay of on-column redox reaction and the chromatographic retention process 316(1984)151

Hydroxy acids
Phosgene: a versatile reagent for enantiomer separation by capillary GC 301(1984)129

Studies of cyclic di-*tert.*-butylsilylene derivatives of diols and hydroxy acids by GC-MS 315(1984)119

Hydroxyalkanesulphonates
Analysis of α-olefinsulphonates by HPLC 325(1985)43

β-Hydroxyamino acids
Quantitation of the enantiomers of β-hydroxyamino acids by GC resolution on an optically active packed column 301(1984)461

10-Hydroxyamitriptyline
GC-MS determination of amitriptyline and its major metabolites in human serum 305(1984)73

***p*-Hydroxyamphetamine**
Simple determination of *p*-hydroxyamphetamine by HPLC with electrochemical detection 306(1984)394

Determination of methamphetamine and its metabolites in rat tissues by GC with a nitrogen-phosphorus detector 318(1985)307

9α-Hydroxyandrost-4-en-3,17-dione
Determination of 9α-hydroxyandrost-4-en-3,17-dione in fermentation solutions by TLC (in German) 319(1985)475

***p*-Hydroxybenzoic acid**
Determination of preservatives in cosmetic products by reversed-phase HPLC 315(1984)465

Hydroxycarboxylic acids
GLC separation of hydroxy monocarboxylic acids and dicarboxylic acids on a fused-silica capillary column 301(1984)273

Hydroxycarboxylic and oxocarboxylic acids in urine: products from branched-chain amino acid degradation and from ketogenesis 309(1984)225

Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

Isotachophoretic analysis of saccharinic acids 320(1985)111

7-Hydroxychlorpromazine
Isotopic fractionation of N-([^{3}H]methyl)-chlorpromazine and N-([^{3}H]methyl)-7-hydroxychlorpromazine by reversed-phase HPLC 303(1984)412

Hydroxycinnamic acid esters
Analysis of hydroxycinnamic acid esters and their glucosides by reversed-phase HPLC after polyamide separation 315(1984)243

Hydroxycinnamic acid glucosides
Analysis of hydroxycinnamic acid esters and their glucosides by reversed-phase HPLC after polyamide separation 315(1984)243

17-Hydroxycorticosteroids
New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188

18-Hydroxycorticosterone
Application of reversed-phase HPLC for radioimmunoassay of plasma 18-hydroxycorticosterone 308(1984)295

7-Hydroxycoumaro[5,6-*c*]cinnoline
Improved analytical and preparative methods for necatorin from *Lactarius necator* (Fr.) Karst mushroom 301(1984)303

10-Hydroxydesmethylnortriptyline
Amitriptyline and its basic metabolites determined in plasma by GC 308(1984)165

20-Hydroxyecdysone
Analytical droplet counter-current chromatography isolation of 20-hydroxyecdysone from *Vitex thyrsiflora* (Verbenaceae) 321(1985)246

5-Hydroxyeicosatetraenoic acid
Preparation of pentafluorobenzyl esters of arachidonic acid lipoxygenase metabolites. Analysis by GC and negative-ion chemical ionization MS 305(1984)3

Hydroxyeicosatetraenoic acids
Simple single-step HPLC method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 308(1984)316

Determination of leukotrienes and prostaglandins in [^{14}C]arachidonic acid labelled human lung tissue by HPLC and radioimmunoassay 311(1984)39

9-Hydroxyellipticine
Use of electrochemical detection in the HPLC determination of hydroxylated ellipticine derivatives 309(1984)170

5-Hydroxyindole-3-acetic acid
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

Rapid and simple method for the simultaneous determination of 3,4-dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by HPLC with electrochemical detection 308(1984)55

Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53

Determination of urinary tryptophan, 5-hydroxytryptamine and 5-hydroxyindoleacetic acid in neonatal hyperbilirubinaemic infants using reversed-phase HPLC with fluorescence detection 309(1984)145

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Analysis of vanillylmandelic acid, homovanillic acid and 5-hydroxyindoleacetic acid in human urine by HPLC and fluorometry 310(1984)379

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Isocratic analysis of 3-methoxy-4-hydroxyphenyl glycol, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in cerebrospinal fluid by HPLC with amperometric detection 311(1984)354

HPLC of 5-hydroxyindole-3-acetic acid in urine with direct sample injection 311(1984)380

α-Hydroxyisobutyrate complexes
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11

Hydroxyl aromatic compounds
Mobile phase optimization for the reversed-phase LC of complex hydroxyl aromatic mixtures 324(1985)1

Hydroxylysine
TLC of hydroxylysine for collagen analysis 311(1984)400

Hydroxylysine glycosides
Measurement of hydroxylysine glycosides in urine and its application to spinal cord injury 308(1984)65

***p*-Hydroxymethamphetamine**
Determination of methamphetamine and its metabolites in rat tissues by GC with a nitrogen-phosphorus detector 318(1985)307

4-Hydroxy-3-methoxyphenylacetic acid
Rapid and simple method for the simultaneous determination of 3,4-dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by HPLC with electrochemical detection 308(1984)55

Isocratic analysis of 3-methoxy-4-hydroxyphenyl glycol, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in cerebrospinal fluid by HPLC with amperometric detection 311(1984)354

Hydroxymethylated porous polystyrene
Determination of creatinine in human, dog and rat urine by HPLC on a column of hydroxymethylated porous polystyrene 307(1984)399

Hydroxymethylfurfural
Single-valve, single-pump, HPLC column-switching analysis of hydrolyzed wood components 318(1985)269

LC of tomato paste serum with sample preparation by ultrafiltration 325(1985)450

Hydroxymethylnucleosides
Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 308(1984)121

Hydroxymethylserine
Quantitation of the enantiomers of β-hydroxyamino acids by GC resolution on an optically active packed column 301(1984)461

3-Hydroxymyristic acid
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

10-Hydroxynortriptyline
GC-MS determination of amitriptyline and its major metabolites in human serum 305(1984)73

N-Hydroxyphenacetin
MS determination of N-hydroxyphenacetin in urine using multiple metastable peak monitoring following TLC 310(1984)179

3-(4-Hydroxyphenyl)propionic acid
Isolation, identification and determination of 3-(4-hydroxyphenyl)propionic acid by HPLC of the phenolic compounds of *Loiseleuria procumbens* Desf. (Ericaceae) (in French) 322(1985)257

***p*-Hydroxyphenytoin**
Analysis of *p*-hydroxyphenytoin in microsomal reactions by HPLC with electrochemical detection 310(1984)418

4-Hydroxypropranolol
Silica gel HPLC for the simultaneous determination of propanolol and 4-hydroxypropranolol enantiomers after chiral derivatization 310(1984)424

4'-Hydroxypropranolol sulphate
Isolation from urine of 4'-hydroxypropranolol sulphate, a major new propranolol metabolite, by ion-pair extraction 305(1984)250

5-Hydroxytryptamine
see Serotonin

5-Hydroxytryptophan
Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Hyfrane
GC analysis of the extraction solvent used in nuclear fuel reprocessing plants 325(1985)75

Hyoscyamine
Qualitative and quantitative analysis of hyoscyamine-atropine and scopolamine in mother tinctures of Solanaceae by HPLC (in French) 319(1985)412

Hypoxanthine
Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Measurement of adenosine, inosine, and hypoxanthine in human plasma 307(1984)404

Simultaneous determination of allopurinol, oxipurinol, hypoxanthine and xanthine in biological fluids by HPLC 307(1984)469

Separation of adenosine and its derivatives by TLC on silica gel 325(1985)317

Hypoxanthine arabinoside
HPLC analysis of hypoxanthine arabinoside in plasma 307(1984)410

I

Ibuprofen
Silica capillary GC determination of ibuprofen in serum 307(1984)206

Determination of ibuprofen by HPLC 308(1984)350

Determination of ibuprofen in human plasma by HPLC 311(1984)443

Ifosfamide
Rapid GC determination of ifosfamide in biological fluids 311(1984)194

Imidazole
HPLC determination of imidazole in biological fluids 308(1984)310

Imidazoleacetic acid
Reversed-phase ion-pair HPLC procedure for determination of histamine and its metabolites in rat urine 312(1984)443

Imidazopyrimidines
HPLC of phytotoxic substances. II. Behaviour of imidazopyrimidines 315(1984)384

Imino acids
Sensitive detection of amino acids in human serum and dried blood disc of 3 mm diameter for diagnosis of inborn errors of metabolism 309(1984)279

Iminodiacetate
Development of immobilized metal affinity chromatography. I. Comparison of two iminodiacetate gels 323(1985)247

Iminopeptides
Isotachophoretic analysis of iminopeptides in the urine of patients with iminopeptiduria 305(1984)204

Imipenem
Determination of imipenem (N-formimidoyl thienamycin) in human plasma and urine by HPLC, comparison with microbiological methodology and stability 310(1984)71

Imipramine
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Immobilized enzymes
Use of immobilized amino acid oxidase as post-column reactor in the HPLC of amino acids 303(1984)396

Immobilized metal affinity chromatography
Development of immobilized metal affinity chromatography. I. Comparison of two iminodiacetate gels 323(1985)247

Development of immobilized metal affinity chromatography. II. Interaction of amino acids with immobilized nickel iminodiacetate 323(1985)255

Development of immobilized metal affinity chromatography. III. Interaction of oligopeptides with immobilized nickel iminodiacetate 323(1985)265

Immobilized protein stationary phases
Direct LC separation of enantiomers on immobilized protein stationary phases. IV. Molecular interaction forces and retention behaviour in chromatography on bovine serum albumin as a stationary phase 316(1984)617

Immunoglobulin D
Purification of glycopeptides of human plasma proteins by HPLC 317(1984)11

Immunoglobulin G
High-performance affinity chromatography: a rapid technique for the isolation and quantitation of IgG from cerebral spinal fluid 317(1984)173

Immunoglobulins
Comparison of chromatographic characteristics of a series of homologous Bence-Jones proteins during size-exclusion chromatography by HPLC and by Sephadex 323(1985)418

Impurities
Dynamics of isotachophoretic separation. II. Impurities in the electrolytes and sample injection 320(1985)175

Indirect UV detection
Indirect UV detection of non-absorbing solutes in reversed-phase HPLC with the help of UV-active, non-ionic mobile phase constituents 312(1984)345

Indole
GC determination and MS confirmation of traces of indole and 3-methylindole (skatole) in pig back fat 324(1985)469

3-Indoleacetaldoxime
High-performance LC separation and some properties of (*E*)- and (*Z*)-3-indoleacetaldoxime 318(1985)95

Indole-3-acetic acid
Comparison of HPLC and GC-MS for the analysis of indole-3-acetic acid in brain tissue 306(1984)338

Simultaneous determination of tryptamine and its metabolites in mouse brain by HPLC with fluorometric detection 311(1984)385

Trace enrichment of indole-3-acetic acid from leaf matrix 325(1985)127

Separation of conjugates and oxidative metabolites of indole-3-acetic acid by HPLC 325(1985)340

Indole alkaloids
Qualitative and quantitative determinations of hallucinogenic components of psilocybe mushrooms by reversed-phase HPLC 312(1984)467

Indoleamines
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

HPLC determination of hallucinogenic indoleamines with simultaneous UV, photometric and voltammetric detection 320(1985)414

Use of impregnated reagents in the pentafluorobenzylation of indoleamine metabolites 325(1985)309

Indomethacin
HPLC method for the determination of indomethacin and its two primary metabolites in urine 306(1984)315

Industrial autofocusing
Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213

Injection techniques
Evaluation of automated splitless and manual on-column injection techniques using capillary GC for pesticide residue analysis 314(1984)413

"Inverse" injection system for GLC with capillary columns 322(1985)206

Band broadening in space in splitless injection 324(1985)251

Injectors
New injection syringe for GC 322(1985)55

Inorganic anions
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography; factors that control retention and selectivity 301(1984)25

Methods for the analysis of inorganic anions. IV. Reversed-phase HPLC with aqueous hydrophobic ion pairs as eluents 302(1984)15

Simultaneous determination of anions and metal cations by single-column ion chromatography with ethylenediaminetetraacetate as eluent and conductivity and UV detection 312(1984)327

Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

UV visualization of inorganic anions by reversed-phase ion-interaction chromatography: factors that control sensitivity and detection 316(1984)211

Studies on sample preconcentration in ion chromatography. I. Design of an automated, single pump preconcentration system with direct UV absorbance detection 318(1985)279

Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Ion chromatography of inorganic anions with potentiometric detection using a metallic copper electrode 321(1985)363

Inorganic compounds
Inorganic GC (Review) 313(1984)3

Recent developments in ion chromatography (Review) 313(1984)47

Isotachophoresis in inorganic chemistry (Review) 313(1984)189

Inorganic ions
Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Application of indirect potentiometric detection with a metallic copper electrode to ion chromatography of transition metal ions 324(1985)319

Inorganic pyrophosphatase
Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Inosine
Measurement of adenosine, inosine, and hypoxanthine in human plasma 307(1984)404

Separation of adenosine and its derivatives by TLC on silica gel 325(1985)317

Insulin
HPLC determination of bovine and porcine insulins in a commercial injection 312(1984)476

Interaction energy parameters
Prediction of retention data by using parameters of intermolecular interaction. I. A model for calculation of non-polar and polar interaction parameters 315(1984)75

Prediction of retention data by using parameters of intermolecular interaction. II. Application of the model to GLC systems 315(1984)85

Interaction indices
Reversed-phase LC of homologous series. A general method for prediction of retention 314(1984)13

Interfaces, GC-MS
Optimizing flow-rates in capillary GC-MS 325(1985)23

Interfaces, LC-MS
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Ten years of LC-MS 323(1985)3

Determination of dibenzothiophene in oils by LC-tandem MS 323(1985)33

Open-tubular LC-MS using direct liquid introduction 323(1985)37

Performance of a moving belt LC-MS interface 323(1985)81

Micro column LC-MS using a capillary interface 323(1985)89

Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Design and construction of an interface for direct liquid introduction coupling of a microbore high-performance liquid chromatograph to a quadrupole mass spectrometer 323(1985)121

Studies with a laboratory-constructed thermospray liquid chromatograph-mass spectrometer interface 323(1985)127

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Interfacial potentials
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433

Interfacial tension
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79

Interferon
High-performance immunosorbent purification of recombinant leukocyte A interferon 303(1984)225

Internal standards
Separation and use of the perdeuterated analogue as an internal standard for the analysis of ethylene dibromide 312(1984)450

Inverse injection system
"Inverse" injection system for GLC with capillary columns 322(1985)206

Iodide
Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Iodinated peptides
Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Iodine
GC identification of alkyl radicals formed in plasma radiofrequency discharges by using iodine as a scavenger 303(1984)61

Microdetermination of iodine in organic compounds by reversed-phase HPLC 324(1985)498

Iodine values
On-line GC evaluation of iodine values in edible fats and oils 314(1984)233

Iodothyroxine
Soluble antibody affinity chromatography technique investigated with ultratrace [^{125}I]thyroxine 314(1984)295

Ion chromatography
Recent developments in ion chromatography (Review) 313(1984)47

Ion-exchange desolvation mechanism
Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

Ion exchangers
Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

Ion-exchange properties of activated carbon filled with hydrous Zr(IV) and Zr(IV)-P(V) oxides 303(1984)13

Recovery of plant amines with carboxylic ion-exchange substrates and their subsequent analysis by TLC 312(1984)357

Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

Use of cation exchangers for the on-line preconcentration of polar anilines in LC 317(1984)557

GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

Ionic charge
Determination of ionic charge by liquid-liquid partition 322(1985)425

Ionization quenching
Ionization quenching of a hydrogen atmosphere flame as a detection method for GC 303(1984)331

Ion-pairing reagents
Selection of mixed ion-pair modifiers for HPLC mobile phases 314(1984)402

Rapid optimization of the concentration of the ion-pairing reagent in ion-pairing reversed-phase LC 316(1984)231

Iothalamate
Simple and micro HPLC method for simultaneous determination of *p*-aminohippuric acid and iothalamate in biological fluids 306(1984)89

Iron
HPLC of homo- and hetero-tetrametallic iron, ruthenium, osmium and nickel clusters 303(1984)404

Iron-alumina catalysts
Effect of activated sorption processes on the results of thermoprogrammed reduction investigations on Fe-Al_2O_3 catalysts 301(1984)270

Isocolchicine
Separation of colchicine and related hydrolysis and photodecomposition products by HPLC, using copper ion complexation 315(1984)233

Isocyanates
Separation of isocyanate prepolymer components as their urea derivatives by reversed-phase HPLC 301(1984)485

Isodesmosine
Separation of the cross-linking amino acids of elastin on thin-layer plates 305(1984)461

Isoniazid
Determination of isoniazid and acetylisoniazid in plasma or serum by HPLC (in French) 307(1984)137

Isoprenaline
Non-radiochemical procedure for the measurement of O-methylation of the stereoisomers of isoprenaline 310(1984)283

Isoprenoid alcohols
Investigation of isoprenoid benzoates and naphthoates by reversed-phase LC. Isocratic elution characteristics of benzoates and naphthoates of C_5-C_{20} terpenoid alcohols 312(1984)121

Isoprenoid esters
GC and GC-MS analysis of diastereomeric acyclic isoprenoid esters using fused-silica capillary columns 312(1984)395

N-Isopropyl-N'-phenyl-*p*-phenylene-diamine
Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141

Isoproterenol sulphate
Determination of *d*-isoproterenol sulphate by HPLC with amperometric detection 306(1984)257

Isoproturon
Reversed-phase HPLC method for the simultaneous microquantitative determination of the herbicide isoproturon and its possible degradation products in soil 318(1985)144

Isoquinolines
GC analysis of endogenous catecholamines, phenolic amines and derived isoquinolines using short glass capillary columns and electron-capture detection 311(1984)17

On-column UV absorption detector for open tubular capillary zone electrophoresis 315(1984)135

Isosorbide dinitrate
Determination of the two mononitrate metabolites of isosorbide dinitrate in human plasma and urine by GC with electron-capture detection 305(1984)95

Determination of isosorbide dinitrate and its mononitrate metabolites in human plasma using Extrelut purification and capillary column GLC 306(1984)165

Determination of the partitioning, stability, and metabolite formation of isosorbide dinitrate in human and rat blood using and improved GLC assay 308(1984)153

Isotachophoresis
Isotachophoresis in inorganic chemistry (Review) 313(1984)189

Unified view of moving boundary electrophoresis: practical implications (plenary lecture) 320(1985)1

Isotachophoresis sensitivity
Enhancing isotachophoresis sensitivity by low-concentration electrolyte cascading 320(1985)45

Isotachophoretic dynamics
Dynamics of isotachophoretic separation. I. Computer simulation 320(1985)167

Dynamics of isotachophoretic separation. II. Impurities in the electrolytes and sample injection 320(1985)175

Isotachophoretic index
Isotachophoretic determination of mobility and pK_a by means of computer simulation. III. Evaluation of mobility and pK_a of fifteen nucleotides and seven phosphorus oxoacids and their isotachophoretic separation 318(1985)195

Isotopically labelled compounds
Separation and use of the perdeuterated analogue as an internal standard for the analysis of ethylene dibromide 312(1984)450

Isoxepac
Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133

Isoxicam
HPLC analysis of isoxicam in human plasma and urine 305(1984)145

J

Juvenile hormone
Convenient method for the determination of picomole amounts of juvenile hormone 323(1985)355

K

Kaolinite
Characterization of adsorption at kaolinite surfaces by heterogeneous GSC 314(1984)111

Keratin
Reversed-phase HPLC of some ultra-heterogeneous and covalently-modified proteins from human hair 324(1985)65

Kerosine fuel
HPLC procedure for direct determination of the compound class composition of diesel and kerosine fuels using refractive index detection 312(1984)237

Keto acids
Characterization of artefacts produced by treatment of organic acids with diazomethane 309(1984)243

Branched-chain α-keto acid analysis in biological fluids: preparative clean-up by anion-exchange and analysis by capillary GC 310(1984)1

Ketoconazole
Micro-determination of ketoconazole in plasma or serum by HPLC 306(1984)436

Keto-D-gluconic acid
Determination of D-gluconic, 5-keto-D-gluconic, 2-keto-D-gluconic and 2,5-diketo-D-gluconic acids by HPLC 312(1984)211

Ketones
Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23

TLC of enamino ketones. II 312(1984)387

HPLC of carbonyl compounds as 2,4-dinitrophenylhydrazones with electrochemical detection 318(1985)427

Determination of carbonyl compounds in an Australian (Rundle) shale oil 319(1985)230

HPLC elution behaviour of oligosaccharides, monosaccharides and sugar degradation products on series-connected ion-exchange resin columns using water as the mobile phase 322(1985)411

Structure-retention studies of model ketones by capillary GC 325(1985)13

6-Keto-prostaglandin
Determination of 6-keto-prostaglandin $F_{1\alpha}$ and thromboxane B_2 in urine by HPLC and radioimmunoassay 306(1984)351

17-Ketosteroid sulphates
Studies on steroids. CC. Determination of 17-ketosteroid sulphates in serum by HPLC with electrochemical detection using pre-column derivatization 307(1984)23

Ketotrilostane
Determination of trilostane and ketotrilostane in human plasma by HPLC 311(1984)434

Khellin
Rapid separation and quantitative determination of khellin and visnagin in *Ammi visnaga* (L.) Lam. fruits by HPLC 301(1984)297

Kinins
Detection and quantitation of fluorescamine-labeled bradykinin, its analogues and metabolites using HPLC 306(1984)109

Krebs cycle acids
Separation of tricarboxylic acid cycle acids and other related organic acids in insect haemolymph by HPLC 318(1985)112

L

Labetalol
HPLC method for the determination of labetalol in plasma using UV detection 305(1984)222

HPLC assay for labetalol in human plasma using a PRP-1 column and fluorometric detection 311(1984)319

Laccase
Determination of laccase activity with various aromatic substrates by HPLC 320(1985)435

Laccol
Analysis of long-chain phenols in the sap of the Burmese lac tree, *Melanorrhoea usitate* by capillary GLC 318(1985)378

Lactam antibiotics
Process scale-up of a β-lactam antibiotic purification by HPLC 316(1984)133

Lactate
HPLC determination of plasma lactate specific radioactivity 307(1984)380

Lactoglobulin
Preparative HPLC of proteins on ion-exchange columns with carboxymethyldextrans as displacers 316(1984)125

Lactones
Analysis of uronic and aldonic acids, their lactones, and related compounds by HPLC on cation-exchange resins 319(1985)159

Lactulose
Automated anion-exchange chromatographic method for the quantitation of lactulose in physiological fluids and processed milks 307(1984)393

Lantanosides
Investigation of conditions of the preparative LC separation of cardiac glycosides present in *Digitalis lanata* 303(1984)179

Lanthanoids
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11

Lasiocarpic acid
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

Lauric acid
Electrokinetic detection of lauric, myristic, stearic and arachidic acids separated by reversed-phase HPLC 321(1985)433

(*R*)-N-Lauroyl-α-(1-naphthyl)-ethylamine
Resolution of α-halogenocarboxylic acids by GC on (*R*)-N-lauroyl-α-(1-naphthyl)ethylamine 318(1985)235

Lectins
Rapid resolution of nucleotide sugars by lectin affinity HPLC 323(1985)434

Leucine polymers
Separation of hydrophobic peptide polymers by size-exclusion and reversed-phase HPLC 317(1984)1

Leukotrienes
Preparation of pentafluorobenzyl esters of arachidonic acid lipoxygenase metabolites. Analysis by GC and negative-ion chemical ionization MS 305(1984)3

Simple single-step HPLC method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 308(1984)316

Determination of leukotrienes and prostaglandins in [^{14}C]arachidonic acid labelled human lung tissue by HPLC and radioimmunoassay 311(1984)39

Levorphanol
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Lidocaine
Determination of lidocaine and active metabolites in blood serum by LC with electrochemical detection 306(1984)269

Simultaneous determination of lidocaine and its deethylated metabolites using GLC with nitrogen-phosphorus detection 307(1984)200

Simultaneous determination of lidocaine and its metabolites in plasma and myocardium 311(1984)291

HPLC determination of *d*-/*l*-epinephrine enantiomer ratio in lidocaine-epinephrine local anesthetics 325(1985)249

Ligand density
Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157

Ligand immobilization
Immobilization of ligands for affinity chromatography. Coupling on a spacer arm gel with N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline as condensation agent: study of coupling conditions by means of a radioimmunologic method 314(1984)445

Ligand-spacer coupling
Immobilization of ligands for affinity chromatography. Coupling on a spacer arm gel with N-ethoxycarbonyl-2-ethoxy-1,2-dihydroquinoline as condensation agent: study of coupling conditions by means of a radioimmunologic method 314(1984)445

Light-scattering detector
Examination of the concentration response of evaporative light-scattering mass detectors 323(1985)297

Evaluation of the mass detector for quantitative detection of triglycerides and fatty acid methyl esters 324(1985)35

Lignan lactones
Solid-state fluorodensitometric quantitation of arylnaphthalene lignan lactones of *Cleistanthus collinus* 303(1984)296

Lignins
Aqueous size exclusion chromatography of industrial lignins 322(1985)129

Linalool
Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23

Linear free energy relationships
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47

Partitioning of oligoglucans in Sephadex G-15 in relation to their conformational structure 318(1985)57

Chromatographic selectivity in LC of polycondensed aromatic hydrocarbons on 3-(2,4-Dinitroanilino)propyl silica 324(1985)261

Linogliride
Determination of linogliride in biological fluids by HPLC 311(1984)418

Lipid A
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Lipids
Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173

Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65

Quantitative densitometric method for the rapid separation and quantitation of the major lipids of tissues and lipoproteins by HPTLC. II. Reduction of the densitometric data 307(1984)81

Capillary GC investigation of plasma lipid alcoholysis during alcohol extraction 307(1984)172

Faecal lipid chromatography. I. Quantitative determination with Chromarods 310(1984)297

Enhanced response of more highly unsaturated lipids on TLC-Iatroscan flame ionization detection after exposing the developed Chromarods to iodine vapour 319(1985)215

Improved one-dimensional TLC technique for polar lipids 322(1985)391

Simplified procedure for the quantitation of radioactive phosphoinositides by TLC 323(1985)462

Lipophilic index
Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. II. Procedure for the determination of a lipophilic index measured by reversed-phase HPLC 320(1985)305

Lipophilicity
see also Hydrophobicity

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. I. Relationship between capacity factors and the methanol concentration in methanol-water eluents 320(1985)293

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. II. Procedure for the determination of a lipophilic index measured by reversed-phase HPLC 320(1985)305

Lipopolysaccharides
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Lipoproteins
Detection of an abnormal lipoprotein in a large colony of pedigreed baboons using high-performance gel exclusion chromatography 308(1984)101

Resolution and quantitation of apolipoproteins A-I and A-II from human high-density lipoprotein by size exclusion HPLC 311(1984)79

Analytical capillary isotachophoresis: a routine technique for the analysis of lipoproteins and lipoprotein subfractions in whole serum 320(1985)253

Lipsticks
Determination of synthetic organic colours in lipsticks by TLC and HPLC 318(1985)149

Liquid chromatography-mass spectrometry
Ten years of LC-MS 323(1985)3

Chromatographic perspective of HPLC-MS 323(1985)13

Determination of dibenzothiophene in oils by LC-tandem MS 323(1985)33

Open-tubular LC-MS using direct liquid introduction 323(1985)37

Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

HPLC-MS of derivatized and underivatized amino acids 323(1985)67

Performance of a moving belt LC-MS interface 323(1985)81

Micro column LC-MS using a capillary interface 323(1985)89

Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

On-line trace enrichment for improved sensitivity in LC with direct liquid introduction MS detection 323(1985)113

Design and construction of an interface for direct liquid introduction coupling of a microbore high-performance liquid chromatograph to a quadrupole mass spectrometer 323(1985)121

Studies with a laboratory-constructed thermospray liquid chromatograph-mass spectrometer interface 323(1985)127

On-line LC-MS analysis of non-ionic surfactants 323(1985)135

Qualitative and quantitative analysis of ranitidine and its metabolites by HPLC-MS 323(1985)143

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Applications of thermospray ionisation LC-MS to compounds of pharmaceutical interest 323(1985)163

Liquid chromatography-mass spectrometry coupling
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Liquid crystals
Liquid crystalline stationary phases with a cyclohexyl ring in the molecule 312(1984)141

Disc-like liquid crystalline stationary phases from the triphenylene derivatives group 315(1984)145

Aryl substituted diastereomeric alkenes: GC behavior on a non-polar *versus* a liquid crystal phase 321(1985)127

High-performance GC on liquid crystal capillaries. VIII. Separation of alkylbenzenes up to C_9 (in German) 323(1985)414

Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

Distribution of the liquid crystal stationary phase on the surface of diatomite supports 324(1985)299

Phase transitions of 4-cyano-4'-*n*-alkoxyformyloxyazobenzenes on silanized supports 324(1985)309

Lofemizole
GLC evaluation of lofemizole in biological samples for pharmacokinetic investigations. Comparison of two analytical methods 310(1984)51

Lolitrem B
Rapid determination of the neurotoxin lolitrem B in perennial ryegrass by HPLC with fluorescence detection 321(1985)217

HPLC with stop-flow UV spectral characterization of lolitrem neurotoxins from perennial ryegrass 322(1985)159

London force interaction
GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19

Loop diuretics
HPLC behaviour of some pharmaceutically important thiazide, loop and potassium-sparing diuretics 325(1985)395

Lorazepam
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Lubricants
Analysis of mechanical lubricant contaminants in aluminium cold rolling oil by reversed-phase LC 319(1985)222

Luciferase
Metal-ion mediated interaction of luciferase with tetraiodofluorescein 315(1984)175

Lumicolchicine
Separation of colchicine and related hydrolysis and photodecomposition products by HPLC, using copper ion complexation 315(1984)233

Luteinizing hormone releasing hormone
Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Luteose
Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Luteotropic hormone
Subunits of bovine lutropin. Hormonal and immunological activity after separation with reversed-phase HPLC 319(1985)195

Lymphocytes
Hydrophobic chromatography of cells: adsorption and resolution on homologous series of alkylagaroses 317(1984)103

Lysine
Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

Isotachophoretic determination of lysine in fermentation broths 320(1985)131

Lysinoalanine
Determination of lysinoalanine as the heptafluorobutyryl isobutyl ester derivative by GLC 303(1984)234

Lysinonorleucine
Separation of the cross-linking amino acids of elastin on thin-layer plates 305(1984)461

Lysodren
The mass spectrometer as a chlorine-selective chromatographic detector. II. Applications to chemical systems 314(1984)93

Lysozyme
Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

On-column UV absorption detector for open tubular capillary zone electrophoresis 315(1984)135

Kinetics of unfolding of proteins on hydrophobic surfaces in reversed-phase LC 317(1984)227

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

M

α_2-Macroglobulin
Fractionation of human serum proteins by immobilized metal affinity chromatography 315(1984)167

Magnesium
Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Major peak elimination
Elimination of the peak of a major ion in anion chromatography with UV detection 325(1985)327

Maprotiline
Fluorimetric determination of maprotiline in urine and plasma after TLC separation 305(1984)508

Measurement of maprotiline and oxaprotiline in plasma by HPLC of fluorescent derivatives 309(1984)107

Marine diesel fuel
Supercritical fluid chromatography and supercritical fluid chromatography-MS of marine diesel fuel 314(1984)253

Mass detector
Examination of the concentration response of evaporative light-scattering mass detectors 323(1985)297

Evaluation of the mass detector for quantitative detection of triglycerides and fatty acid methyl esters 324(1985)35

Mass-flow-modulated detector
Design and performance of a mas-flow-modulated detector for GC 319(1985)263

Mass spectrometer
The mass spectrometer as a chlorine-selective chromatographic detector. I. Description and evaluation of the technique 314(1984)83

The mass spectrometer as a chlorine-selective chromatographic detector. II. Applications to chemical systems 314(1984)93

MCPA
Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

McReynolds probes
Indirect determination of specific retention volumes in GC. Use of retentions of the McReynolds probes relative to *n*-decane 312(1984)69

MDL 17,043
Simultaneous analysis of a new cardiotonic agent, MDL 17,043, and its major sulfoxide metabolite in plasma by HPLC 306(1984)249

Mechanical lubricants
Analysis of mechanical lubricant contaminants in aluminium cold rolling oil by reversed-phase LC 319(1985)222

Mecoprop
Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Mefloquine
Determination of the antimalarial mefloquine in humans plasma by GC with electron-capture detection 311(1984)411

Meibomian gland lipids
Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173

α-Melanotropin
HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99

Reversed-phase HPLC studies of α-MSH fragments 314(1984)313

Melatonin
Analysis of melatonin, 5-methoxytryptophol and 5-methoxyindoleacetic acid in the pineal gland and retina of hamster by capillary column GC-MS 311(1984)1

Use of impregnated reagents in the pentafluorobenzylation of indoleamine metabolites 325(1985)309

Membrane proteins
Isolation and purification of HLA-DR antigens from tumour cells by affinity chromatography and chromatofocusing 308(1984)322

Analysis of membrane proteins by ion-exchange HPLC 322(1985)212

Fractionation of thylakoid membrane components by extraction in aqueous polymer two-phase systems containing detergent 323(1985)363

Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173

Membranes
Liquid-liquid extraction of membranes from calf brain using conventional and centrifugal counter-current distribution techniques 311(1984)277

Affinity chromatography of membrane fragments. Separation of carbonic anhydrase-enriched and -depleted brain membranes 312(1984)435

Menadione
Fluorimetric determination of menadione sodium bisulphite (vitamin K_3) in animal feed and premixes by HPLC with post-column derivatization 301(1984)441

Menthol
HPLC assay of menthol using indirect photometric detection 303(1984)436

Menthyl esters
GC and GC-MS analysis of diastereomeric acyclic isoprenoid esters using fused-silica capillary columns 312(1984)395

Mephenytoin
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Direct enantiomeric resolution of mephenytoin and its N-demethylated metabolite in plasma and blood using chiral capillary GC 307(1984)121

Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444

Mepixanthone
HPLC determination of mepixanthone in serum 309(1984)214

Mequitazin
Determination of mequitazin in human plasma and urine by capillary column GLC-MS 309(1984)391

Mercaptans
GC study of the analysis and elution mechanism of hydrogen sulphide, carbonyl sulphide and light mercaptans in petroleum gases 321(1985)59

2-Mercaptoethanesulphonate
Determination of sodium 2-mercaptoethanesulphonate by HPLC using post-column reaction colorimetry or electrochemical detection 311(1984)234

6-Mercaptopurine
Novel method for the quantitation of 6-mercaptopurine in human plasma using HPLC with fluorescence detection 309(1984)409

Merodesmosine
Separation of the cross-linking amino acids of elastin on thin-layer plates 305(1984)461

Mesaconitine
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Metabolites
Advantages and limitations of isotachophoretic determinations of metabolites in tissue extracts 320(1985)105

Metal carbonyls
Inorganic GC (Review) 313(1984)3

Metal cations
Isotachophoresis in inorganic chemistry (Review) 313(1984)189

Metal chelates
Inorganic GC (Review) 313(1984)3

Determination of molybdenum and tungsten with tiron by reversed-phase LC 320(1985)403

Volatility studies of metal chelates. VI. GC behaviour of aryl β-diketones and their metal chelates 322(1985)1

Volatility studies of metal chelates. VII. GC behaviour of chelates derived from the aromatic β-diketones 1-phenylbutane-1,3-dione and 4,4,4-trifluoro-1-(pentafluorophenyl)butane-1,3-dione 322(1985)23

Metal complexes
Substitution-inert metal complex mobile phases in non-suppressed anion chromatography 312(1984)99

Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59

Reversed-phase high-performance ion-pair chromatography of cobalt(III) coordination complexes (Review) 313(1984)93

Chromatographic study of optical resolution. XI. Optimum conditions for optical resolution of uncharged complexes 319(1985)35

Separation of anionic chloro complexes by capillary tube isotachophoresis in non-aqueous systems 322(1985)139

Metal β-diketonates
Improved elution of metal β-diketonates on fused-silica wall-coated open-tubular columns 324(1985)75

Metal ions
Simultaneous determination of anions and metal cations by single-column ion chromatography with ethylenediaminetetraacetate as eluent and conductivity and UV detection 312(1984)327

Recent developments in ion chromatography (Review) 313(1984)47

α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81

Metallocenes
Inorganic GC (Review) 313(1984)3

Metalloporphyrins
Non-aqueous LC of some metalloporphyrins on a reversed-phase column 315(1984)111

HPLC of dicarboxylic porphyrins and metalloporphyrins: retention behaviour and biomedical applications 317(1984)333

Metallothionein
Separation of metallothionein into isoforms by column switching on gel permeation and ion-exchange columns with HPLC-atomic-absorption spectrophotometry 303(1984)131

Metanephrine
Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Metanephrines
Selective retention of catecholamines and their derivatives in reversed-phase ion-pair partition chromatography 322(1985)43

Methacrylate copolymers
GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

Methacrylate monomers
Chromatography of monomers. II. Glass capillary GC of C_1-C_{18} alkyl esters of acrylic and methacrylic acids 325(1985)367

Methacycline
Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

Methamidophos
Improved analysis of acephate and methamidophos in biological samples by selective ion monitoring GC-MS 301(1984)465

Methamphetamine
Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Determination of methamphetamine and its metabolites in rat tissues by GC with a nitrogen-phosphorus detector 318(1985)307

Methanol
Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79

Quantitation of methanol formed in cell culture cytotoxicity assays and as a metabolite in microsome suspensions 305(1984)171

Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.*-butanol 319(1985)247

Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

Methohexital
HPLC method for methohexital and thiopental in plasma or whole blood 307(1984)481

Methoxyanilines
Separation of methoxyanilines by competing-ion reversed-phase HPLC 316(1984)227

6-Methoxy-8-(6-diethylaminohexylamino)-4-methylquinoline
HPLC method for the analysis of a candidate 8-aminoquinoline antileishmanial drug using oxidative electrochemical detection 311(1984)117

Methoxy-5-hydroxyflavones
TLC behaviour and chemical structure of 6- and 8-methoxy-5-hydroxyflavones 315(1984)101

3-Methoxy-4-hydroxyphenylacetic acid
Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

3-Methoxy-4-hydroxyphenylethylene glycol
Direct determination of 3-methoxy-4-sulphonyloxyphenylglycol (MHPG sulphate) in urine using GLC 305(1984)438

HPLC method for determining plasma and urine 3-methoxy-4-hydroxyphenylglycol by amperometric detection 306(1984)358

Measurement of 3-methoxy-4-hydroxyphenylglycol sulfate ester in brain using reversed-phase LC and electrochemical detection 306(1984)364

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

Determination of free 3-methoxy-4-hydroxyphenylglycol with several other monoamine metabolites in plasma by HPLC with amperometric detection 311(1984)51

Isocratic analysis of 3-methoxy-4-hydroxyphenyl glycol, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in cerebrospinal fluid by HPLC with amperometric detection 311(1984)354

Method for total 3-methoxy-4-hydroxyphenylglycol extraction from urine, plasma and brain tissue using bonded-phase materials: comparison with the ethyl acetate extraction method 311(1984)361

Determination of urinary 3-methoxy-4-hydroxyphenylglycol by LC with electrochemical detection 317(1984)327

Selective retention of catecholamines and their derivatives in reversed-phase ion-pair partition chromatography 322(1985)43

5-Methoxyindoleacetic acid
Analysis of melatonin, 5-methoxytryptophol and 5-methoxyindoleacetic acid in the pineal gland and retina of hamster by capillary column GC-MS 311(1984)1

3-Methoxyisoprenaline
Non-radiochemical procedure for the measurement of O-methylation of the stereoisomers of isoprenaline 310(1984)283

Methoxyphenamine
GC quantitation of methoxyphenamine and three of its metabolites in plasma 310(1984)307

3-Methoxy-4-sulphonyloxyphenyl-ethylene glycol
Direct determination of 3-methoxy-4-sulphonyloxyphenylglycol (MHPG sulphate) in urine using GLC 305(1984)438

5-Methoxytryptophol
Analysis of melatonin, 5-methoxytryptophol and 5-methoxyindoleacetic acid in the pineal gland and retina of hamster by capillary column GC-MS 311(1984)1

3-Methoxytyramine
Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Methylalkanes
Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333

Methylamine
Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Methylamino acids
Phosgene: a versatile reagent for enantiomer separation by capillary GC 301(1984)129

Determination of free trimethyllysine in plasma and tissue specimens by HPLC 306(1984)79

GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

Rapid ion-exchange method for the determination of 3-methylhistidine in rat urine and skeletal muscle 311(1984)69

Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

α-Methyl-α-amino acids
Enantiomeric separation of underivatized α-methyl-α-amino acids by HPLC 318(1985)117

9-(N-Methylaminomethyl)anthracene
Improved chromatographic procedure for determination of 9-(N-methylaminomethyl)anthracene isocyanate derivatives by HPLC 301(1984)481

Methylated nucleobases
Improved fluorometric method for quantitative HPLC analysis of methylated guanine derivatives in DNA 324(1985)484

Separation of naturally occurring or induced methylated nucleobases of DNA by reversed-phase HPLC 325(1985)207

9β-Methylcarbacyclin
HPLC method for the separation of isomers of 9β-methylcarbacyclin (ciprostene) and the analysis of pharmaceutical samples 321(1985)209

Methyl carboprost
HPLC determination of acid-catalyzed degradation products of methyl carboprost in a polymeric controlled-release device 314(1984)369

2-Methylcardol
Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

Methylcotinium
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291

***S*-Methylcysteine sulphoxide**
Determination of *S*-methyl cysteine sulfoxide in *Brassica* extracts by HPLC 319(1985)450

Methylcytosine
GC-MS characteristics of methylated perfluoroacyl derivatives of cytosine and 5-methylcytosine 314(1984)211

Separation of naturally occurring or induced methylated nucleobases of DNA by reversed-phase HPLC 325(1985)207

N^6-Methyldeoxyadenosine
Quantitative reversed-phase HPLC of major and modified nucleosides in DNA 301(1984)199

5-Methyldeoxycytidine
Quantitative reversed-phase HPLC of major and modified nucleosides in DNA 301(1984)199

Methyldopa
Sensitive HPLC assay using electrochemical detection for a novel prodrug ester of methyldopa, pivaloyloxyethyl 3-(3,4-

dihydroxyphenyl)-2-methylalaninate, in plasma and urine 308(1984)251

HPLC determination of plasma catecholamines during α-methyldopa therapy 308(1984)301

Methylephedrine
Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

Methylguanidine
HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257

Methylguanine
Improved fluorometric method for quantitative HPLC analysis of methylated guanine derivatives in DNA 324(1985)484

Methylhistamine
Reversed-phase ion-pair HPLC procedure for determination of histamine and its metabolites in rat urine 312(1984)443

Methylhistidine
GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

Rapid ion-exchange method for the determination of 3-methylhistidine in rat urine and skeletal muscle 311(1984)69

Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

1-Methylimidazole-4-acetic acid
Sensitive GLC procedure for urinary Nτ-methylimidazole acetic acid, an index of histamine turnover 306(1984)69

Methylimidazoleacetic acid
Reversed-phase ion-pair HPLC procedure for determination of histamine and its metabolites in rat urine 312(1984)443

3-Methylindole
GC determination and MS confirmation of traces of indole and 3-methylindole (skatole) in pig back fat 324(1985)469

Methyl ketononanoates
Use of retention time plots as a means of identifying isomeric methyl ketononanoates 324(1985)440

Methylnoradrenaline
HPLC determination of plasma catecholamines during α-methyldopa therapy 308(1984)301

Methyl oleate
Enhanced response of more highly unsaturated lipids on TLC-Iatroscan flame ionization detection after exposing the developed Chromarods to iodine vapour 319(1985)215

Methylprednisolone
Analysis of cortisol, methylprednisolone, and methylprednisolone hemisuccinate. Absence of effects of troleandomycin on ester hydrolysis 305(1984)271

Determination of methylprednisolone and methylprednisolone acetate in synovial fluid using HPLC 309(1984)385

Methylpseudoephedrine
Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

0-Methyltransferases
HPLC of proteins. Purification of plant enzymes by ion-exchange chromatography 315(1984)333

N-Methyltryptamine
HPLC determination of hallucinogenic indoleamines with simultaneous UV photometric and voltammetric detection 320(1985)414

Methyl ureas
Analysis of selected pesticides by HPLC-MS 312(1984)221

3-Methylxanthine
Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107

Metocurine iodide
Determination of D-tubocurarine chloride or metocurine iodide in human plasma by HPLC with UV detection 306(1984)398

Metoprolol
Determination of metoprolol and its α-hydroxylated metabolite in human plasma by HPLC 305(1984)411

Determination of two metoprolol metabolites in human urine by HPLC 309(1984)198

Simultaneous determination of metoprolol and metabolites in urine by capillary column GC as oxazolidineone and trimethylsilyl derivatives 309(1984)317

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Metronidazole
Effect of precipitating agents on the analysis of metronidazole by HPLC 303(1984)222

Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325

Rapid determination of metronidazole in human serum and urine using a normal-phase HPLC column with aqueous solvents 309(1984)436

Metropolol
Influence of solute polarity in column-switching chromatography for the assay of drugs in plasma and urine 317(1984)493

Metroprolol
Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

Mexiletine
HPLC method for the measurement of mexiletine and flecainide in blood plasma or serum 307(1984)439

GC method for the routine serum monitoring of mexiletine 309(1984)165

Mianserin
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Micellar chromatography
Gradient elution micellar LC 316(1984)183

Microbore columns
see also Capillary columns and columns

Preparation of efficient and stable reversed-phase microbore columns for HPLC 321(1985)450

Micronomicin
Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373

Minicolumns
Protein separations on reversed-phase HPLC minicolumns 317(1984)119

Misonidazole
Determination of misonidazole and desmethylmisonidazole in plasma by HPLC with reductive electrochemical detection 310(1984)159

Mitomycin C
LC determination of mitomycin C in human plasma and urine 306(1984)444

LC determination of mitomycin C in plasma and urine 306(1984)446

Mitotane
Mitotane (*o,p'*-DDD) emulsion and tablet analysis by HPLC 319(1985)467

Mixed ligand complexes
Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161

Mixed mode retention
Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157

Mobile phase composition
Basis of retention in normal-phase HPLC with cyano-propyl columns 303(1984)1

Computer-assisted optimization of eluent concentration and pH in ion chromatography 303(1984)321

Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Substitution-inert metal complex mobile phases in non-suppressed anion chromatography 312(1984)99

Automated optimisation of reversed-phase HPLC separations. An improved method using the sequential simplex procedure 316(1984)69

Influence of mobile phase additives in LC of phenoxypropanolamines 316(1984)291

Role of mobile phase components for multi-functional solutes in silica gel LSC. Resolution of protected deoxyribonucleotide derivatives for oligomer synthesis 316(1984)389

Ammonium acetate: a general purpose buffer for clinical applications of HPLC 316(1984)397

Mobile phase effects in reversed-phase chromatography. VII. Dependence of retention on mobile phase composition and column temperature 318(1985)1

Reversed-phase chromatography of proteins with diphenyl-silica column and hydro-organic eluents containing two organic solvents 318(1985)11

Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. I. Relationship between capacity factors and the methanol concentration in methanol-water eluents 320(1985)293

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. II. Procedure for the determination of a lipophilic index measured by reversed-phase HPLC 320(1985)305

Chromatography with sub- and supercritical eluents: influence of temperature, pressure and flow-rate on the behaviour of lower alkanes 321(1985)273

Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. I. Sorption isotherms of *n*-alcohols on octyl-agarose 322(1985)399

HPLC analysis of basic drugs on silica columns using non-aqueous ionic eluents. I. Factors influencing retention, peak shape and detector response 323(1985)173

Mobile phase optimization for multi-functional solutes in LSC. Resolution of synthetic intermediates of oligodeoxyribonucleotides 323(1985)227

Mobile phase optimization for the reversed-phase LC of complex hydroxyl aromatic mixtures 324(1985)1

The use of retention indices to measure the solvent selectivity of ternary eluents in reversed-phase LC 324(1985)243

Mobilities
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11

Isotachophoretic determination of mobility and pK_a by means of computer simulation. III. Evaluation of mobility and pK_a of fifteen nucleotides and seven phosphorus oxoacids and their isotachophoretic separation 318(1985)195

Determination of relative ionic mobilities by capillary isotachophoresis 320(1985)139

Determination of absolute mobilities and pK values by isotachophoresis 320(1985)147

Molecular orbital calculations
Study of GC behaviour of alkenes based on molecular orbital calculations 302(1984)107

Molecular topology
Calculation of retention indices by molecular topology: chlorinated alkanes 314(1984)1

Molecular weight determination
Molecular weight determination of polychromophore polymers using HPLC and size-exclusion chromatography with UV detection 322(1985)338

Further experiments on a new, fast method for determining molecular weights of diffusing species in a liquid phase 322(1985)443

Experimental evaluation of the errors in high-performance GPC measurements 325(1985)37

Determination of polymer molecular weights by PC 325(1985)363

Molten salts
Influence of phase loading on the performance of whisker-walled open tubular columns coated with organic molten salts 324(1985)415

Molybdate
Capillary-tube isotachophoretic separation of molybdate VI, Tungstate VI and Vanadate V ions using ion pairing equilibria 325(1985)179

Molybdenum
GC studies on group VIB organometallic complexes. The analytical and preparative separation of chromium, molybdenum and tungsten trifluorophosphine carbonyl complexes 301(1984)253

Determination of molybdenum and tungsten with tiron by reversed-phase LC 320(1985)403

Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Monoacetyldapsone
Quantification of antimalarial drugs. II. Simultaneous measurement of dapsone, monoacetyldapsone and pyrimethamine in human plasma 307(1984)426

Monoalkyltin
Speciation of alkyltin compounds by HPLC with a cyanopropyl-bonded silica column 315(1984)351

Monoamine oxidase
Determination of monoamine oxidase B activity by HPLC 309(1984)156

Monochlorobenzoates
GLC analyses. XXXVII. Capillary column studies of benzoyl and monochlorobenzoyl esters of lower saturated branched-chain alcohols 323(1985)331

GLC Analyses. XXXVIII. Capillary column studies of unsaturated esters of benzoic and monochlorobenzoic acids 324(1985)113

Monochlorobenzoic acid
GLC analyses. XXXV. Capillary column studies of C_1-C_{12} n-alkyl esters of benzoic and monochlorobenzoic acids 322(1985)83

Monochlorobutanoates
GC of homologous esters. XXXI. Butanoyl and monochlorobutanoyl esters of lower saturated branched chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 325(1985)61

Monochloropropanoyl esters
GC of homologous esters. XXIX. Propanoyl and monochlorpropanoyl esters of lower saturated brached-chain and unsaturated alcohols 324(1985)343

Monoclonal antibodies
One-step procedure for the rapid isolation of mouse monoclonal antibodies and their antigen binding fragments by fast protein LC on a Mono Q anion-exchange column 319(1985)67

Micropreparative isotachophoretic electrodesorption of monoclonal antibodies from an affinity adsorbent 320(1985)81

Monoethylglycine xylidide
Determination of lidocaine and active metabolites in blood serum by LC with electrochemical detection 306(1984)269

Monoethylglycylxylidide
Simultaneous determination of lidocaine and its deethylated metabolites using GLC with nitrogen-phosphorus detection 307(1984)200

Simultaneous determination of lidocaine and its metabolites in plasma and myocardium 311(1984)291

Monohydroxamate-Sepharose 6B
α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81

Separation of serum proteins on a Fe^{3+}-monohydroxamate adsorbent 321(1985)105

Monohydroxy fatty acids
Lipoxygenase products. A novel GC-MS assay for monohydroxy fatty acids 306(1984)1

Monolocalized solute retention
Influence of water in LSC systems on retention data. II. Fixation of monolocalized solutes with class N solvents and silica gels 316(1984)373

Monomethylalkanes
Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333

Monomethyllysine
GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

Monosaccharides
HPLC elution behaviour of oligosaccharides, monosaccharides and sugar degradation products on series-connected ion-exchange resin columns using water as the mobile phase 322(1985)411

Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

Monoterpenes
GLC study of monoterpene hydrocarbons of conifer balsams and ring opening products of pinane and carane structure 321(1985)240

Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

Morphine
Determination of morphine in biological samples by GC-MS. Evidence for persistent tissue binding in rats twenty-two days post-withdrawal 309(1984)73

Determination of morphine in serum and cerebrospinal fluid by GC and selected ion monitoring after reversed-phase column extraction 310(1984)193

Determination of pseudomorphine in morphine injection by HPLC 312(1984)473

Analysis of opium and crude morphine samples by capillary GC. Comparison of impurity profiles 315(1984)404

Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Motilin
Rapid high-yield purification of canine intestinal motilin and its complete sequence determination 321(1985)421

Moving-belt interface
Ten years of LC-MS 323(1985)3

Performance of a moving belt LC-MS interface 323(1985)81

Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Moving boundary electrophoresis
Unified view of moving boundary electrophoresis: practical implications (plenary lecture) 320(1985)1

Multi-channel detectors
Multi-channel detectors for micro HPLC: examination of flow cell structures 316(1984)441

Munitions components
Determination of munitions components in water by resin adsorption and HPLC electrochemical detection 302(1984)51

Muramic acid
Amperometric detection of muramic acid in HPLC with a post-column reaction 316(1984)495

Mutagenic artifacts
Production of mutagenic artifacts by the action of residual chlorine on XAD-4 resin 325(1985)95

Mutagenicity testing
Sorption of organic compounds from urine in mutagenicity testing: choice of sorbent 310(1984)41

Mycotoxins
Confirmation of nivalenol and deoxynivalenol by on-line LC-MS and GC-MS. Comparison of methods 318(1985)103

Optimisation of conditions for the trimethylsilylation of trichothecene mycotoxins 319(1985)376

Myelin basic protein
Simple and rapid identification of phosphorylated peptides from bovine brain myelin basic protein by reversed-phase HPLC 319(1985)359

Myoglobin
Myoglobin determination by HPLC 317(1984)87

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Myricetin
Isolation of quercetin, myricetin, and their respective dihydro-compounds by Sephadex LH-20 chromatography 324(1985)495

Myricoside
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Myristic acid
Determination of acids in whole lipopolysaccharide and in free lipid A from *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 308(1984)19

Electrokinetic detection of lauric, myristic, stearic and arachidic acids separated by reversed-phase HPLC 321(1985)433

N

NAD
LC separations of dimers of nicotinamide adenine dinucleotide and nicotinamide adenine dinucleotide phosphate 319(1985)341

Nadolol
Estimation of nadolol levels in plasma using HPLC with recirculating eluent flow 305(1984)489

Ion-exchange HPLC in drug assay in biological fluids. IV. Nadolol diastereomers: demonstration of pharmacokinetic and binding equivalence 309(1984)421

NADP
LC separations of dimers of nicotinamide adenine dinucleotide and nicotinamide adenine dinucleotide phosphate 319(1985)341

Nafazatrom
HPLC determination of nafazatrom in human plasma using fluorescence detection 305(1984)163

Nalbuphine
Detection of nalbuphine in plasma: an improved HPLC assay 311(1984)223

Naloxone
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

HPLC analysis of naloxone in human serum 311(1984)213

Naltrexone
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Naphthalenesulphonic acids
Spectrophotometric detection in capillary isotachophoresis: determination of the composition of mixed zones 320(1985)185

2-Naphthoic acid
Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

1-(1-Naphthyl)ethyl isocyanate
Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

1-Naphthyl isocyanate
Determination of aliphatic amines in air by reversed-phase HPLC using 1-naphthyl isocyanate derivatives 312(1984)482

1-Naphthyl polysiloxane
Polarizable polysiloxane stationary phases for capillary column GC 302(1984)303

Naphthylurethanes
Separation and determination of diacylglycerols as their naphthylurethanes by HPLC 307(1984)387

Narcotic analgesics
HPLC retention data for 84 basic drugs of forensic interest on a silica column using an aqueous methanol eluent 301(1984)165

Naringin
Micro column LC-MS using a capillary interface 323(1985)89

Nebramycin factors
LC separation of aminoglycosides with pulsed amperometric detection 324(1985)407

Nebulizers
Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Necatorin
Improved analytical and preparative methods for necatorin from *Lactarius necator* (Fr.) Karst mushroom 301(1984)303

Netilmicin
Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373

Neuraminic acids
Reversed-phase HPLC of hydrazones of 3-deoxy-D-*manno*-oct-2-ulosonic acid and neuraminic acids 315(1984)451

Neuropeptides
Ion-exchange chromatographic assay of peptidases acting on the C-terminal hexapeptide sequence of substance P 301(1984)415

HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99

Combined HPLC-radioimmunoassay for the characterization and quantitative measurement of neuropeptides 312(1984)273

Separation of the enkephalins from proteins in an aqueous medium by chromatofocusing 314(1984)499

Separation of enkephalins from their metabolites and precursor by a single pump gradient HPLC method 325(1985)467

Neurotensin
HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99

Neurotransmitters
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53

Nickel
HPLC of homo- and hetero-tetrametallic iron, ruthenium, osmium and nickel clusters 303(1984)404

Nickel chelates
Volatility and GC of some derivatives of Schiff base chelates 318(1985)350

Nickel iminodiacetate
Development of immobilized metal affinity chromatography. II. Interaction of amino acids with immobilized nickel iminodiacetate 323(1985)255

Development of immobilized metal affinity chromatography. III. Interaction of oligopeptides with immobilized nickel iminodiacetate 323(1985)265

Nicotine
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291

Nicotine-N'-oxide
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291

Nicotinic acid
Determination of nicotinic acid in serum by HPLC with fluorescence detection 309(1984)309

Nifedipine
LC determination of nifedipine in plasma and of its main metabolite in urine 308(1984)209

HPLC determination of nifedipine in plasma 310(1984)219

Niflumic acid
Purity assay of niflumic acid by reversed-phase HPLC 303(1984)440

Nikkomycins
Screening for new microbial products by HPLC using a photodiode array detector 316(1984)487

Nimodipine
GC and LC analyses of nimodipine calcium antagonist in blood plasma and cerebrospinal fluid 305(1984)105

Nitramines
Determination of munitions components in water by resin adsorption and HPLC electrochemical detection 302(1984)51

Nitrate
Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Sensitive determination of nitrite and nitrate by ion-exchange chromatography 322(1985)491

Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Nitrate esters
Determination of munitions components in water by resin adsorption and HPLC electrochemical detection 302(1984)51

Nitrazepam
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

HPLC determination of nitrazepam and its metabolites in human urine 310(1984)213

Nitrilotriacetic acid
Determination of nitrilotriacetic acid in water by high-resolution GC 312(1984)413

Nitrite
Determination of the stable isotope of nitrite flux in the blood of mice by GC-MS with selected ion monitoring 306(1984)51

Sensitive determination of nitrite and nitrate by ion-exchange chromatography 322(1985)491

Nitroaromatic compounds
On-line trace enrichment for improved sensitivity in LC with direct liquid introduction MS detection 323(1985)113

Nitrodiphenylamines
Influence of solute polarity in column-switching chromatography for the assay of drugs in plasma and urine 317(1984)493

Nitrogen-containing compounds
Steam GSC: a flexible separation technique 301(1984)11

Separation and identification by GC and GC-MS of the nitrogen compounds from a deasphalted heavy oil. Evolution of their distribution after a catalytic hydrotreatment (in French) 324(1985)87

Nitrogen heterocycles
LC systems for the separation of polycyclic aromatic hydrocarbon and nitrogen heterocycle compounds present in coal liquids 321(1985)343

Nitroglycerin
Application of HPLC with synchronized accumulating radioisotope detector to analysis of glyceryl trinitrate and its metabolites in rat plasma 305(1984)119

Rapid HPLC procedure for nitroglycerin and its degradation products 320(1985)445

Nitroimidazoles
Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325

Nitroimidazo[2,1-*b*]thiazoles
R_m values, retention times and π values of a series of potentially mutagenic nitroimidazo[2,1-*b*]thiazoles 320(1985)281

Nitro-organic compounds
Determination of munitions components in water by resin adsorption and HPLC electrochemical detection 302(1984)51

Nitro-polynuclear aromatic hydrocarbons
Detection of nitro-polycyclic aromatic hydrocarbons in LC by zinc reduction and peroxyoxalate chemiluminescence 316(1984)507

HPLC studies of environmental carcinogens in China 317(1984)545

GLC analyses. XXXIV. Separation and retention indices with retention increments of some nitrated polynuclear aromatic hydrocarbons on a low-polarity (SE-30) capillary column 322(1985)71

Nitrosamines
Capillary GC determination of volatile and tobacco-specific N-nitrosamines with the thermo-energy-analyzer (in German) 321(1985)475

Nitroso-alkylmethylamines
Retention behavior of long chain quaternary ammonium homologues and related nitroso-alkylmethylamines 324(1985)209

Nitrosourea nucleosides
Preparative and analytical HPLC methods in the synthesis and analysis of decomposition of nitrosourea nucleosides 322(1985)455

Nitrotoluenes
Determination of munitions components in water by resin adsorption and HPLC electrochemical detection 302(1984)51

Nivalenol
Confirmation of nivalenol and deoxynivalenol by on-line LC-MS and GC-MS. Comparison of methods 318(1985)103

Nolvadex
Applications of thermospray ionisation LC-MS to compounds of pharmaceutical interest 323(1985)163

Nonhistone chromosomal proteins
Purification of high mobility group nonhistone chromosomal proteins by LC on a column containing immobilized histone H5 314(1984)436

Nonylphenyl-oligo(ethylene glycol)
LC fractionation of oligomers of nonylphenyl-oligo(ethylene glycol) 318(1985)289

Noradrenaline
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

HPLC determination of plasma catecholamines during α-methyldopa therapy 308(1984)301

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Dihydrocaffeic acid: a common contaminant in the LC-electrochemical measurement of plasma catecholamines in man 311(1984)148

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their

metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

Nordihydrocapsaicin
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288

11-Nor-D^9-τετραθυδροχανναβηνολ-9-χαρβοξυληχ αχηδ
Detection of marijuana metabolite 11-nor-Δ^9-tetrahydrocannabinol-9-carboxylic acid in human urine by bonded-phase adsorption and TLC 306(1984)441

Norephedrine
Resolution of the enantiomers of ephedrine, norephedrine and pseudoephedrine by HPLC 307(1984)220

Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Selection of mixed ion-pair modifiers for HPLC mobile phases 314(1984)402

Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

Norepinephrine
see Noradrenaline

Norfenfluramine
Determination of fenfluramine and norfenfluramine in plasma using a nitrogen-sensitive detector 310(1984)412

Norfloxacin
HPLC procedure for the quantitation of norfloxacin in urine, serum and tissues 309(1984)177

Norlaudanosine
Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Normetanephrine
Derivatization and MS behaviour of catecholamines and their 3-O-methylated metabolites 310(1984)243

Nornicotine
HPLC method for the determination of N-methylated metabolites of nicotine 306(1984)291

Norpseudoephedrine
Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

Norspermine
GC-MS of polyamines as their N-ethyloxycarbonyl derivatives and identification of *sym*-homospermidine and *sym*-norspermine in mosses and ferns 303(1984)99

Nortriptyline
GC-MS determination of amitriptyline and its major metabolites in human serum 305(1984)73

Amitriptyline and its basic metabolites determined in plasma by GC 308(1984)165

Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Norzimeldine
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Noscapine
Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Nuclear magnetic resonance spectroscopy
Continuous-flow carbon-13 nuclear magnetic resonance spectroscopy 312(1984)91

Nucleic acids
Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 308(1984)121

Affinity chromatography of *Drosophila melanogaster* ribosomal proteins to 5S rRNA 319(1985)367

Separation of large DNA restriction fragments by high-performance gel filtration on TSKgel DNA-PW 320(1985)440

Rapid fractionation of tRNA on benzoylated DEAE-porous glass 324(1985)480

Nucleobases
see also Purines and pyrimidines

Separation of nucleobases on polar amino cyano HPLC columns 303(1984)185

Masking effects in "charge-transfer" TLC 303(1984)190

Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201

Microbore packed-column anion-exchange LC of nucleobases, nucleosides and nucleotides 320(1985)335

Improved fluorometric method for quantitative HPLC analysis of methylated guanine derivatives in DNA 324(1985)484

Separation of naturally occurring or induced methylated nucleobases of DNA by reversed-phase HPLC 325(1985)207

Nucleoside diphosphate kinase
Cibacron Blue 3G-A-substituted cross-linked agar beads: an inexpensive, fast-flow affinity medium for large scale applications 319(1985)218

Nucleosides
Quantitative reversed-phase HPLC of major and modified nucleosides in DNA 301(1984)199

Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Rapid isolation, hydrolysis and chromatography of formaldehyde-modified DNA 308(1984)121

HPLC separations on dihydroxyboryl-agarose 316(1984)301

Microbore packed-column anion-exchange LC of nucleobases, nucleosides and nucleotides 320(1985)335

Preparative and analytical HPLC methods in the synthesis and analysis of decomposition of nitrosourea nucleosides 322(1985)455

Mobile phase optimization for multi-functional solutes in LSC. Resolution of synthetic intermediates of oligodeoxyribonucleotides 323(1985)227

Nucleotides
HPLC method for the simultaneous determination of myocardial creatine phosphate and adenosine nucleotides 305(1984)183

Sensitive fluorimetry of adenine-containing compounds with HPLC 309(1984)63

Role of mobile phase components for multi-functional solutes in silica gel LSC. Resolution of protected deoxyribonucleotide derivatives for oligomer synthesis 316(1984)389

Measurement of plant nucleotides by HPLC 317(1984)271

HPLC purification of deoxynucleoside 5'-triphosphates and their use in a sensitive electrophoretic assay of misincorporation during DNA synthesis 317(1984)283

Ion-pair chromatography of nucleotides 317(1984)301

Isotachophoretic determination of mobility and pK_a by means of computer simulation. III. Evaluation of mobility and pK_a of fifteen nucleotides and seven phosphorus oxoacids and their isotachophoretic separation 318(1985)195

Structure-retention relationships: chromatographic behaviour and properties of nucleotide-cation complexes 318(1985)221

HPLC of nucleotides and nucleotide sugars extracted from wheat embryo and vegetable seed 318(1985)433

Preparative separation of nucleotides by ion exchange HPLC using a volatile buffer system 319(1985)461

Advantages and limitations of isotachophoretic determinations of metabolites in tissue extracts 320(1985)105

Microbore packed-column anion-exchange LC of nucleobases, nucleosides and nucleotides 320(1985)335

HPLC determination of adenine nucleotides in biological materials. Improvements and adaptations to routine analysis 325(1985)314

Nucleotide sugars
HPLC of nucleotides and nucleotide sugars extracted from wheat embryo and vegetable seed 318(1985)433

Rapid resolution of nucleotide sugars by lectin affinity HPLC 323(1985)434

+N-(1,3-Dimethylbutyl)-N'-phenyl-*p*-phenylenediamine
Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141

O

Ocimum gratissimum
GLC-MS in the analysis of essential oils 303(1984)109

Octadecenyl acetate
Microanalytical methods in the structure elucidation of sex-specific components in the large pine weevil, *Hylobius abietis* L. (*Coleoptera, Curculionidae*) 323(1985)343

Octadecyl acetate
Microanalytical methods in the structure elucidation of sex-specific components in the large pine weevil, *Hylobius abietis* L. (*Coleoptera, Curculionidae*) 323(1985)343

Octadecyl-silica
Structural investigations on reversed-phase silicas. Determination of ligand functionality 323(1985)273

Octyl-agarose
Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. I. Sorption isotherms of *n*-alcohols on octyl-agarose 322(1985)399

Octyl-silica
Structural investigations on reversed-phase silicas. Determination of ligand functionality 323(1985)273

Oestradiol
Rapid measurement of oestradiol and oestriol by HPLC after automatic pretreatment 311(1984)249

Oestradiol valerate
HPLC analysis of estradiol valerate-testosterone enanthate in oily formulations 301(1984)292

Oestriol
Rapid measurement of oestradiol and oestriol by HPLC after automatic pretreatment 311(1984)249

Oestrogen conjugates
Separation of oestrogen conjugates in urine and synthetic mixtures by HPLC methods (Review) 309(1984)1

Oestrogen receptors
High-performance size exclusion chromatography as a rapid method for the separation of steroid hormone receptors 307(1984)39

Oestrogens
Separation of oestrogen conjugates in urine and synthetic mixtures by HPLC methods (Review) 309(1984)1

Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313

General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470

Oils
On-line GC evaluation of iodine values in edible fats and oils 314(1984)233

Oleanane saponins
HPLC of oleanane saponins 315(1984)441

α-Olefinsulphonates
Analysis of α-olefinsulphonates by HPLC 325(1985)43

Oleic acid
Analysis of phospholipids in cow's milk by high-temperature injection GC and HPLC 315(1984)223

Oligo(D-galactosiduronic acids)
Product analysis of the action of *Trichoderma reesei* polygalacturonase by TLC 312(1984)492

Oligoglucans
Partitioning of oligoglucans in Sephadex G-15 in relation to their conformational structure 318(1985)57

Oligopeptides
Development of immobilized metal affinity chromatography. III. Interaction of oligopeptides with immobilized nickel iminodiacetate 323(1985)265

Oligosaccharides
Determination of fructan oligomers of degree of polymerization 2-30 by HPLC 303(1984)417

Comparison of gel permeation and ion-exchange chromatographic procedures for the separation of hyaluronate oligosaccharides 320(1985)351

HPLC of oligosaccharides in water on a reversed phase doped with primary amino groups 320(1985)408

Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

HPLC elution behaviour of oligosaccharides, monosaccharides and sugar degradation products on series-connected ion-exchange resin columns using water as the mobile phase 322(1985)411

Oligostyrenes
Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Oltipraz
Method for the determination of oltipraz, a new antischistosomal agent, in blood 305(1984)465

Omeprazole
Determination of omeprazole and metabolites in plasma and urine by LC 309(1984)347

Oxygen effects in amperometric LC detection of omeprazole at a mercury electrode 321(1985)375

Open-tubular columns
Open-tubular LC-MS using direct liquid introduction 323(1985)37

Opiate receptor
Liquid-liquid extraction of membranes from calf brain using conventional and centrifugal counter-current distribution techniques 311(1984)277

Opiates
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Opium
Analysis of opium and crude morphine samples by capillary GC. Comparison of impurity profiles 315(1984)404

Opium alkaloids
Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Optimization
Window analysis optimization of GC separations using mixed Porapaks 302(1984)119

Computer-assisted optimization of eluent concentration and pH in ion chromatography 303(1984)321

Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

Automated optimisation of reversed-phase HPLC separations. An improved method using the sequential simplex procedure 316(1984)69

Rapid optimization of the concentration of the ion-pairing reagent in ion-pairing reversed-phase LC 316(1984)231

Retention-optimization strategy for the HPLC ion-pair separation of samples containing basic compounds 316(1984)241

Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23

Optimization in isotachophoresis. The concept of selectivity and separation speed 320(1985)49

General theory for correction for compressibility effects in binary coupled GC columns and the procedure for window diagram optimisation of relative lengths 321(1985)249

Mobile phase optimization for multi-functional solutes in LSC. Resolution of synthetic intermediates of oligodeoxyribonucleotides 323(1985)227

Mobile phase optimization for the reversed-phase LC of complex hydroxyl aromatic mixtures 324(1985)1

Theory of the efficiency of serially connected GC columns and the effect on optimized speed of analysis 325(1985)1

Oral contraceptives
Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313

General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470

Organic acids
see also Carboxylic acids

Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23

Profiles of strongly polar and less polar acids obtained from human blood, plasma and serum by two-step ultrafiltration 309(1984)17

Hydroxycarboxylic and oxocarboxylic acids in urine: products from branched-chain amino acid degradation and from ketogenesis 309(1984)225

Characterization of artefacts produced by treatment of organic acids with diazomethane 309(1984)243

HPLC method for screening disorders of aromatic acid metabolism using a multi-detection system 310(1984)273

Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133

Derivatization at the liquid-solid interface. Application to the trace analysis of organic acids in aqueous samples 314(1984)421

HPLC of organic acids with potentiometric detection using a metallic copper electrode 315(1984)261

Chromatography of aromatic acids on ion exchangers 316(1984)261

Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267

Separation of tricarboxylic acid cycle acids and other related organic acids in insect haemolymph by HPLC 318(1985)112

Determination of dissociation constants of weak electrolytes by capillary isotachophoresis 320(1985)33

LC of organic acids in silage extracts using dual detection 322(1985)352

Organic compounds
Trace analysis of organic compounds by HPLC with photoionization detection 302(1984)43

Chromatographic analysis of organic compounds in the atmosphere 302(1984)173

Organic modifiers
Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79

Organic phosphites
TLC of age resisters for elastomers (phenols and organic phosphites) (in French) 314(1984)349

Organic synthesis
Preparative HPLC as an aid in organic synthesis 316(1984)113

Organoarsenic compounds
High-resolution GC and LC of organoarsenic compounds 325(1985)53

Organochlorine compounds
Combination of automated pre-column trace enrichment and on-line evaporation in HPLC 301(1984)335

Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448

Capillary fused-silica on-column injection of chlorinated pesticides with an ultra-low volume rotary valve 302(1984)205

Selected ion monitoring MS method for the determination of hexachlorocyclohexane isomers in soils 312(1984)403

Evaluation of automated splitless and manual on-column injection techniques using capillary GC for pesticide residue analysis 314(1984)413

Inexpensive, precise method for the determination of chlorinated pesticide residues in soil 321(1985)480

Organometallic compounds
GC studies on group VIB organometallic complexes. The analytical and preparative separation of chromium, molybdenum and tungsten trifluorophosphine carbonyl complexes 301(1984)253

Ionization quenching of a hydrogen atmosphere flame as a detection method for GC 303(1984)331

Inorganic GC (Review) 313(1984)3

Developments in the HPLC of metallo-organic compounds 317(1984)569

GC investigation of organometallic compounds and their carbon analogues. VI. Determination, calculation and correlation of Kováts retention indices for alkylalkoxysilanes 318(1985)41

Organophosphorus compounds
Evaluation of a GC method for calculating vapor pressures with organophosphorus pesticides 314(1984)37

Evaluation of automated splitless and manual on-column injection techniques using capillary GC for pesticide residue analysis 314(1984)413

LC-chloride-attachment negative chemical ionization MS 319(1985)273

Post-column reaction detection system for the determination of organophosphorus compounds by LC 324(1985)53

Analysis of soman and sarin in blood utilizing a sensitive GC-MS method 324(1985)163

Determination of trace amounts of organophosphorus pesticides and related compounds in soils and sediments using capillary GC and a nitrogenphosphorus detector 325(1985)231

Organosulphur compounds
Retention of sulphur and sulphur organics in reversed-phase LC 317(1984)589

Organotin compounds
Speciation of alkyltin compounds by HPLC with a cyanopropyl-bonded silica column 315(1984)351

Ornidazole
Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325

Orobanchoside
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Orotidine
Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Orotidylate
Enzymatic coupled assay procedures that employ HPLC: the synthesis of orotidylate from ribose 317(1984)245

Orphenadrine
Isothermal GC analysis of diphenhydramine after direct injection onto a fused-silica capillary column 307(1984)295

Oryzalin
Derivatization of Oryzalin for characterization by GC-MS 314(1984)450

Osmium
HPLC of homo- and hetero-tetrametallic iron, ruthenium, osmium and nickel clusters 303(1984)404

OV-17
Supercritical fluid chromatography of various samples with different polarities 315(1984)39

Ovalbumin
Ligand-exchange chromatography on copper(II)-modified silica gel. Improvements and use for screening of protein hydrolyzate and quantitation of dipeptides and amino acid fractions 317(1984)41

Protein separations on reversed-phase HPLC minicolumns 317(1984)119

Fractionation of peptic glycopeptides from chicken ovalbumin by reversed-phase HPLC 320(1985)426

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Oxalic acid
Determination of oxalic acid in biological matrices by LC with amperometric detection 315(1984)271

Oxaprotiline
Measurement of maprotiline and oxaprotiline in plasma by HPLC of fluorescent derivatives 309(1984)107

Oxaprozin
Rapid HPLC analysis of oxaprozin, a non-steroidal anti-inflammatory agent 310(1984)445

Oxcarbazepine
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Oximes
Retention times of oximes in reversed-phase HPLC 301(1984)492

Oxipurinol
HPLC with polarographic and voltammetric anodic detection: simultaneous determination of allopurinol, oxipurinol and uric acid in body fluids 306(1984)205

Simultaneous determination of allopurinol, oxipurinol, hypoxanthine and xanthine in biological fluids by HPLC 307(1984)469

Oxiracetam
Influence of solute polarity in column-switching chromatography for the assay of drugs in plasma and urine 317(1984)493

Oxmetidine
Improved assay procedure for oxmetidine and its metabolites in plasma, urine and bile samples 305(1984)214

HPLC determination of oxmetidine in human plasma: comparison of liquid-liquid and liquid-solid extraction techniques 317(1984)475

Oxoalcohol ethoxylates
On-line LC-MS analysis of non-ionic surfactants 323(1985)135

3-Oxocarboxylic acid esters
Reaction gas chromatographic studies of 3-oxocarboxylic acid esters with a tautomeric enol form. I. A semi-quantitative study (in German) 303(1984)375

Oxocarboxylic acids
Hydroxycarboxylic and oxocarboxylic acids in urine: products from branched-chain amino acid degradation and from ketogenesis 309(1984)225

Δ_{5-4} 3-Oxosteroid isomerase
Sepharose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 319(1985)43

Oxovanadium
Determination of vanadate(V) by conductometric anion chromatography 320(1985)450

Oxprenolol
Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Oxycodone
Structural influences on the amperometric detection of opiates in HPLC 301(1984)39

Determination of oxycodone in human plasma by HPLC with electrochemical detection 308(1984)359

Oxygen
Oxygen effects in amperometric LC detection of omeprazole at a mercury electrode 321(1985)375

Oxytetracycline
Improvement of chemical analysis of antibiotics. VIII. Application of prepacked C_{18} Cartridge for the analysis of tetracycline residues in animal liver 325(1985)265

Quantitative analysis of doxycycline and related substances by HPLC 325(1985)413

P

Pairing ion chain length
Studies in reversed-phase ion-pair chromatography. IV. The role of the chain length of the pairing ion 303(1984)29

Palladium chelates
Volatility and GC of some derivatives of Schiff base chelates 318(1985)350

Panax ginseng
Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497

D-Panthenol
Direct determination of D-panthenol in pharmaceutical preparations by ion-pair chromatography 303(1984)291

Papain
Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

Kinetics of unfolding of proteins on hydrophobic surfaces in reversed-phase LC 317(1984)227

Parabens
Determination of preservatives in cosmetic products by reversed-phase HPLC 315(1984)465

Paracetamol
Improved HPLC separation of urinary paracetamol metabolites using radially compressed columns 306(1984)215

Simultaneous HPLC assay of acetylsalicylic acid and salicylic acid in film-coated aspirin tablets 317(1984)507

Paraffin-coated silica gel
Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133

Paraquat
Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Paraxanthine
Simultaneous determination of caffeine and its N-demethylated metabolites in umbilical cord plasma using HPLC 306(1984)191

Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107

Particle size
Silica gel particle size and coating thicknesses of Chromarods-S and -SII 323(1985)410

Partition coefficients
Selection of GC stationary phase pairs for characterization of the 1-octanol-water partition coefficient 301(1984)355

Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Comparison of phenyl- and octadecyl-modified silica gel as stationary phase for the prediction of *n*-octanol-water partition coefficients by HPLC 320(1985)271

Lipophilicity measurements of protonated basic compounds by reversed-phase HPLC. II. Procedure for the determination of a lipophilic index measured by reversed-phase HPLC 320(1985)305

Mixed solvents in GLC. Activity coefficients for chloroform, dichloromethane and ethyl acetate in squalane-dinonyl phthalate mixtures at 303°K 324(1985)231

Rôle of displacement effects in trapping trace components from gases in sorbent-packed columns 324(1985)269

Partitioning
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47

Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Partitioning of oligoglucans in Sephadex G-15 in relation to their conformational structure 318(1985)57

Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.*-butanol 319(1985)247

Determination of ionic charge by liquid-liquid partition 322(1985)425

PCBs
Separation of polychlorinated biphenyls (Aroclor 1254) by HPLC 302(1984)159

High resolution GC analysis of cytochrome P-448 inducing polychlorinated biphenyl congeners in hazardous waste 314(1984)482

Polychlorinated biphenyl retention time standards obtained by chemical dechlorination of polychlorinated biphenyl isomers 318(1985)211

Determination of polychlorinated biphenyls in copper phthalocyanine pigments 325(1985)456

Peak capacity
Calculation of the peak capacity in capillary GC 312(1984)1

Effect of high speed on peak capacity in LC 316(1984)81

Peak distortion
Characterization of extracolumn and concentration-dependent distortion of chromatographic peaks by Edgeworth-Cramér series 315(1984)67

Peak overlap
Analytical solution for the area under chromatographic peaks obtained from the deconvolution method of Vaidya and Hester 303(1984)177

Application of photodiode array UV detector to unresolved peak analysis 316(1984)473

Peak purity
Limitation of assessing HPLC peak purity with photodiode array detectors 319(1985)387

Peak shape
Effects of amine modifiers on retention and peak shape in reversed-phase HPLC 320(1985)313

HPLC analysis of basic drugs on silica columns using non-aqueous ionic eluents. I. Factors influencing retention, peak shape and detector response 323(1985)173

Pectin
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Penicillamine
Determination of penicillamine in encapsulated formulations by HPLC 318(1985)404

Penicillin G
Determination of penicillin G anmd cloxacillin residues in beef and pork tissue by HPLC 317(1984)311

Studies on residual antibacterials in foods. III. HPLC determination of penicillin G in animal tissues using an on-line pre-column concentration and purification system 318(1985)299

Penicillins
Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

Pentachlorophenol
Identification of bromotetrachlorophenol in commercial pentachlorophenol samples 314(1984)476

1,3,3,5,5,-Pentakis-(azaridino)-λ^6,2,4,6,3λ^5,5λ^5-thia-triazadiphosphorine-1-oxide
GLC determination of a new anticancer agent, 1,3,3,5,5-pentakis-(azaridino)-λ^6,2,4,6,3λ^5,5λ^5-thiatriazadiphosphorine-1-oxide, for pharmacokinetic investigations (in French) 305(1984)335

Pentoxifylline
Determination of 3,7-dimethyl-1-(5-oxohexyl)-xanthine (pentoxifylline) and its 3,7-dimethyl-1-(5-hydroxyhexyl)xanthine metabolite in the plasma of patients with multiple diseases using HPLC 305(1984)419

GC determination of pentoxifylline and its major metabolites in human breast milk 310(1984)61

Pepsin
Fractionation of peptic glycopeptides from chicken ovalbumin by reversed-phase HPLC 320(1985)426

Peptidases
Ion-exchange chromatographic assay of peptidases acting on the C-terminal hexapeptide sequence of substance P 301(1984)415

Peptide mapping
Analytical peptide mapping by ion-exchange HPLC: application to haemoglobin variants 305(1984)456

Peptides
Purification of a cholecystokinin fragment peptide in the horizontal flow-through coil planet centrifuge 301(1984)277

Sequence analysis of derivatized peptides by HPLC-MS 301(1984)425

Reversed-phase HPLC and separation of peptide hormones in human tissues and plasma with reference to cholecystokinins 305(1984)43

Separation of peptides on a polystyrene resin column 307(1984)29

Combined HPLC-radioimmunoassay for the characterization and quantitative measurement of neuropeptides 312(1984)273

Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Reversed-phase HPLC studies of α-MSH fragments 314(1984)313

Separation of the enkephalins from proteins in an aqueous medium by chromatofocusing 314(1984)499

Separation of hydrophobic peptide polymers by size-exclusion and reversed-phase HPLC 317(1984)1

Ligand-exchange chromatography on copper(II)-modified silica gel. Improvements and use for screening of protein hydrolyzate and quantitation of dipeptides and amino acid fractions 317(1984)41

Effects of HPLC solvents and hydrophobic matrices on the secondary and quaternary matrices on the secondary and quaternary structure of a model protein. Reversed-phase and size-exclusion HPLC 317(1984)129

Separation of sequence isomeric dipeptides by high-resolution GC 318(1985)384

Simple and rapid identification of phosphorylated peptides from bovine brain myelin basic protein by reversed-phase HPLC 319(1985)359

Selection of optimum conditions for the preparative-scale separation of a dipeptide diastereomer mixture on the basis of measurements made in an analytical column 320(1985)397

Rapid high-yield purification of canine intestinal motilin and its complete sequence determination 321(1985)421

Separation and purification of diastereomers of angiotensin I by weak anion-exchange HPLC 322(1985)117

Use of benzoin as pre-column fluorescence derivatization reagent for the HPLC of angiotensins 322(1985)473

Development of immobilized metal affinity chromatography. III. Interaction of oligopeptides with immobilized nickel iminodiacetate 323(1985)265

Complete automatization of peptide maps by reversed-phase LC using *o*-phthalaldehyde pre-column derivatization 323(1985)373

Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395

Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173

Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

Determination of epimeric peptides for assessing enantiomeric purity of starting materials and studying racemization in peptide synthesis using HPLC 325(1985)111

Prediction of retention data of protected dipeptides in normal phase HPLC 325(1985)282

Perfluorocarboxylic acids
Use of perfluorinated carboxylic acids in the separation of aminoglycoside antibiotics by ion-pair reversed-phase HPLC 303(1984)277

Perhexiline
Single-dose pharmacokinetics of perhexiline administered orally to humans 305(1984)401

Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Perhydro-1,4-thiazepine-3,5-dicarboxylic acid
Quantitative analysis of cystathionine and perhydro-1,4-thiazepine-3,5-dicarboxylic acid in the urine of a patient with cystathioninuria using isotachophoresis 311(1984)183

Perloline
Determination of the alkaloid perloline in grasses by HPLC 322(1985)261

Permethrin
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Peropyrene
Solvated structure-retention relationships of peropyrene-type polycyclic aromatic hydrocarbons 322(1985)275

Peroxyoxalate chemiluminescence
Solvent and pH effects on peroxyoxalate chemiluminescence detection for LC 314(1984)155

Pesticides
Improved analysis of acephate and methamidophos in biological samples by selective ion monitoring GC-MS 301(1984)465

Capillary fused-silica on-column injection of chlorinated pesticides with an ultra-low volume rotary valve 302(1984)205

Comparison of a new electrochemical detector for GC analysis with the electrolytic conductivity detector 302(1984)269

Anomalies in the HPLC determination of diquat in water samples 303(1984)206

HPLC analysis of rotenone and rotenonone in water by direct injection 303(1984)263

Determination of 3-amino-1,2,4-triazole (amitrole) in urine by ion-pair HPLC 303(1984)267

Detection of phosmethylan and its major metabolites by TLC 303(1984)309

Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Analysis of selected pesticides by HPLC-MS 312(1984)221

Selected ion monitoring MS method for the determination of hexachlorocyclohexane isomers in soils 312(1984)403

Evaluation of a GC method for calculating vapor pressures with organophosphorus pesticides 314(1984)37

The mass spectrometer as a chlorine-selective chromatographic detector. II. Applications to chemical systems 314(1984)93

Evidence for surface effects in an electron-capture detector 314(1984)219

Evaluation of automated splitless and manual on-column injection techniques using capillary GC for pesticide residue analysis 314(1984)413

Derivatization of Oryzalin for characterization by GC-MS 314(1984)450

Simultaneous determination of pynamin forte, resmethrin and piperonyl butoxide in insecticide formulations by GC 314(1984)471

Identification of bromotetrachlorophenol in commercial pentachlorophenol samples 314(1984)476

High-efficiency resolution of isomeric rotenone compounds by HPLC 317(1984)383

Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Determination of benomyl and its metabolites by cation-exchange HPLC 317(1984)527

Determination of phenylurea pesticides by HPLC with UV and photoconductivity detectors in series 317(1984)533

Reversed-phase HPLC method for the simultaneous microquantitative determination of the herbicide isoproturon and its possible degradation products in soil 318(1985)144

LC-chloride-attachment negative chemical ionization MS 319(1985)273

Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Inexpensive, precise method for the determination of chlorinated pesticide residues in soil 321(1985)480

HPLC analysis of the photodegradation products of phosmethylan 322(1985)265

On-line trace enrichment for improved sensitivity in LC with direct liquid introduction MS detection 323(1985)113

Post-column reaction detection system for the determination of organophosphorus compounds by LC 324(1985)53

Determination of trace amounts of organophosphorus pesticides and related compounds in soils and sediments using capillary GC and a nitrogenphosphorus detector 325(1985)231

Application of electrochemical detection to the determination of residues of some carbamate insecticides by HPLC (in French) 325(1985)239

Petroleum distillation residues
Determination of the molar mass of petroleum distillation residues using GPC 312(1984)261

Petroleum sulphonates
Separation of sulfonate and carboxylate mixtures by ion-exchange HPLC 302(1984)65

pH gradients
Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

pH optimization
Computer-assisted optimization of eluent concentration and pH in ion chromatography 303(1984)321

Phase loading
Influence of phase loading on the performance of whisker-walled open tubular columns coated with organic molten salts 324(1985)415

Phase ratio
Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1

Pheliposide
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Phenacetin
MS determination of N-hydroxyphenacetin in urine using multiple metastable peak monitoring following TLC 310(1984)179

Phenanthroline
Effects of various strengths of acids on the electrophoretic behavior of weak bases in a non-aqueous solvent 314(1984)65

Pheneturide
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Phenobarbital
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Automated analysis of antiepileptic drugs in serum by column-switching HPLC 309(1984)357

Routine determination of eight common antiepileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Phenol
Use of LC with dual-electrode electrochemical detection in the investigation of glutathione oxidation during benzene metabolism 317(1984)579

Phenolic amines
GC analysis of endogenous catecholamines, phenolic amines and derived isoquinolines using short glass capillary columns and electron-capture detection 311(1984)17

Phenolic compounds
Analysis of hydroxycinnamic acid esters and their glucosides by reversed-phase HPLC after polyamide separation 315(1984)243

Chromatographic separation and identification of naturally occurring chlorogenic acids by ^1H nuclear magnetic resonance spectroscopy and MS 315(1984)253

Isolation, identification and determination of 3-(4-hydroxyphenyl)propionic acid by HPLC of the phenolic compounds of *Loiseleuria procumbens* Desf. (Ericaceae) (in French) 322(1985)257

Determination of 2,6-di-*tert.*-butyl-4-methylphenol in aviation turbine fuel by LC with electrochemical detection 322(1985)376

Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Phenolic ketones
HPLC separation of pungency components of ginger 319(1985)471

Phenols
Dependence of the retention of phenols upon Van der Waals volume, π-energy and hydrogen-bonding effects 302(1984)89

Long-chain phenols. XXV. Quantitative analysis of natural cashew nut-shell liquid (*Anacardium occidentale*) by HPLC 303(1984)137

Determination of germicidal phenols in blood by capillary column GC 307(1984)420

TLC of age resisters for elastomers (phenols and organic phosphites) (in French) 314(1984)349

Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267

Analysis of long-chain phenols in the sap of the Burmese lac tree, *Melanorrhoea usitate* by capillary GLC 318(1985)378

Identification of phenols of 7,12-dimethylbenz[*a*]anthracene by conversion into their methyl ethers 319(1985)153

GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

Application of electrochemical detection to the determination of residues of some carbamate insecticides by HPLC (in French) 325(1985)239

Phenothiazines
Screening procedure for detection of phenothiazine and analogous neuroleptics and their metabolites in urine using a computerized GC-MS technique 306(1984)125

Assay of plasma thioridazine and metabolites by HPLC with amperometric detection 307(1984)457

Determination of mequitazin in human plasma and urine by capillary column GLC-MS 309(1984)391

Phenoxy acids
Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Phenoxyphenyl polysiloxane
Polarizable polysiloxane stationary phases for capillary column GC 302(1984)303

Phenoxypropane-1,2-diols
HPTLC assay for epoxide hydrolase activity and the determination of phenoxypropane-1,2-diols 318(1985)343

Phenoxypropanolamines
Influence of mobile phase additives in LC of phenoxypropanolamines 316(1984)291

Phenyl acetates
GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

Phenylalanine
Enantiomeric resolution of an optically active guanine derivative by HPLC with phenylalanine-Cu(II) in the mobile phase 303(1984)217

Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

Phenylalanine methyl ester
R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

1-Phenylbutane-1,3-dione
Volatility studies of metal chelates. VII. GC behaviour of chelates derived from the aromatic β-diketones 1-phenylbutane-1,3-dione and 4,4,4-trifluoro-1-(pentafluorophenyl)butane-1,3-dione 322(1985)23

Phenylephrine
Selection of mixed ion-pair modifiers for HPLC mobile phases 314(1984)402

5-Phenyl-5-ethylhydantoin
Direct enantiomeric resolution of mephenytoin and its N-demethylated metabolite in plasma and blood using chiral capillary GC 307(1984)121

Phenylethylmalonamide
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Phenylglycine
R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Phenylglycinol
R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

N-Phenyl-2-naphthylamine
Quantitative analysis of aromatic amines in human urine by GC-MS-selected-ion monitoring 311(1984)141

***o*-Phenylphenol**
Determination of germicidal phenols in blood by capillary column GC 307(1984)420

Phenylpropanolamine
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Selection of mixed ion-pair modifiers for HPLC mobile phases 314(1984)402

Phenylureas
Determination of phenylurea pesticides by HPLC with UV and photoconductivity detectors in series 317(1984)533

On-line trace enrichment for improved sensitivity in LC with direct liquid introduction MS detection 323(1985)113

Phenytoin
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Automated analysis of antiepileptic drugs in serum by column-switching HPLC 309(1984)357

Routine determination of eight common anti-epileptic drugs and metabolites by HPLC

using a column-switching system for direct injection of serum samples 310(1984)97

Determination of plasma phenytoin by capillary GC with nitrogen-phosphorus detection and with selective ion monitoring 310(1984)199

Pheromones
Microanalytical methods in the structure elucidation of sex-specific components in the large pine weevil, *Hylobius abietis* L. (*Coleoptera, Curculionidae*) 323(1985)343

Phoenicoxanthins
Direct diastereomeric resolution of carotenoids. I. 3-Hydroxy-4-oxo β-end group 318(1985)122

Phosgene
Phosgene: a versatile reagent for enantiomer separation by capillary GC 301(1984)129

Phosmet
Rapid separation of pesticides by HPLC with 3-μm columns 317(1984)523

Phosmethylan
Detection of phosmethylan and its major metabolites by TLC 303(1984)309

HPLC analysis of the photodegradation products of phosmethylan 322(1985)265

Phosphate
Simultaneous determination of phosphate and phosphonate by flow injection analysis and HPLC with a series detection system 318(1985)319

HPLC separation of glucose-1-phosphate, fructose, sucrose and inorganic orthophosphate 322(1985)220

Phosphatidic acid
Formation and separation of fluorescent derivatives of phosphatidic acid 320(1985)421

Phosphatidylcholine
Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65

HPLC of phospholipids using deuterated solvents for IR detection 307(1984)261

Separation of molecular species of phosphatidylcholine by HPLC on a PLRP-S column 325(1985)473

Phosphatidylethanolamine
HPLC of phospholipids using deuterated solvents for IR detection 307(1984)261

Phosphites
TLC of age resisters for elastomers (phenols and organic phosphites) (in French) 314(1984)349

Phosphoamino acids
Automatic amino acid and sugar analysis of glycoproteins 317(1984)35

Phosphoenolpyruvate carboxylase
Isolation of plant phosphoenolpyruvate carboxylase by high-performance size-exclusion chromatography 315(1984)430

3-Phosphoglycerate kinase
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Phosphoinositides
Simplified procedure for the quantitation of radioactive phosphoinositides by TLC 323(1985)462

Phospholipase C
Fast protein LC for the isolation of *Clostridium perfringens* type A α-toxin 321(1985)235

Phospholipids
Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65

HPLC of phospholipids using deuterated solvents for IR detection 307(1984)261

Rapid isocratic HPLC purification of platelet activating factor (PAF) and *lyso*-PAF from human skin 309(1984)160

Analysis of phospholipids in cow's milk by high-temperature injection GC and HPLC 315(1984)223

Separation of molecular species of phosphatidylcholine by HPLC on a PLRP-S column 325(1985)473

Phosphonate
Simultaneous determination of phosphate and phosphonate by flow injection analysis and HPLC with a series detection system 318(1985)319

Phosphonodipeptides
Chromatographic assignment of the configuration of 1-aminoalkanephosphonic acids 324(1985)455

Phosphoribosyl α-1-pyrophosphate synthetase
Enzymatic coupled assay procedures that employ HPLC: the synthesis of orotidylate from ribose 317(1984)245

Phosphoribosyltransferase
Enzymatic assay procedures that employ HPLC: competition between phosphoribosyltransferases for a common substrate 316(1984)43

Phosphoric acid
Factors influencing the reproducibility of isotachophoretic determinations of ions present in wet-process phosphoric acid 325(1985)161

Phosphorus compounds
New thermionic detector utilizing orthogonal nebulization for capillary column LC 314(1984)103

Phosphorus oxoacids
Isotachophoretic determination of mobility and pK_a by means of computer simulation. III. Evaluation of mobility and pK_a of fifteen nucleotides and seven phosphorus oxoacids and their isotachophoretic separation 318(1985)195

Phosphorylated peptides
Simple and rapid identification of phosphorylated peptides from bovine brain myelin basic protein by reversed-phase HPLC 319(1985)359

Photoionization detection
Trace analysis of organic compounds by HPLC with photoionization detection 302(1984)43

Phthalates
Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53

Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Supercritical fluid chromatography of various samples with different polarities 315(1984)39

Development of an on-line high-pressure pre-column concentration method for micro HPLC and its application to the rapid determination of dibutyl phthalate in water 321(1985)159

Phytanic acid
Improved method for GC determination of phytanic acid 325(1985)336

Picein
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Picrorrhiza kurroa
Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497

Picrosides
Quantitative determination of biologicaly active constituents in crude extracts of medicinal plants by TLC-densitometry. II. *Eleutherococcus senticosus* Maxim., *Panax ginseng* Meyer and *Picrorrhiza kurroa* Royle 312(1984)497

Pindolol
Determination of pindolol in human plasma by HPLC with amperometric detection 311(1984)430

Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Pipecolic acid
HPLC of amino acids in urine and cerebrospinal fluid 302(1984)21

Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251

Piperonyl butoxide
Simultaneous determination of pynamin forte, resmethrin and piperonyl butoxide in insecticide formulations by GC 314(1984)471

Piracetam
Determination of 3,7-dimethyl-1-(5-oxohexyl)-xanthine (pentoxifylline) and its 3,7-dimethyl-1-(5-hydroxyhexyl)xanthine metabolite in the plasma of patients with multiple diseases using HPLC 305(1984)419

Pirlimycin
HPLC assay of pirlimycin in human serum and urine using 9-fluorenylmethylchloroformate 309(1984)95

Pirlindole
Determination of pirlindole in plasma and urine by HPLC 309(1984)115

Piroxicam
Rapid method for the determination of either piroxicam or tenoxicam in plasma using HPLC 310(1984)455

Pivaloyloxyethyl3-(3,4-dihydroxyphenyl)-2-methylalaninate
Sensitive HPLC assay using electrochemical detection for a novel prodrug ester of methyldopa, pivaloyloxyethyl 3-(3,4-dihydroxyphenyl)-2-methylalaninate, in plasma and urine 308(1984)251

Plant hormones
Trace enrichment of indole-3-acetic acid from leaf matrix 325(1985)127

Separation of conjugates and oxidative metabolites of indole-3-acetic acid by HPLC 325(1985)340

Less hazardous derivatization procedure for GC of plant hormone abscisic acid 325(1985)425

Platelet activating factor
Rapid isocratic HPLC purification of platelet activating factor (PAF) and *lyso*-PAF from human skin 309(1984)160

Platinum
Sensitive HPLC assay for platinum in plasma ultrafiltrate 306(1984)417

Platinum complexes
Analysis of platinum complexes by LC with quenched phosphorescence detection 312(1984)337

Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161

Analysis of antitumour [1,1-bis(aminomethyl)cyclohexane]platinum(II) complexes derived from spiroplatin by HPLC with differential pulse amperometric detection 320(1985)379

Polarity
Influence of solute polarity in column-switching chromatography for the assay of drugs in plasma and urine 317(1984)493

Study of Kováts retention indices of aliphatic saturated esters and their relation to the polarity of the stationary phase 318(1985)187

Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1

Poliumoside
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Polyamines
GC-MS of polyamines as their N-ethyloxycarbonyl derivatives and identification of *sym*-homospermidine and *sym*-norspermine in mosses and ferns 303(1984)99

Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59

N-Permethylation of polyamines at trace levels for GC analysis 322(1985)169

Polyanions
Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161

Polychlorinated biphenyls
see PCBs

Polychromophore polymers
Molecular weight determination of polychromophore polymers using HPLC and size-exclusion chromatography with UV detection 322(1985)338

Polycyclic aromatic hydrocarbons
see Polynuclear aromatic hydrocarbons

Polyether polyurethanes
Chromatographic analysis of polyether based polyurethanes after alkali fusion 319(1985)121

Polyethylene
Products obtained during combustion of polymers under simulated incinerator conditions. I. Polyethylene 314(1984)263

Poly(ethylene oxide)
Sepharose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 319(1985)43

Preparative fractionation of fluorescent-dye-labeled poly(ethylene oxide) by aqueous size-exclusion chromatography 319(1985)241

Polygalacturonase
Product analysis of the action of *Trichoderma reesei* polygalacturonase by TLC 312(1984)492

Polyhydroxyl compounds
Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

Polyion ion-pair chromatography
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Polymers
Products obtained during combustion of polymers under simulated incinerator conditions. I. Polyethylene 314(1984)263

Separation of hydrophobic peptide polymers by size-exclusion and reversed-phase HPLC 317(1984)1

Effects of HPLC solvents and hydrophobic matrices on the secondary and quaternary matrices on the secondary and quaternary structure of a model protein. Reversed-phase and size-exclusion HPLC 317(1984)129

Effect of solute size and shape on the elution volume in size-exclusion chromatography 319(1985)115

Preparative fractionation of fluorescent-dye-labeled poly(ethylene oxide) by aqueous size-exclusion chromatography 319(1985)241

Semi-micro size-exclusion chromatography of polymers. Calibration of columns and comparison with conventional columns 320(1985)343

Characterization of epoxy novolac resins using phenyl-bonded phase HPLC 321(1985)325

Molecular weight determination of polychromophore polymers using HPLC and size-exclusion chromatography with UV detection 322(1985)338

Determination of polymer molecular weights by PC 325(1985)363

Polymethylsiloxane
Separation of underivatised barbiturates by capillary column GC. Preparation of medium polar polymethylsiloxane columns to optimize selectivity 318(1985)255

Polynuclear aromatic hydrocarbons
Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448

Comparison of methods for separating polycyclic aromatic hydrocarbons by HPTLC 302(1984)149

Sample enrichment for GC-MS analysis of polynuclear aromatic hydrocarbons in water and in organic mixtures 302(1984)167

New schemes for the electron-capture sensitization of aromatic hydrocarbons 302(1984)243

Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53

Detection of GC eluates by simultaneous absorbance and fluorescence measurements 303(1984)341

Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Quantitative analysis of polycyclic aromatic compounds in diesel exhaust particulate extracts by combined chromatographic techniques 312(1984)183

Determination of polycyclic aromatic hydrocarbons in diesel exhaust particulate matter and diesel fuel oil 312(1984)247

Supercritical fluid chromatography and supercritical fluid chromatography-MS of marine diesel fuel 314(1984)253

Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Products obtained during combustion of polymers under simulated incinerator conditions. II. Polystyrene 315(1984)201

Products obtained during combustion of polymers under simulated incinerator conditions. III. Polyvinyl chloride 315(1984)211

Computer-assisted retention prediction system for reversed-phase micro HPLC 316(1984)1

Direct resolution of mono- and diol enantiomers of unsubstituted and methyl-substituted benz[a]anthracene and benzo[a]pyrene by HPLC with a chiral stationary phase 316(1984)569

HPLC studies of environmental carcinogens in China 317(1984)545

Identification of phenols of 7,12-dimethylbenz[a]anthracene by conversion into their methyl ethers 319(1985)153

Separation of polyaromatic hydrocarbons on caffeine-bonded silica gel 319(1985)432

Chromatography with sub- and supercritical eluents: influence of temperature, pressure and flow-rate on the behaviour of lower alkanes 321(1985)273

LC systems for the separation of polycyclic aromatic hydrocarbon and nitrogen heterocycle compounds present in coal liquids 321(1985)343

Fingerprinting of crude oil spills 322(1985)107

Solvated structure-retention relationships of peropyrene-type polycyclic aromatic hydrocarbons 322(1985)275

Chromatographic selectivity in LC of polycondensed aromatic hydrocarbons on 3-(2,4-Dinitroanilino)propyl silica 324(1985)261

Identification of trace organic compounds in dimethyl sulphoxide solution using HPLC and GC-MS 324(1985)373

Polypeptides
Isolation and structural studies of porcine, ovine and murine thymosin β_4 by HPLC 301(1984)221

Fractionation of thylakoid membrane components by extraction in aqueous polymer two-phase systems containing detergent 323(1985)363

Polyphosphates
Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Polysaccharides
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 319(1985)205

Analyses of homogeneous D-gluco-oligosaccharides and -polysaccharides (degree of polymerization up to about 35) by HPLC and TLC 321(1985)145

Polyselenandisulphonates
Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161

Poly(silarylene-methylphenylsiloxane)
Use of poly(silarylene-methylphenylsiloxane) block copolymer as a thermostable stationary phase in capillary column GC 301(1984)265

Polysilicic acid
Chromatographic studies in the field of silicate chemistry (Review) 313(1984)33

Polysiloxanes
Polarizable polysiloxane stationary phases for capillary column GC 302(1984)303

Polystyrene
Solvent selectivity studies using isomers of polystyrene 314(1984)55

Products obtained during combustion of polymers under simulated incinerator conditions. II. Polystyrene 315(1984)201

Semi-micro size-exclusion chromatography of polymers. Calibration of columns and comparison with conventional columns 320(1985)343

Molecular weight determination of polychromophore polymers using HPLC and size-exclusion chromatography with UV detection 322(1985)338

Poly(styrene-co-acrylonitrile)
Two-dimensional analysis of copolymers by size-exclusion chromatography and gradient-elution reversed-phase precipitation chromatography 317(1984)615

Poly(styrene-divinylbenzene) column
Separation of peptides on a polystyrene resin column 307(1984)29

Polystyrenes
High-performance liquid-solid adsorption chromatography of spiropyran-end-labelled polystyrenes 303(1984)361

Pyrolysis-fluorescence spectroscopy, pyrolysis-MS and pyrolysis-LC of linear and cross-linked polystyrenes 314(1984)283

Supercritical fluid chromatography of various samples with different polarities 315(1984)39

Effect of solute size and shape on the elution volume in size-exclusion chromatography 319(1985)115

Multi-mode LC of polystyrenes on Partisil PAC 324(1985)436

Polysulphandisulphonates
Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161

Polysulphanes
Retention of sulphur and sulphur organics in reversed-phase LC 317(1984)589

Polysulphides
Retention of sulphur and sulphur organics in reversed-phase LC 317(1984)589

Polyurethane foam
Theoretical plate measurements and collection efficiencies for high-volume air samplers using polyurethane foam 301(1984)448

Polyurethanes
Chromatographic analysis of polyether based polyurethanes after alkali fusion 319(1985)121

Chromatographic analysis of polyester-based polyurethane elastomers after alkali fusion 321(1985)295

Poly(vinyl alcohol)
High-performance gel filtration of water-soluble samples on polyvinyl alcohol columns 320(1985)369

Polyvinyl chloride
Products obtained during combustion of polymers under simulated incinerator conditions. III. Polyvinyl chloride 315(1984)211

Multi-stage chromatographic procedure for the isolation of a single oligomeric species (vinyl chloride tetramer) from poly(vinyl chloride) resin used for food packaging applications 320(1985)361

Populin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Porapak
Window analysis optimization of GC separations using mixed Porapaks 302(1984)119

Pore size distribution
Theoretical models for size-exclusion chromatography and calculation of pore size distribution from size-exclusion chromatography data 316(1984)311

Porphyrins
Computerised GC-MS analysis of complex mixtures of alkyl porphyrins 301(1984)107

Separation of haem compounds by reversed-phase ion-pair HPLC and its application in the assay of ferrochelatase activity 310(1984)31

HPLC of dicarboxylic porphyrins and metalloporphyrins: retention behaviour and biomedical applications 317(1984)333

Post-column reactors
Electrochemical detection of cyanogenic glycosides after enzymatic post-column cleavage 303(1984)67

Cyanogenic glycosides and cyanohydrins in plant tissues. Qualitative and quantitative determination by enzymatic post-column cleavage and electrochemical detection, after separation by HPLC 303(1984)77

Sub-picomol chemiluminescence detection of Dns-amino acids separated by HPLC with gradient elution 303(1984)173

Use of immobilized amino acid oxidase as post-column reactor in the HPLC of amino acids 303(1984)396

Determination of (endogenous) vitamin K_1 in human plasma by reversed-phase HPLC using fluorometric detection after post-column electrochemical reduction. Comparison with UV, single and dual electrochemical detection 305(1984)61

New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188

HPLC analyser for guanidino compounds using benzoin as a fluorogenic reagent 305(1984)281

Determination of free trimethyllysine in plasma and tissue specimens by HPLC 306(1984)79

Automated anion-exchange chromatographic method for the quantitation of lactulose in physiological fluids and processed milks 307(1984)393

Determination of cyanide and thiocyanate in blood plasma and red cells by HPLC with fluorometric detection 308(1984)133

Determination of sodium 2-mercaptoethanesulphonate by HPLC using post-column reaction colorimetry or electrochemical detection 311(1984)234

Amperometric detection of muramic acid in HPLC with a post-column reaction 316(1984)495

Microbore HPLC and chemiluminescence detection of Dns-amino acids 316(1984)501

Enzymes adsorbed on an ion exchanger as a post-column reactor: application to acetylcholine measurement 317(1984)343

Post-column reaction of amino acids with the pentane-2,4-dione-formaldehyde system for their automated analysis 318(1985)367

Post-column reaction detection system for the determination of organophosphorus compounds by LC 324(1985)53

Determination of urea and ammonia using ion-pair LC with on-line post-column derivatization in an enzymatic solid-phase reactor 325(1985)255

Potassium
Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Potassium-sparing diuretics
HPLC behaviour of some pharmaceutically important thiazide, loop and potassium-sparing diuretics 325(1985)395

Potassium thiocyanate
Chromatography of Astra Blue dyes with eluents containing potassium thiocyanate in organic solvents 319(1985)400

Pre-column
Optimization of precolumn design in LC 301(1984)325

Conical precolumn as loading buffer for the main column 314(1984)183

Pre-column concentration
Development of an on-line high-pressure pre-column concentration method for micro HPLC and its application to the rapid determination of dibutyl phthalate in water 321(1985)159

Pre-column venting technique
Procedures for direct injections of untreated blood plasma into LC columns with emphasis on a pre-column venting technique 317(1984)213

Prednisolone
Application of a high-performance gel permeation LC procedure to the determination of binding of prednisolone to high-affinity binding sites in human serum 305(1984)194

Pregnenolone
Use of steroid profiling in the resolution of pregnenolone metabolites from porcine testicular preparations 301(1984)189

Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Premazepam
Quantitative determination of premazepam in human plasma by HPLC 309(1984)415

Prenalterol
Determination of prenalterol in plasma and urine by LC with electrochemical detection 307(1984)230

Preparative chromatography
Preparative HPLC as an aid in organic synthesis 316(1984)113

Preparative HPLC of proteins on ion-exchange columns with carboxymethyldextrans as displacers 316(1984)125

Process scale-up of a β-lactam antibiotic purification by HPLC 316(1984)133

Selection of optimum conditions for the preparative-scale separation of a dipeptide diastereomer mixture on the basis of measurements made in an analytical column 320(1985)397

Pre-separation
Large sample volume preseparation for trace analysis in isotachophoresis 320(1985)89

Primaquine
Determination of primaquine in biological fluids by reversed-phase HPLC 305(1984)239

HPLC-electrochemical assay method for primaquine in plasma and urine 307(1984)451

Primidone
Simultaneous determination of eleven antiepileptic compounds in serum by HPLC 305(1984)127

Automated analysis of antiepileptic drugs in serum by column-switching HPLC 309(1984)357

Routine determination of eight common antiepileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Procainamide
Ion-exchange HPLC in drug assay in biological fluids. IV. Nadolol diastereomers: demonstration of pharmacokinetic and binding equivalence 309(1984)421

Progabide
Determination of progabide and its main acid metabolite in biological fluids using HPLC and electrochemical detection. Application to the measurement of blood/plasma partition ratio 308(1984)229

Progesterone
Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Progestin receptors
High-performance size exclusion chromatography as a rapid method for the separation of steroid hormone receptors 307(1984)39

Progestogens
Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313

General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470

Prolidase
Measurement of prolidase activity in erythrocytes using isotachophoresis 310(1984)401

Proline
Quantification by selected ion monitoring of pipecolic acid, proline, γ-aminobutyric acid and glycine in rat brain 310(1984)251

Rapid and sensitive method for the determination of proline by reversed-phase HPLC with automated pre-column fluorescence derivatization 318(1985)132

Promethazine
Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Prometryne
Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Propanidid
HPLC of propanidid in rat plasma 308(1984)376

2-Propanol
Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79

Propanol
HPLC of low-molecular-weight aliphatic alcohols using indirect photometric detection 314(1984)488

2-Propanol
Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

Propantheline
Analysis of propantheline bromide in serum by HPLC 306(1984)424

Propiomazine
Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Propranolol
Direct enantiomeric determination of (-)- and (+)-propranolol in human serum by HPLC on a chiral stationary phase 306(1984)405

Silica gel HPLC for the simultaneous determination of propanolol and 4-hydroxypropranolol enantiomers after chiral derivatization 310(1984)424

R-α-Methylbenzyl isothiocyanate, a new and convenient chiral derivatizing agent for the separation of enantiomeric amino compounds by HPLC 314(1984)275

Optical resolution of β-blocking agents by TLC and HPLC as diastereomeric *R*-(-)-1-(1-naphthyl)ethylureas 314(1984)462

Determination of secondary amino drugs as their metal dithiocarbamate complexes by reversed-phase HPLC with electrochemical detection 317(1984)513

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Propylamine
Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Prostaglandins
HPLC assay for prostaglandin E_1 in various ointment vehicles. Separation and stability testing 303(1984)165

Determination of 6-keto-prostaglandin $F_{1\alpha}$ and thromboxane B_2 in urine by HPLC and radioimmunoassay 306(1984)351

Rapid, sensitive method for detecting different arachidonic acid metabolites by TLC: the use of autoradiography 307(1984)185

Quantitative analysis of prostaglandins in cell culture medium by high-resolution GC with electron-capture detection 307(1984)251

Picogram detection of eicosanoids by UV absorbance after narrow-bore HPLC. Comparison with conventional-bore column 308(1984)31

Simple single-step HPLC method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 308(1984)316

GC-MS profiling with negative-ion chemical ionization detection of prostaglandins and their 15-keto and 15-keto-13,14-dihydro catabolites in rat blood 310(1984)233

Determination of leukotrienes and prostaglandins in [^{14}C]arachidonic acid labelled human lung tissue by HPLC and radioimmunoassay 311(1984)39

HPLC determination of acid-catalyzed degradation products of methyl carboprost in a polymeric controlled-release device 314(1984)369

HPLC method for the separation of isomers of 9β-methylcarbacyclin (ciprostene) and the analysis of pharmaceutical samples 321(1985)209

Silver-modified mobile phase for normal-phase LC determination of prostaglandins and their 5,6-*trans* isomers in prostaglandin bulk drugs and triacetin solutions 321(1985)353

Protein conformation
Protein conformation and reversed-phase HPLC 317(1984)93

Proteins
see also Enzymes

Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

Separation of metallothionein into isoforms by column switching on gel permeation and ion-exchange columns with HPLC-atomic-absorption spectrophotometry 303(1984)131

High-performance immunosorbent purification of recombinant leukocyte A interferon 303(1984)225

Fast protein liquid chromatofocusing of lyophilized native haemoglobin and its chemically modified derivatives 303(1984)422

Separation of glyoxylated hemoglobin with varying oxygen affinity by polyanion fast protein LC 303(1984)425

Analytical peptide mapping by ion-exchange HPLC: application to haemoglobin variants 305(1984)456

Separation of peptides on a polystyrene resin column 307(1984)29

Analysis of the globins from fast human haemoglobins by isoelectrofocusing on polyacrylamide gel rods 307(1984)91

Determination of glycosylated haemoglobin by isoelectric focusing in non-linear pH gradients 307(1984)103

Separation of collagen α-chains by reversed-phase HPLC. Comparison of column alkyl stationary phases and temperature effects 308(1984)111

Isolation and purification of HLA-DR antigens from tumour cells by affinity chromatography and chromatofocusing 308(1984)322

HPLC of proteins: purification of α-fetoprotein from fetal calf serum 310(1984)19

Resolution and quantitation of apolipoproteins A-I and A-II from human high-density lipoprotein by size exclusion HPLC 311(1984)79

Detection of changes in rabbit serum proteins after partial hepatectomy by means of two-dimensional electrophoresis under non-denaturing conditions 311(1984)93

HPLC determination of bovine and porcine insulins in a commercial injection 312(1984)476

Purification of high mobility group nonhistone chromosomal proteins by LC on a column containing immobilized histone H5 314(1984)436

Separation of the enkephalins from proteins in an aqueous medium by chromatofocusing 314(1984)499

Fractionation of human serum proteins by immobilized metal affinity chromatography 315(1984)167

HPLC of proteins. Purification of plant enzymes by ion-exchange chromatography 315(1984)333

Preparative HPLC of proteins on ion-exchange columns with carboxymethyldextrans as displacers 316(1984)125

HPLC separations on dihydroxyboryl-agarose 316(1984)301

Wide-pore silica-based ether-bonded phases for separation of proteins by high-performance hydrophobic-interaction and size-exclusion chromatography 316(1984)519

Purification of glycopeptides of human plasma proteins by HPLC 317(1984)11

Hydrophobic interaction HPLC of proteins 317(1984)55

Salt-mediated retention of proteins in hydrophobic interaction chromatography. Application of solvophobic theory 317(1984)67

Myoglobin determination by HPLC 317(1984)87

Protein conformation and reversed-phase HPLC 317(1984)93

Protein separations on reversed-phase HPLC minicolumns 317(1984)119

Effects of HPLC solvents and hydrophobic matrices on the secondary and quaternary matrices on the secondary and quaternary structure of a model protein. Reversed-phase and size-exclusion HPLC 317(1984)129

Comparison of hydrophobic-interaction and reversed-phase chromatography of proteins 317(1984)141

Effect of surfactants on the separation of proteins by reversed-phase HPLC. I. Non-ionic surfactants (Tween) 317(1984)157

High-performance affinity chromatography: a rapid technique for the isolation and quantitation of IgG from cerebral spinal fluid 317(1984)173

Application of HPLC techniques to the separation of ribosomal proteins of different organisms 317(1984)181

Application of HPLC to the reconstitution of ribosomal subunits 317(1984)201

Kinetics of unfolding of proteins on hydrophobic surfaces in reversed-phase LC 317(1984)227

Reversed-phase chromatography of proteins with diphenyl-silica column and hydro-organic eluents containing two organic solvents 318(1985)11

Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157

Separation of hemoglobin variants by ion-exchange chromatography on monobead resins 318(1985)325

Comparison of high-performance ion-exchange chromatography and gel electrophoresis in protein separations 318(1985)358

Separation and quantitative assay of three pathogenesis-related (b)proteins from tobacco mosaic virus hypersensitive *Nicotiana tabacum* by reversed-phase HPLC 318(1985)417

Sepharose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 319(1985)43

Rapid ion-exchange chromatography for preparative separation of proteins. Application to porcine and bovine carbonic anhydrases 319(1985)59

One-step procedure for the rapid isolation of mouse monoclonal antibodies and their antigen binding fragments by fast protein LC on a Mono Q anion-exchange column 319(1985)67

Determination of protein spots separated by two-dimensional polyacrylamide gel electrophoresis 319(1985)79

Subunits of bovine lutropin. Hormonal and immunological activity after separation with reversed-phase HPLC 319(1985)195

Affinity chromatography of *Drosophila melanogaster* ribosomal proteins to 5S rRNA 319(1985)367

Silica-bound polyethyleneglycol as stationary phase for separation of proteins by HPLC 319(1985)396

Rapid fluorescamine based protein assay usable in the presence of interfering substances 319(1985)446

Micropreparative isotachophoretic electrodesorption of monoclonal antibodies from an affinity adsorbent 320(1985)81

Characterization of a heterogeneous chicken plasma protein, HEF, by analytical isotachophoresis in agarose gel 320(1985)119

Isotachophoresis of allergenic extracts 320(1985)205

Industrial autofocusing: a new technology for large-scale isoelectric focusing 320(1985)213

α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Separation of serum proteins on a Fe^{3+}-monohydroxamate adsorbent 321(1985)105

Synthesis of new hydrophobic adsorbents based on homologous series of uncharged alkyl sulphide agarose derivatives 321(1985)305

Reversed-phase HPLC of human haemoglobin chains 321(1985)407

Analysis of membrane proteins by ion-exchange HPLC 322(1985)212

Further experiments on a new, fast method for determining molecular weights of diffusing species in a liquid phase 322(1985)443

Comparison of chromatographic characteristics of a series of homologous Bence-Jones proteins during size-exclusion chromatography by HPLC and by Sephadex 323(1985)418

Reversed-phase HPLC of some ultra-heterogeneous and covalently-modified proteins from human hair 324(1985)65

Identification of fish species by HPLC 324(1985)199

Highly stable pH-gradients formed by polyhydroxyl compounds, boric acid and borates. Theoretical aspects and application to isoelectric focusing of proteins, peptides and amino acids 324(1985)285

Comparison by gel filtration chromatography and reversed-phase HPLC of the immunoreactive growth hormone composition of a human pituitary extract 324(1985)385

Electrophoretic studies of blood globin preparations 324(1985)395

Microbore reversed-phase chromatography of proteins with conventional gradient equipment for HPLC 325(1985)187

***Proteus* species**
Volatile products from acetylcholine as markers in the rapid urine test using head-space GLC 307(1984)11

Proxyphylline
Separation of (*R*)- and (*S*)-proxyphylline as diastereoisomeric camphanates by reversed-phase LC 303(1984)433

Pseudoephedrine
Resolution of the enantiomers of ephedrine, norephedrine and pseudoephedrine by HPLC 307(1984)220

Determination of ephedrine alkaloids by isotachophoresis 324(1985)503

Pseudomorphine
Determination of pseudomorphine in morphine injection by HPLC 312(1984)473

Psilocin
Qualitative and quantitative determinations of hallucinogenic components of psilocybe mushrooms by reversed-phase HPLC 312(1984)467

Psilocybin
Qualitative and quantitative determinations of hallucinogenic components of psilocybe mushrooms by reversed-phase HPLC 312(1984)467

Pumps
Simple device for sequential linear gradient elution with a single, conventional, one-piston reciprocating pump 316(1984)415

Purines
see also Nucleobases

Putrescine
Sensitive assay for diamine oxidase activity using HPLC 309(1984)151

Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59

Pynamin forte
Simultaneous determination of pynamin forte, resmethrin and piperonyl butoxide in insecticide formulations by GC 314(1984)471

Pyrazines
Calculation of retention indices of pyrazines on the basis of molecular structure 324(1985)428

Pyrene
Masking effects in "charge-transfer" TLC 303(1984)190

N-(1--Pyrene)maleimide
Cysteinylglycine in urine determined by HPLC 308(1984)75

Pyrethroides
Combination of automated pre-column trace enrichment and on-line evaporation in HPLC 301(1984)335

1-(2-Pyridyl)piperazine
Separation of isocyanate prepolymer components as their urea derivatives by reversed-phase HPLC 301(1984)485

Pyrimethamine
Quantification of antimalarial drugs. I. Simultaneous measurement of sulphadoxine, N_4-acetylsulphadoxine and pyrimethamine in human plasma 305(1984)502

Rapid GC determination of pyrimethamine in human plasma and urine 306(1984)388

Quantification of antimalarial drugs. II. Simultaneous measurement of dapsone, monoacetyldapsone and pyrimethamine in human plasma 307(1984)426

HPLC assay of pyrimethamine, sulfadoxine and its N^4-acetyl metabolite in serum and urine after ingestion of Suldox 308(1984)217

HPLC method for the determination of pyrimethamine and its 3-N-oxide metabolite in biological fluids 308(1984)363

Pyrimidine nucleoside phosphorylase
Pyrimidine nucleoside phosphorylase assay by automated HPLC 319(1985)351

Pyrimidines
see also Nucleobases

Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201

Pyroglutamic acid
Analysis of glutamine, glutamic acid and pyroglutamic acid in protein hydrolysates by HPLC 322(1985)248

Q

Quaternary ammonium compounds
Retention behavior of long chain quaternary ammonium homologues and related nitroso-alkylmethylamines 324(1985)209

Quercetin
Isolation of quercetin, myricetin, and their respective dihydro-compounds by Sephadex LH-20 chromatography 324(1985)495

Quinidine
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Inter-laboratory study on the separation of cinchona alkaloids by HPLC 318(1985)362

Quinine
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Chromatographic separation of enantiomers of acids with quinine as chiral counter ion 316(1984)553

Determination of quinine in beverages, pharmaceutical preparations and urine by isotachophoresis 320(1985)245

Quinonoid dihydrobiopterin
Determination of quinonoid dihydrobiopterin by LC and electrochemical detection 317(1984)407

R

Radially compressed polythene cartridges
HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99

Radioassay
High-sensitivity radioassay in chromatographic effluents 302(1984)31

Ranitidine
Qualitative and quantitative analysis of ranitidine and its metabolites by HPLC-MS 323(1985)143

Reaction detector
Reaction detector for LC with electrochemical generation and detection to excess bromine 324(1985)333

Relative ionic mobilities
Determination of relative ionic mobilities by capillary isotachophoresis 320(1985)139

Relative molar heats of solution
GC separation and identification of linear and branched-chain alkyl bromides 324(1985)363

Reserpine
Micro column LC-MS using a capillary interface 323(1985)89

Resin acids
Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

Resmethrin
Simultaneous determination of pynamin forte, resmethrin and piperonyl butoxide in insecticide formulations by GC 314(1984)471

Response factor calibration
Determinations of chlorinated hydrocarbons by GC using response factor calibration 314(1984)243

Retention behaviour
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography; factors that control retention and selectivity 301(1984)25

Three-mode factor analysis of data on retention in normal-phase HPLC 301(1984)311

Retention times of oximes in reversed-phase HPLC 301(1984)492

Interfacial tension effects of polar modifiers in nonpolar-nonpolar LC 302(1984)79

Basis of retention in normal-phase HPLC with cyano-propyl columns 303(1984)1

Mobile phase effects in reversed-phase chromatography. VII. Dependence of retention on mobile phase composition and column temperature 318(1985)1

HPLC analysis of basic drugs on silica columns using non-aqueous ionic eluents. I. Factors influencing retention, peak shape and detector response 323(1985)173

Retention behavior of long chain quaternary ammonium homologues and related nitroso-alkylmethylamines 324(1985)209

Retention gaps

Speed of temperature increase when using large retention gaps in capillary GC 301(1984)1

Method for testing retention gaps and data on the GC retention power of differently deactivated capillary tubes 323(1985)237

Band broadening in space in splitless injection 324(1985)251

Retention indices

Study of GC behaviour of alkenes based on molecular orbital calculations 302(1984)107

Evaluation of a calibration marker scheme for open tubular column GC with on-column injection and electron-capture detection 302(1984)289

Structure assignment by retention index in gas-liquid radiochromatography of substituted cyclohexenes 312(1984)75

Calculation of retention indices by molecular topology: chlorinated alkanes 314(1984)1

Retention indices of dimethylformamidines, dimethylacetamidines and tetramethylguanidines on a non-polar column 315(1984)95

GLC analyses. XXXI. Retention increments of isomeric chlorophenols on low-polarity (SE-30) and polar (FFAP) capillary columns 315(1984)185

GC investigation of organometallic compounds and their carbon analogues. VI. Determination, calculation and correlation of Kováts retention indices for alkylalkoxysilanes 318(1985)41

Study of Kováts retention indices of aliphatic saturated esters and their relation to the polarity of the stationary phase 318(1985)187

Calculation of retention indices by molecular topology. Chlorinated benzenes 319(1985)1

GC of homologous esters. XXVII. Retention increments of C_1-C_{18} primary alkanols and their 2-chloropropanoyl and 3-chloropropanoyl derivatives on SE-30 and OV-351 capillary columns 319(1985)131

GC of homologous esters. XXVIII. Retention increments of aliphatic C_1-C_{18} *n*-alkyl esters of butanoic acid and its monochloro derivatives on SE-30 and OV-351 capillary columns 320(1985)325

GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

Heats of formation-GC retention relationships for aliphatic hydrocarbons and alkylbenzenes 322(1985)35

GLC analyses. XXXIV. Separation and retention indices with retention increments of some nitrated polynuclear aromatic hydrocarbons on a low-polarity (SE-30) capillary column 322(1985)71

GLC analyses. XXXVII. Capillary column studies of benzoyl and monochlorobenzoyl esters of lower saturated branched-chain alcohols 323(1985)331

GLC analyses. XXXIX. ω-Chloroethanols on low-polarity (SE-30) and polar (OV-351) capillary columns 324(1985)181

GLC analyses. XL. Incremental effects of a formyl group introduced into the *ortho*- and *para*-positions of isomeric chlorophenols 324(1985)192

The use of retention indices to measure the solvent selectivity of ternary eluents in reversed-phase LC 324(1985)243

GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

GC of homologous esters. XXIX. Propanoyl and monochlorpropanoyl esters of lower saturated brached-chain and unsaturated alcohols 324(1985)343

GC separation and identification of linear and branched-chain alkyl bromides 324(1985)363

Calculation of retention indices of pyrazines on the basis of molecular structure 324(1985)428

Structure-retention studies of model ketones by capillary GC 325(1985)13

GC of homologous esters. XXXI. Butanoyl and monochlorobutanoyl esters of lower saturated branched chain and unsaturated alcohols on SE-30 and OV-351 capillary columns 325(1985)61

Chromatography of monomers. II. Glass capillary GC of C_1-C_{18} alkyl esters of acrylic and methacrylic acids 325(1985)367

GLC analyses. XLI. Ethyl and ω-chloroethyl esters of benzoic and monochlorobenzoic acids on SE-30 and OV-351 capillary columns 325(1985)433

Retention mechanisms
Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

Retention models
Influence of water in LSC systems on retention data. II. Fixation of monolocalized solutes with class N solvents and silica gels 316(1984)373

Retention optimization
Retention-optimization strategy for the HPLC ion-pair separation of samples containing basic compounds 316(1984)241

Retention prediction
Reversed-phase LC of homologous series. A general method for prediction of retention 314(1984)13

Prediction of retention data by using parameters of intermolecular interaction. I. A model for calculation of non-polar and polar interaction parameters 315(1984)75

Prediction of retention data by using parameters of intermolecular interaction. II. Application of the model to GLC systems 315(1984)85

Computer-assisted retention prediction system for reversed-phase micro HPLC 316(1984)1

Prediction of retention data of protected dipeptides in normal phase HPLC 325(1985)282

Retention time standards
Polychlorinated biphenyl retention time standards obtained by chemical dechlorination of polychlorinated biphenyl isomers 318(1985)211

Retention volumes
Indirect determination of specific retention volumes in GC. Use of retentions of the McReynolds probes relative to *n*-decane 312(1984)69

Prediction of retention data by using parameters of intermolecular interaction. II. Application of the model to GLC systems 315(1984)85

Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

GC separation and identification of linear and branched-chain alkyl bromides 324(1985)363

Retinals
HPLC of retinals, retinols (vitamin A_1) and their dehydro homologues (vitamin A_2): improvements in resolution and spectroscopic characterization of the stereoisomers 301(1984)65

Retinoids
Analysis of retinoids by HPLC using programmed gradient separation 305(1984)199

Gradient reversed-phase HPLC separation of naturally occurring retinoids 309(1984)299

Stability-indicating reversed-phase HPLC assay for fenretinide drug substances 322(1985)380

Retinol
HPLC of retinals, retinols (vitamin A_1) and their dehydro homologues (vitamin A_2): improvements in resolution and spectroscopic characterization of the stereoisomers 301(1984)65

Determination of serum and plasma concentrations of retinol using HPLC 311(1984)239

Quantitative HPLC method for determining the isomer distribution of retinols (vitamin A_1) and 3-dehydroretinol (vitamin A_2) in fish oils 312(1984)423

Determination of vitamin A and vitamin A acetate by HPLC with fluorescence detection 317(1984)349

Rhamnose
Differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* based on carbohydrates in lipopolysaccharide 310(1984)261

Riboflavin
Simultaneous analysis of thiamin and riboflavin in foods by HPLC 318(1985)412

LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Ribokinase
Enzymatic coupled assay procedures that employ HPLC: the synthesis of orotidylate from ribose 317(1984)245

Ribonuclease
Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Ribose
Enzymatic coupled assay procedures that employ HPLC: the synthesis of orotidylate from ribose 317(1984)245

Ribosomal proteins
Application of HPLC techniques to the separation of ribosomal proteins of different organisms 317(1984)181

Application of HPLC to the reconstitution of ribosomal subunits 317(1984)201

Affinity chromatography of *Drosophila melanogaster* ribosomal proteins to 5S rRNA 319(1985)367

Ribosomal subunits
Application of HPLC to the reconstitution of ribosomal subunits 317(1984)201

Riboxamine
Analysis of riboxamide in urine by HPLC 307(1984)216

Rifampicine
Micro column LC-MS using a capillary interface 323(1985)89

R_M values
R_m values, retention times and π values of a series of potentially mutagenic nitroimidazo[2,1-*b*]thiazoles 320(1985)281

rRNA
Affinity chromatography of *Drosophila melanogaster* ribosomal proteins to 5S rRNA 319(1985)367

tRNA
Temperature gradient chromatography of transfer RNA on hydroxyapatite 314(1984)408

HPLC separations on dihydroxyboryl-agarose 316(1984)301

Mixed-mode chromatographic matrices for the resolution of transfer ribonucleic acids 317(1984)251

Rapid fractionation of tRNA on benzoylated DEAE-porous glass 324(1985)480

Rolling oil
Analysis of mechanical lubricant contaminants in aluminium cold rolling oil by reversed-phase LC 319(1985)222

Rotating disc multiwavelength detector
Use of a rotating disc multiwavelength detector operating in the visible region of the spectrum for monitoring ball pen inks separated by HPLC 303(1984)211

Rotenone
HPLC analysis of rotenone and rotenonone in water by direct injection 303(1984)263

High-efficiency resolution of isomeric rotenone compounds by HPLC 317(1984)383

Rotenonone
HPLC analysis of rotenone and rotenonone in water by direct injection 303(1984)263

Ro 13-6438
Determination of the imidazo quinazoline derivative Ro 13-6438 in biological fluids by HPLC 305(1984)153

Rubidium
Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Λ-Ru(1,10-phenanthroline)$_3^{2+}$-montmorillonite
Clay CC for optical resolution. Resolution of aromatic compounds on a Λ-Ru(1,10-phenanthroline)$_3^{2+}$-montmorillonite column 319(1985)299

Ruthenium
HPLC of homo- and hetero-tetrametallic iron, ruthenium, osmium and nickel clusters 303(1984)404

Ruthenium complexes
Electrophoretic method for the separation and isolation of polyanions and mixed ligand complexes of homologous series. A review with special emphasis on own results (in German) 313(1984)161

S

Saccharides
see also Sugars and Carbohydrates

Determination of fructan oligomers of degree of polymerization 2-30 by HPLC 303(1984)417

Saccharinic acids
Isotachophoretic analysis of saccharinic acids 320(1985)111

Saikosaponin
Quantitative analysis of the metabolites of saikosaponin a using HPLC 307(1984)488

Salazosulphapyridine
Determination of sodium azodisalicylate, salazosulphapyridine and their metabolites in serum, urine and faeces by HPLC 305(1984)470

Salbutamol
Reversed-phase HPLC determination of salbutamol in rabbit plasma 305(1984)485

Determination of salbutamol in human serum by reversed-phase HPLC with amperometric detection 311(1984)311

Salicin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Salicortin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Salicylic acid
Rapid and sensitive determination of acetylsalicylic acid and its metabolites using reversed-phase HPLC 310(1984)343

Improved method for the determination of aspirin and its metabolites in biological fluids by HPLC: applications to human and animal studies 311(1984)301

Simultaneous HPLC assay of acetylsalicylic acid and salicylic acid in film-coated aspirin tablets 317(1984)507

Salicyluric acid
Rapid and sensitive determination of acetylsalicylic acid and its metabolites using reversed-phase HPLC 310(1984)343

Improved method for the determination of aspirin and its metabolites in biological fluids by HPLC: applications to human and animal studies 311(1984)301

Salidrosid
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Salvia garedznii
GLC-MS in the analysis of essential oils 303(1984)109

Sample enrichment
Simple precolumn sample enrichment in HPLC for determination of phenoxy acid herbicides in water samples from exposure studies 319(1985)427

Sample-introduction patterns
Sampling and flow effects in sedimentation field flow fractionation 315(1984)45

Sample preconcentration
Studies on sample preconcentration in ion chromatography. I. Design of an automated, single pump preconcentration system with direct UV absorbance detection 318(1985)279

Sample splitter
Electric sample splitter for capillary zone electrophoresis 320(1985)159

Sample volume
Effect of sample volume on peak width in high-performance gel filtration chromatography 324(1985)422

Sampling device
New injection syringe for GC 322(1985)55

Sapogenins
Analysis of soybean sapogenins by HPLC 325(1985)275

Saponins
HPLC of oleanane saponins 315(1984)441

Sarcosine
Fluorometric evaluation of sarcosine in urine and serum 311(1984)405

Sarin
Analysis of soman and sarin in blood utilizing a sensitive GC-MS method 324(1985)163

Satietin
Molecular weight measurement of human satietin 317(1984)165

Satranidazole
Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325

Schiff base chelates
Volatility and GC of some derivatives of Schiff base chelates 318(1985)350

Schiff base complexes
Polymeric Schiff base complexes as solid phases in GC 322(1985)63

Scintillation fluids
Use of Sep-Pak cartridges for the recovery of compounds from scintillation fluids 325(1985)323

Scopolamine
Qualitative and quantitative analysis of hyoscyamine-atropine and scopolamine in mother tinctures of Solanaceae by HPLC (in French) 319(1985)412

Scoville heat units
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288

Secnidazole
Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325

Sedimentation field flow fractionation
Sampling and flow effects in sedimentation field flow fractionation 315(1984)45

Selectivity
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography; factors that control retention and selectivity 301(1984)25

Influence of substrate parameters on column selectivity with alkylbonded-phase sorbents 316(1984)163

Effect of functional group structure and exchange capacity on the selectivity of anion exchangers for divalent anions 316(1984)201

Selectivity effects between ionic and neutral solutes using hydrophobic pairing ions 316(1984)279

Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23

Optimization in isotachophoresis. The concept of selectivity and separation speed 320(1985)49

Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1

The use of retention indices to measure the solvent selectivity of ternary eluents in reversed-phase LC 324(1985)243

Chromatographic selectivity in LC of polycondensed aromatic hydrocarbons on 3-(2,4-Dinitroanilino)propyl silica 324(1985)261

Sennosides
Comparison between HPTLC fluorometry and HPLC for the determination of sennosides A and B in Senna (Cassia spp.) pods and leaves 303(1984)391

Sensitivity
UV visualization of inorganic anions by reversed-phase ion-interaction chromatography: factors that control sensitivity and detection 316(1984)211

Enhancing isotachophoresis sensitivity by low-concentration electrolyte cascading 320(1985)45

Separation speed
Optimization in isotachophoresis. The concept of selectivity and separation speed 320(1985)49

Sephadex G-10
Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Sephadex G-15
Partitioning of aliphatic alcohols in Sephadex G-15 at 25°C with water as solvent 301(1984)47

Partitioning of oligoglucans in Sephadex G-15 in relation to their conformational structure 318(1985)57

Sephadex G-25
Alkali cation selectivity of Sephadex G-25 in water and aqueous mixtures of methanol, ethanol and *tert.* -butanol 319(1985)247

Sep-Pak Silica
Sorption of organic compounds from urine in mutagenicity testing: choice of sorbent 310(1984)41

Sequential simplex procedure
Automated optimisation of reversed-phase HPLC separations. An improved method using the sequential simplex procedure 316(1984)69

Serine
Quantitation of the enantiomers of β-hydroxyamino acids by GC resolution on an optically active packed column 301(1984)461

Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

Serotonin
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Revised method for the quantitative determination of 5-hydroxytryptamine in human plasma by GC-MS-selected ion monitoring 307(1984)1

Quantification of a number of biogenic amines and their metabolites in brain homogenates by GC-MS 307(1984)241

Simultaneous determination of norepinephrine, dopamine, 5-hydroxytryptamine and their main metabolites in rat brain using HPLC with electrochemical detection. Enzymatic hydrolysis of metabolites prior to chromatography 307(1984)271

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Rapid and simple method for the simultaneous determination of 3,4-dihydroxyphenylacetic acid, 5-hydroxyindole-3-acetic acid and 4-hydroxy-3-methoxyphenylacetic acid in human plasma by HPLC with electrochemical detection 308(1984)55

Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53

Determination of urinary tryptophan, 5-hydroxytryptamine and 5-hydroxyindoleacetic acid in neonatal hyperbilirubinaemic infants using reversed-phase HPLC with fluorescence detection 309(1984)145

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Ascorbic acid oxidase speeds up analysis for catecholamines, indoleamines and their metabolites in brain tissue using HPLC with electrochemical detection 311(1984)167

HPLC determination of hallucinogenic indoleamines with simultaneous UV photometric and voltammetric detection 320(1985)414

Serum proteins
Detection of changes in rabbit serum proteins after partial hepatectomy by means of two-dimensional electrophoresis under non-denaturing conditions 311(1984)93

Fractionation of human serum proteins by immobilized metal affinity chromatography 315(1984)167

Sesquiterpene lactones
New reagent for the detection of sesquiterpene lactones by TLC 303(1984)306

Sesquiterpenes
GC characterization of frequently occuring sesquiterpenes in essential oils 318(1985)125

Settling time
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography. III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

Shale oil
Determination of carbonyl compounds in an Australian (Rundle) shale oil 319(1985)230

Shogaols
HPLC separation of pungency components of ginger 319(1985)471

Siderophores
Separation of ferrichromes and other hydroxamate siderophores of fungal origin by reversed-phase chromatography 301(1984)247

Silanization
Role of solvent and base in the silanization reaction of silicas for reversed-phase HPLC 316(1984)193

Effect of alkyl chain length on surface silanization of silica 318(1985)173

Silanol groups
Characterization of silanol reactivity and acidity on octadecyl-bonded chromatographic supports by ^{29}Si solid-state nuclear magnetic resonance spectroscopy and surface titration 319(1985)9

Silica-bound polyethyleneglycol
Silica-bound polyethyleneglycol as stationary phase for separation of proteins by HPLC 319(1985)396

Silica gel
Comparison of chromatographic behaviour of silica and chemically bonded phases in supercritical fluid chromatography 315(1984)31

Influence of substrate parameters on column selectivity with alkylbonded-phase sorbents 316(1984)163

Silicalite
Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333

Silicic acid
Chromatographic studies in the field of silicate chemistry (Review) 313(1984)33

Silvex
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Simazine
Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Simplex procedure
Automated optimisation of reversed-phase HPLC separations. An improved method using the sequential simplex procedure 316(1984)69

Sinapic acid
Direct coupling of packed fused-silica LC columns to a magnetic sector mass spectrometer and application to polar thermolabile compounds 323(1985)47

Sisomicin
Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Determination of sisomicin, netilmicin, astromicin and micronomicin in serum by HPLC 305(1984)373

Size-exclusion chromatography
Theoretical models for size-exclusion chromatography and calculation of pore size distribution from size-exclusion chromatography data 316(1984)311

Semi-micro size-exclusion chromatography of polymers. Calibration of columns and comparison with conventional columns 320(1985)343

Skatole
GC determination and MS confirmation of traces of indole and 3-methylindole (skatole) in pig back fat 324(1985)469

Snyder equation
Three-mode factor analysis of data on retention in normal-phase HPLC 301(1984)311

Sobrerol
Sensitive GLC-electron-capture detection method for determination of sobrerol in biological fluids 308(1984)189

Sodium
Isotachophoresis as a candidate reference method in analytical chemistry. Determination of sodium in serum 320(1985)193

Simultaneous determination of sodium, ammonium, potassium, rubidium, caesium, magnesium and calcium ions by a conventional HPLC system with a common ion-exchange column 323(1985)443

Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Sodium alkanesulphonates
Studies in reversed-phase ion-pair chromatography. IV. The role of the chain length of the pairing ion 303(1984)29

Sodium dodecyl sulphate
Effect of sodium dodecyl sulphate on the extraction of ubiquinone-10 in the determination of plasma samples 310(1984)204

Sodium ethylenediaminetetraacetatocobaltate(III)
Substitution-inert metal complex mobile phases in non-suppressed anion chromatography 312(1984)99

Sodium octanesulphonate
Rapid optimization of the concentration of the ion-pairing reagent in ion-pairing reversed-phase LC 316(1984)231

Soluble antibody affinity chromatography
Soluble antibody affinity chromatography technique investigated with ultratrace [^{125}I]thyroxine 314(1984)295

Solution behaviour
Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

Solvents
Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

Solvent selectivity
Solvent selectivity studies using isomers of polystyrene 314(1984)55

Solvophobic effects
Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1

Solvophobic theory
Salt-mediated retention of proteins in hydrophobic interaction chromatography. Application of solvophobic theory 317(1984)67

Soman
Analysis of soman and sarin in blood utilizing a sensitive GC-MS method 324(1985)163

Sorbents
Sorption of organic compounds from urine in mutagenicity testing: choice of sorbent 310(1984)41

Sorbic acid
Determination of preservatives in cosmetic products by reversed-phase HPLC 315(1984)465

Sorption
GC solute-sorbent interactions on macroporous methacrylate ion exchangers 324(1985)277

Sorption isotherms
Influence of additives to the eluent on hydrophobic interaction chromatography of simple compounds. I. Sorption isotherms of *n*-alcohols on octyl-agarose 322(1985)399

Sotalol
Determination of the beta-adrenoceptor blocking drug sotalol in plasma and tissues of the rat by HPLC with UV detection 309(1984)187

Spacers
Introduction of ω-carboxyl spacers onto cross-linked agarose gel beads by O-alkylation for the preparation of affinity adsorbents 315(1984)161

Specific retention volumes
Indirect determination of specific retention volumes in GC. Use of retentions of the McReynolds probes relative to *n*-decane 312(1984)69

Speed of analysis
Effect of high speed on peak capacity in LC 316(1984)81

Theory of the efficiency of serially connected GC columns and the effect on optimized speed of analysis 325(1985)1

Spermidine
Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59

Spermine
Analysis of dansyl derivatives of di- and polyamines in mouse brain, human serum and duodenal biopsy specimens by HPLC on a standard reversed-phase column 311(1984)59

Spheron
Ion-exchange derivatives of Spheron. V. Sulphate and sulpho derivatives 301(1984)93

Sphingomyelin
Quantitative densitometric method for the rapid separation and quantitation of the major tissue and lipoprotein lipids by HPTLC. I. Sample preparation, chromatography, and densitometry 307(1984)65

HPLC of phospholipids using deuterated solvents for IR detection 307(1984)261

Improved HPLC of sphingomyelin 322(1985)240

Spiroplatin
Analysis of antitumour [1,1-bis(aminomethyl)cyclohexane]platinum(II) complexes derived from spiroplatin by HPLC with differential pulse amperometric detection 320(1985)379

Spiropyran-end-labelled polystyrenes
High-performance liquid-solid adsorption chromatography of spiropyran-end-labelled polystyrenes 303(1984)361

Splitter
Electric sample splitter for capillary zone electrophoresis 320(1985)159

Spray reagents
New reagent for the detection of sesquiterpene lactones by TLC 303(1984)306

Sulphosalicylic acid as spray reagent for the detection of sugars on thin-layer chromatograms 315(1984)401

Use of π-acceptors as spray reagents for the detection of penicillins on thin-layer plates 323(1985)424

Spreading pressure
GC determination of adsorption isotherms, spreading pressures, London force interactions and equations of state for *n*-alkanes on graphite and carbon blacks 324(1985)19

Squalene
Enhanced response of more highly unsaturated lipids on TLC-Iatroscan flame ionization detection after exposing the developed Chromarods to iodine vapour 319(1985)215

Square-planar complexes
Effect of the geometric configuration of square-planar complexes on their R_F values obtained by TLC on silica gel 324(1985)431

Stability constants
Complex-forming equilibria in isotachophoresis. VI. Simulation of isotachophoretic equilibria of lanthanoids and determination of mobilities and stability constants of acetate and α-hydroxyisobutyrate complexes 312(1984)11

Staphidine
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Staphisine
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Starch
Rapid analysis of starch, amylose and amylopectin by high-performance size-exclusion chromatography 319(1985)205

Starvation-refeeding experiments
Effects of starvation and refeeding on the excretion of urinary steroid metabolites in mice with different genetic background 308(1984)1

Stationary phase dielectric constant
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433

Stationary phases
see also Chiral stationary phases

Stationary phases, GC
Use of poly(silarylene-methylphenylsiloxane) block copolymer as a thermostable stationary phase in capillary column GC 301(1984)265

Selection of GC stationary phase pairs for characterization of the 1-octanol-water partition coefficient 301(1984)355

GC-MS analysis of compounds generated upon thermal degradation of some stationary phases in capillary GC 302(1984)95

Polarizable polysiloxane stationary phases for capillary column GC 302(1984)303

Cyanosilicones as stationary phases in GC. III. Synthesis, characterization and evaluation 302(1984)319

Indirect determination of specific retention volumes in GC. Use of retentions of the McReynolds probes relative to *n*-decane 312(1984)69

Liquid crystalline stationary phases with a cyclohexyl ring in the molecule 312(1984)141

Disc-like liquid crystalline stationary phases from the triphenylene derivatives group 315(1984)145

Separation of underivatised barbiturates by capillary column GC. Preparation of medium polar polymethylsiloxane columns to optimize selectivity 318(1985)255

Aryl substituted diastereomeric alkenes: GC behavior on a non-polar *versus* a liquid crystal phase 321(1985)127

Polymeric Schiff base complexes as solid phases in GC 322(1985)63

Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

Distribution of the liquid crystal stationary phase on the surface of diatomite supports 324(1985)299

Phase transitions of 4-cyano-4'-*n*-alkoxyformyloxyazobenzenes on silanized supports 324(1985)309

Stationary phases, LC
Unmodified and acylated cyclodextrin stationary phases for LC separation of aromatic compounds 301(1984)345

Separation of polychlorinated biphenyls (Aroclor 1254) by HPLC 302(1984)159

HPLC of neuropeptides using radially compressed polythene cartridges 306(1984)99

Separation of peptides on a polystyrene resin column 307(1984)29

Determination of creatinine in human, dog and rat urine by HPLC on a column of hydroxymethylated porous polystyrene 307(1984)399

Separation of collagen α-chains by reversed-phase HPLC. Comparison of column alkyl stationary phases and temperature effects 308(1984)111

Preparation and retention behaviour of chemically bonded methylated-cyclodextrin stationary phases for LC 314(1984)193

Influence of substrate parameters on column selectivity with alkylbonded-phase sorbents 316(1984)163

Role of solvent and base in the silanization reaction of silicas for reversed-phase HPLC 316(1984)193

HPLC separations on dihydroxyboryl-agarose 316(1984)301

Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333

Wide-pore silica-based ether-bonded phases for separation of proteins by high-performance hydrophobic-interaction and size-exclusion chromatography 316(1984)519

Hydrophobic interaction HPLC of proteins 317(1984)55

Reversed-phase chromatography of proteins with diphenyl-silica column and hydro-organic eluents containing two organic solvents 318(1985)11

Simultaneous selectivity optimization of mobile and stationary phases in reversed-phase LC for isocratic separations of phenylthiohydantoin amino acid derivatives 318(1985)23

Stationary phase contributions to retention in high-performance anion-exchange protein chromatography: ligand density and mixed mode effects 318(1985)157

Effect of alkyl chain length on surface silanization of silica 318(1985)173

Sepharose 6B substituted with polyethylene oxide-like derivatives for chromatographic fractionation of protein mixtures 319(1985)43

Sepharose derivatives containing citric acid as affinity ligand. Purification of fumarase 319(1985)173

Cibacron Blue 3G-A-substituted cross-linked agar beads: an inexpensive, fast-flow affinity medium for large scale applications 319(1985)218

Pyrolysis-MS investigations of reversed-phase HPLC stationary phases 319(1985)285

Silica-bound polyethyleneglycol as stationary phase for separation of proteins by HPLC 319(1985)396

Comparison of phenyl- and octadecyl-modified silica gel as stationary phase for the prediction of *n*-octanol-water partition coefficients by HPLC 320(1985)271

High-performance gel filtration of water-soluble samples on polyvinyl alcohol columns 320(1985)369

Separation of large DNA restriction fragments by high-performance gel filtration on TSKgel DNA-PW 320(1985)440

Characterization of silica-based reversed-phase columns with respect to retention selectivity. Solvophobic effects 321(1985)1

α-Aminoacyl hydroxamate adsorbents: a new type of immobilized chelator 321(1985)81

Fe^{3+}-Hydroxamate as immobilized metal affinity-adsorbent for protein chromatography 321(1985)93

Separation of serum proteins on a Fe^{3+}-monohydroxamate adsorbent 321(1985)105

Synthesis of new hydrophobic adsorbents based on homologous series of uncharged alkyl sulphide agarose derivatives 321(1985)305

Carbon analysis for the study of bonded phases in LC 321(1985)319

Quantitative determination of the ligand content and the negative and positive charges on butylamine-substituted Sepharose 4B 321(1985)335

Structural investigations on reversed-phase silicas. Determination of ligand functionality 323(1985)273

Synthesis, solid-state nuclear magnetic resonance characterization and chromatographic evaluation of a diphenylphosphine-octadecylsilane hybrid bonded phase 323(1985)281

Resolution of racemic aminoalcohols (β-blockers), amines and acids as enantiomeric derivatives using a chiral α_1-acid glycoprotein column 325(1985)379

Stationary phases, supercritical fluid chromatography
Comparison of chromatographic behaviour of silica and chemically bonded phases in supercritical fluid chromatography 315(1984)31

Stationary phases, TLC
Comparison of methods for separating polycyclic aromatic hydrocarbons by HPTLC 302(1984)149

Factors affecting the ion-pair reversed-phase TLC of organic acids on paraffin-coated and C_{18} bonded silica gel 312(1984)133

Stationary phase structure
Characterization of silanol reactivity and acidity on octadecyl-bonded chromatographic supports by ^{29}Si solid-state nuclear magnetic resonance spectroscopy and surface titration 319(1985)9

Steam gas-solid chromatography
Steam GSC: a flexible separation technique 301(1984)11

Stearic acid
Electrokinetic detection of lauric, myristic, stearic and arachidic acids separated by reversed-phase HPLC 321(1985)433

Steric effects
GC of homologous esters. XXX. Unexpected predominancy of polar effects over steric effects in crowded aliphatic esters: a GLC appraisal of specific interactions 324(1985)355

Steroid glucuronates
Rapid chromatographic purification of urinary steroid glucuronates for determination of aldosterone secretion rate 317(1984)367

Steroid receptors
High-performance size exclusion chromatography as a rapid method for the separation of steroid hormone receptors 307(1984)39

Steroids
see also Hormones

Identification of meibomian gland lipids by GC-MS: application to the meibomian lipids of the mouse 301(1984)173

Use of steroid profiling in the resolution of pregnenolone metabolites from porcine testicular preparations 301(1984)189

Analysis of flumethasone pivalate formulations by HPLC 301(1984)477

Computer simulation as an aid to optimizing continuous-development two-dimensional TLC 302(1984)135

Solvent optimization of reversed-phase HPLC for polar adrenal steroids using computer-predicted retentions 305(1984)13

New fluorimetric determination of 17-hydroxycorticosteroids after HPLC using post-column derivatization with benzamidine 305(1984)188

Application of a high-performance gel permeation LC procedure to the determination of binding of prednisolone to high-affinity binding sites in human serum 305(1984)194

Semi-micro quantitative analysis of complex urinary steroid mixtures in healthy and diseased states 306(1984)39

Simultaneous analysis of cortexolone and cortisol by HPLC for use in the metyrapone test 306(1984)345

Studies on steroids. CC. Determination of 17-ketosteroid sulphates in serum by HPLC with electrochemical detection using pre-column derivatization 307(1984)23

Effects of starvation and refeeding on the excretion of urinary steroid metabolites in mice with different genetic background 308(1984)1

Quantitative metabolic profiling of testicular steroid secretions with bonded-phase capillary GC 308(1984)11

Rapid analysis of C_{19}-steroid metabolism by HPLC and in-line monitoring of radioactivity 308(1984)289

Application of reversed-phase HPLC for radioimmunoassay of plasma 18-hydroxycorticosterone 308(1984)295

Separation of oestrogen conjugates in urine and synthetic mixtures by HPLC methods (Review) 309(1984)1

Determination of methylprednisolone and methylprednisolone acetate in synovial fluid using HPLC 309(1984)385

Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Rapid measurement of oestradiol and oestriol by HPLC after automatic pretreatment 311(1984)249

Determination of trilostane and ketotrilostane in human plasma by HPLC 311(1984)434

Reversed-phase HPLC for the determination of steroid hormones in oral contraceptives 315(1984)313

Qualitative detection of corticosteroids in equine biological fluids and the comparison of relative dexamethasone metabolite/dexamethasone concentration in equine urine by micro-LC-MS 315(1984)359

General reversed-phase HPLC procedure for the analysis of oral contraceptive formulations 315(1984)470

Analysis of steroids in bulk pharmaceuticals by LC with light-scattering detection 316(1984)461

Determination of trace levels of steroids in blood plasma by LC with peroxyoxalate chemiluminescence detection 317(1984)355

Clinical application of the rapid HPLC determination of serum cortisol 317(1984)377

HPLC determination of ecdysteroid titers in the house fly 317(1984)413

Laser fluorimetry for capillary column LC: high-sensitivity detection of derivatized biological compounds 317(1984)443

TLC of ecdysteroids 318(1985)373

Fast HPLC screening method for the presence of trenbolone and its major metabolite in urine of slaughter cattle 319(1985)436

Determination of 9α-hydroxyandrost-4-en-3,17-dione in fermentation solutions by TLC (in German) 319(1985)475

Analytical droplet counter-current chromatography isolation of 20-hydroxyecdysone from *Vitex thyrsiflora* (Verbenaceae) 321(1985)246

HPTLC on amino-bonded silica gel: application to barbiturates and steroids 322(1985)149

Isocratic reversed-phase HPLC of free and sulphoconjugated steroids 322(1985)371

Steroid sulphates
Isocratic reversed-phase HPLC of free and sulphoconjugated steroids 322(1985)371

Stevioside
Role of column switching in semi-preparative LC. Isolation of the sweetener stevioside 316(1984)105

Stirocainide
Assay of the antiarrhythmic compound stirocainide in plasma by fused-silica GLC and nitrogen-selective detection 307(1984)190

Strontium
Characterization of enzymatic hydrolysis of inorganic polyphosphates by flow injection analysis and HPLC 325(1985)385

Structure-activity relationships
Comparison of phenyl- and octadecyl-modified silica gel as stationary phase for the prediction of *n*-octanol-water partition coefficients by HPLC 320(1985)271

R_m values, retention times and π values of a series of potentially mutagenic nitroimidazo[2,1-*b*]thiazoles 320(1985)281

Structure-retention relationships
Structure assignment by retention index in gas-liquid radiochromatography of substituted cyclohexenes 312(1984)75

Calculation of retention indices by molecular topology: chlorinated alkanes 314(1984)1

Reversed-phase retention behaviour of Dns-amides as a function of eluent composition and molecular structure in ethanol-water and methanol-water 314(1984)141

TLC behaviour and chemical structure of 6- and 8-methoxy-5-hydroxyflavones 315(1984)101

GC investigation of organometallic compounds and their carbon analogues. VI. Determination, calculation and correlation of Kováts retention indices for alkylalkoxysilanes 318(1985)41

Structure-retention relationships: chromatographic behaviour and properties of nucleotide-cation complexes 318(1985)221

Solvated structure-retention relationships of peropyrene-type polycyclic aromatic hydrocarbons 322(1985)275

Calculation of retention indices of pyrazines on the basis of molecular structure 324(1985)428

Effect of the geometric configuration of square-planar complexes on their R_F values obtained by TLC on silica gel 324(1985)431

Structure-retention studies of model ketones by capillary GC 325(1985)13

Styrene oligomers
Phenyl-bonded phase LC of styrene oligomers 312(1984)313

Substance P
Ion-exchange chromatographic assay of peptidases acting on the C-terminal hexapeptide sequence of substance P 301(1984)415

Substituent effects
GLC analyses. XXXII. Incremental effects of hydroxy, methoxy and acetoxy groups introduced into isomeric chlorobenzenes 321(1985)115

GLC analyses. XL. Incremental effects of a formyl group introduced into the *ortho-* and *para-*positions of isomeric chlorophenols 324(1985)192

Substitution-inert metal complexes
Substitution-inert metal complex mobile phases in non-suppressed anion chromatography 312(1984)99

Succinimides
Application of HPLC chiral stationary phases to pharmaceutical analysis. Resolution of enantiomeric barbiturates, succinimides and related molecules on four commercially available chiral stationary phases 324(1985)444

Sucrose
HPLC separation of glucose-1-phosphate, fructose, sucrose and inorganic orthophosphate 322(1985)220

Micro column LC-MS using a capillary interface 323(1985)89

Sugars
see also Carbohydrates and saccharides

Automated anion-exchange chromatographic method for the quantitation of lactulose in physiological fluids and processed milks 307(1984)393

Whole-cell methanolysis as a rapid method for differentiation between *Actinobacillus actinomycetemcomitans* and *Haemophilus aphrophilus* 311(1984)347

Capillary GC of partially methylated alditol acetates on a high-polarity bonded-phase vitreous-silica column 315(1984)373

Sulphosalicylic acid as spray reagent for the detection of sugars on thin-layer chromatograms 315(1984)401

Single-valve, single-pump, HPLC column-switching analysis of hydrolyzed wood components 318(1985)269

Sulphadoxine
Quantification of antimalarial drugs. I. Simultaneous measurement of sulphadoxine, N_4-acetylsulphadoxine and pyrimethamine in human plasma 305(1984)502

HPLC assay of pyrimethamine, sulfadoxine and its N^4-acetyl metabolite in serum and urine after ingestion of Suldox 308(1984)217

Sulphamethazine
Simultaneous analysis of three antimicrobial agents in feed premixes by reversed-phase HPLC 323(1985)447

Sulphamethoxazole
HPTLC determination of trimethoprim and sulphamethoxazole in pharmaceutical dosage forms 323(1985)456

Sulphamethoxypyridazine
Simultaneous analysis of three antimicrobial agents in feed premixes by reversed-phase HPLC 323(1985)447

Sulphanilamides
Determination of chloroacetanilides and chloroanilines in sulphanilamides and waste water by HPLC 319(1985)99

Sulphate
Determination of anions and cations in atmospheric aerosols by single column ion chromatography 324(1985)153

Sulphides
Reaction GC-MS. VI. Use of deuterium as the carrier and reagent gas in reaction GC-MS 301(1984)77

Sulphinpyrazone
Simple and rapid HPLC method for the analysis of sulfinpyrazone and four of its metabolites in human plasma 310(1984)438

Sulphite
Determination of sulphite in fixers and photographic effluents by ion chromatography 322(1985)478

Sulphoconjugated steroids
Isocratic reversed-phase HPLC of free and sulphoconjugated steroids 322(1985)371

Sulphonates
Separation of sulfonate and carboxylate mixtures by ion-exchange HPLC 302(1984)65

Sulphonic acids
Reversed-phase ion-pair chromatography of organic acids on silica with cetrimide as cationic ion-pair reagent 316(1984)267

Sulphosalicylic acid
Sulphosalicylic acid as spray reagent for the detection of sugars on thin-layer chromatograms 315(1984)401

Sulphotransferase
Rapid and sensitive assays using HPLC to measure the actitivies of phase II drug metabolising enzymes: glucuronyl transferase and sulfotransferase 301(1984)282

Sulphoxides
GC separation of enantiomeric sulphur compounds on Chirasil-Val 320(1985)393

Sulphur
Retention of sulphur and sulphur organics in reversed-phase LC 317(1984)589

Sulphur-containing compounds
GC separation of enantiomeric sulphur compounds on Chirasil-Val 320(1985)393

GC study of the analysis and elution mechanism of hydrogen sulphide, carbonyl sulphide and light mercaptans in petroleum gases 321(1985)59

Sulphydryl esterase
Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

Supercritical fluid chromatography
Supercritical fluid chromatography and supercritical fluid chromatography-MS of marine diesel fuel 314(1984)253

Thermodynamic model for supercritical fluid chromatography 315(1984)1

Chromatography with sub- and supercritical eluents. The influence of temperature, pressure and flow-rate on the behaviour of dimethyl and diethyl ether 315(1984)19

Comparison of chromatographic behaviour of silica and chemically bonded phases in supercritical fluid chromatography 315(1984)31

Supercritical fluid chromatography of various samples with different polarities 315(1984)39

Chromatography with sub- and supercritical eluents: influence of temperature, pressure and flow-rate on the behaviour of lower alkanes 321(1985)273

Superoxide dismutase
Comparison of high-performance ion-exchange chromatography and gel electrophoresis in protein separations 318(1985)358

Surface effects
Evidence for surface effects in an electron-capture detector 314(1984)219

Surfactants
Effect of surfactants on the separation of proteins by reversed-phase HPLC. I. Non-ionic surfactants (Tween) 317(1984)157

On-line LC-MS analysis of non-ionic surfactants 323(1985)135

Sweeteners
Isocratic LC method for the simultaneous determination of aspartame and other additives in soft drinks 303(1984)256

Role of column switching in semi-preparative LC. Isolation of the sweetener stevioside 316(1984)105

Synaptic plasma membranes
Affinity chromatography of membrane fragments. Separation of carbonic anhydrase-enriched and -depleted brain membranes 312(1984)435

T

2,4,5-T
Micro post-column extraction system for interfacing reversed-phase micro LC and MS 312(1984)153

Tandem mass spectrometry
Determination of dibenzothiophene in oils by LC-tandem MS 323(1985)33

Tannins
Fractionation of condensed tannins by counter-current chromatography 318(1985)85

Tartaric acid monoesters
LC separation of enantiomeric alkanolamines via diastereomeric tartaric acid monoesters 316(1984)605

***d*-Tartrate**
Mechanism of chromatographic separation of optically active metal complexes (Review) 313(1984)59

Taurine
Quantitative GLC of taurine 306(1984)61

Taurocyamine
HPLC measurement of guanidino compounds of clinical importance in human urine and serum by pre-column fluorescence derivatization using benzoin 311(1984)257

Tea
Isolation of trypsin by column chromatography on tea particles 315(1984)478

Temperature effects
Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

Temperature programming
Speed of temperature increase when using large retention gaps in capillary GC 301(1984)1

Temperature gradient chromatography of transfer RNA on hydroxyapatite 314(1984)408

Tenoxicam
Rapid method for the determination of either piroxicam or tenoxicam in plasma using HPLC 310(1984)455

Determination of tenoxicam, and the isolation, identification and determination of Ro 17-6661, its major metabolite, in human urine 317(1984)483

Terazosin
Rapid and sensitive analysis of terazosin in plasma, peritoneal dialysis solution, and urine using HPLC with fluorescence detection 311(1984)206

Terpenoid alcohols
Investigation of isoprenoid benzoates and naphthoates by reversed-phase LC. Isocratic elution characteristics of benzoates and naphthoates of C_5-C_{20} terpenoid alcohols 312(1984)121

Terpenoids
GLC-MS in the analysis of essential oils 303(1984)109

Improved procedure for the TLC of terpenoids on silver ion-silica gel layers 320(1985)455

Testosterone
Simultaneous determination of fifteen steroid hormones from a single serum sample by HPLC and radioimmunoassay 310(1984)386

Testosterone enanthate
HPLC analysis of estradiol valerate-testosterone enanthate in oily formulations 301(1984)292

Tetrabutylammonium salicylate
Methods for the analysis of inorganic anions. IV. Reversed-phase HPLC with aqueous hydrophobic ion pairs as eluents 302(1984)15

Tetracaine
Simultaneous determination of chlorhexidine, tetracaine and their degradation products by ion-pair LC 315(1984)457

Tetrachloroquinone
Masking effects in "charge-transfer" TLC 303(1984)190

Tetracyanoethylene
Masking effects in "charge-transfer" TLC 303(1984)190

Tetracyclines
Improvement of chemical analysis of antibiotics. VII. Comparison of analytical methods for determination of impurities in tetracycline pharmaceutical preparations 314(1984)303

Reversed-phase HPLC separations of tetracycline derivatives using volatile mobile phases 321(1985)462

Improvement of chemical analysis of antibiotics. VIII. Application of prepacked C_{18} Cartridge for the analysis of tetracycline residues in animal liver 325(1985)265

Tetraethylenepentamine
Separation and analysis of ethylenediamines as their *tert.* -butyldimethylsilyl derivatives by GLC and MS 314(1984)455

Tetrahydrobiopterin
Determination of quinonoid dihydrobiopterin by LC and electrochemical detection 317(1984)407

Tetrahydrocannabinol
Detection and quantitation of tetrahydrocannabinol in serum using TLC and fluorometry 307(1984)493

Comparison of the analysis of Δ^9-tetrahydrocannabinol capsules by HPLC and capillary GC 317(1984)455

Δ^9-Tetrahydrocannabinol-11-oic acid
Combination of LC with UV detection and GC with electron-capture detection for the determination of Δ^9-tetrahydrocannabinol-11-oic acid in urine 306(1984)183

Tetrahydrofuran
Automated GC method for the determination of residual solvents in bulk pharmaceuticals 321(1985)137

1-(Tetrahydro-2-furanyl)-5-fluorouracil
Fluorescence determination of 5-fluorouracil and 1-(tetrahydro-2-furanyl)-5-fluorouracil in blood serum by HPLC 310(1984)151

Tetrahydrouridine
Analysis of candidate anticancer drugs by thermospray HPLC-MS 323(1985)383

Tetraiodofluorescein
Metal-ion mediated interaction of luciferase with tetraiodofluorescein 315(1984)175

Tetrametallic clusters
HPLC of homo- and hetero-tetrametallic iron, ruthenium, osmium and nickel clusters 303(1984)404

Tetramethylguanidines
Retention indices of dimethylformamidines, dimethylacetamidines and tetramethylguanidines on a non-polar column 315(1984)95

Tetraphenylarsonium
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433

Tetraphenylarsonium chloride
Non-aqueous LC of some metalloporphyrins on a reversed-phase column 315(1984)111

Tetraphenylarsonium perchlorate
Non-aqueous LC of some metalloporphyrins on a reversed-phase column 315(1984)111

Tetraphenylborate
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433

Tetraphenylsilane
Establishment and determination of interfacial potentials and stationary phase dielectric constant in reversed-phase LC 322(1985)433

Thebaine
Isolation, separation and detection via HPLC of acidic and neutral acetylated rearrangement products of opium alkaloids 317(1984)427

Theobromine
Simultaneous determination of caffeine and its N-demethylated metabolites in umbilical cord plasma using HPLC 306(1984)191

Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107

Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Use of microbore HPLC for the determination of caffeine, theobromine and theophylline in cocoa 318(1985)408

Theophylline
Simultaneous determination of caffeine and its N-demethylated metabolites in umbilical cord plasma using HPLC 306(1984)191

Therapeutic drug monitoring using high-speed LC and rapid sample preparation: an assay for serum theophylline 308(1984)354

Routine determination of eight common anti-epileptic drugs and metabolites by HPLC using a column-switching system for direct injection of serum samples 310(1984)97

Automated HPLC assay for monitoring caffeine and its metabolites in biological fluids of monkeys consuming caffeine 310(1984)107

Computer-aided strategies for archive retrieval and sensitivity enhancement in the identification of drugs by photodiode array detection in HPLC 316(1984)423

Use of microbore HPLC for the determination of caffeine, theobromine and theophylline in cocoa 318(1985)408

Alkylation with dialkylsulfate and diisopropylethylamine 319(1985)382

Thermal desorption
Preliminary evaluation of thermal desorption-GC analysis of airborne particulate matter on dichotomous filters 303(1984)53

Thermionic detector
New thermionic detector utilizing orthogonal nebulization for capillary column LC 314(1984)103

Thermodynamic functions
Linear relationship in the thermodynamics of GLC 321(1985)458

Thermoset resins
Thermoset resin microstructure by GPC: an anhydride-cured epoxy 316(1984)343

Thermospray interface
Ten years of LC-MS 323(1985)3

Developments in interfacing microbore HPLC with MS (a Review) 323(1985)99

Studies with a laboratory-constructed thermospray liquid chromatograph-mass spectrometer interface 323(1985)127

Thermospray ionization
LC-MS interfacing on a magnetic sector mass spectrometer. Techniques and applications 323(1985)153

Applications of thermospray ionisation LC-MS to compounds of pharmaceutical interest 323(1985)163

Analysis of candidate anticancer drugs by thermospray HPLC-MS 323(1985)383

Thiamine
Determination of thiamine and thiamine phosphates in excitable tissues as thiochrome derivatives by reversed-phase HPLC on octadecyl silica 307(1984)283

Simultaneous analysis of thiamin and riboflavin in foods by HPLC 318(1985)412

Thiazides
HPLC behaviour of some pharmaceutically important thiazide, loop and potassium-sparing diuretics 325(1985)395

Thiocyanate
Determination of cyanide and thiocyanate in blood plasma and red cells by HPLC with fluorometric detection 308(1984)133

Butylation of inorganic anions for simultaneous determination by GC 319(1985)143

Thiols
Electrophoresis of thiols in cellulose gels. III. Group analysis of urinary thiols 307(1984)416

Thiopental
Rapid GC assay for serum thiopental 305(1984)210

HPLC method for methohexital and thiopental in plasma or whole blood 307(1984)481

Thiophene
Nickel(II)-exchanged molecular sieve 13X as a column packing for the GC determination of thiophene in benzene with flame ionization detection 325(1985)430

Thioridazine
Assay of plasma thioridazine and metabolites by HPLC with amperometric detection 307(1984)457

Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Thiothixene
Improved HPLC method for the quantitation of *cis*-thiothixene in plasma samples using *trans*-thiothixene as internal standard 311(1984)424

Thitsiol
Analysis of long-chain phenols in the sap of the Burmese lac tree, *Melanorrhoea usitate* by capillary GLC 318(1985)378

Three-mode factor analysis
Three-mode factor analysis of data on retention in normal-phase HPLC 301(1984)311

Threonine
Quantitation of the enantiomers of β-hydroxyamino acids by GC resolution on an optically active packed column 301(1984)461

Application of reversed-phase HPLC using pre-column derivatization with *o*-phthaldialdehyde for the quantitative analysis of amino acids in adult and fetal sheep plasma, animal feeds and tissues 315(1984)321

Thromboxanes
Determination of 6-keto-prostaglandin $F_{1\alpha}$ and thromboxane B_2 in urine by HPLC and radioimmunoassay 306(1984)351

Rapid, sensitive method for detecting different arachidonic acid metabolites by TLC: the use of autoradiography 307(1984)185

Picogram detection of eicosanoids by UV absorbance after narrow-bore HPLC. Comparison with conventional-bore column 308(1984)31

Simple single-step HPLC method for the separation of cyclooxygenase and lipoxygenase enzyme metabolites of arachidonic acid 308(1984)316

Thylakoid membrane
Fractionation of thylakoid membrane components by extraction in aqueous polymer two-phase systems containing detergent 323(1985)363

Thymine
Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201

Thymosin β_4
Isolation and structural studies of porcine, ovine and murine thymosin β_4 by HPLC 301(1984)221

Thyroglobulin
Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395

Thyroxine
Soluble antibody affinity chromatography technique investigated with ultratrace [^{125}I]thyroxine 314(1984)295

Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395

Tiapamil
Determination of tiapamil and of its two main metabolites in plasma and in urine by HPLC 310(1984)119

Tiaprofenic acid
HPLC analysis of tiaprofenic acid and its metabolites in plasma and urine by direct injection 310(1984)327

Tiazofurin
Quantification of tiazofurin in plasma by HPLC 307(1984)361

Timolol
Determination of timolol in plasma and breast milk using HPLC with electrochemical detection 305(1984)244

Tinidazole
Electron-capture GC assays of 5-nitroimidazole class of antimicrobials in blood 305(1984)325

Tinuvin 622
Determination of polymeric hindered-amine light stabilisers in plastics by pyrolysis-GC 325(1985)87

Tiron
Determination of molybdenum and tungsten with tiron by reversed-phase LC 320(1985)403

Tobramycin
Automated HPLC method for the determination of aminoglycosides in serum using pre-column sample clean-up and derivatization 305(1984)345

Pre-column derivatization of aminoglycosides with 1-fluoro-2,4-dinitrobenzene 322(1985)321

Tocainide
Simultaneous determination of the enantiomers of tocainide in blood plasma using GLC with electron-capture detection 306(1984)155

Monitoring of *S*- and *R*-tocainide in human plasma after heptafluorobutyrylation, separation on Chirasil-Val and electron-capture detection 308(1984)181

α-Tocopherol
Determination of α-tocopherol in human plasma by HPLC with electrochemical detection 311(1984)176

Tocopherols
Simultaneous determination by HPLC of tocopherol isomers, α-tocopheryl quinone, and cholesterol in red blood cells and plasma 306(1984)371

α-Tocopheryl quinone
Simultaneous determination by HPLC of tocopherol isomers, α-tocopheryl quinone, and cholesterol in red blood cells and plasma 306(1984)371

Toloxatone
Reversed-phase HPLC separation of ^{14}C-labelled toloxatone and its metabolites 311(1984)329

Toluenediisocyanate
Detection of urinary amine metabolites in toluene diisocyanate exposed rats 323(1985)429

Toluene-4-sulphonic acid
Polyion ion-pair LC. I. A new method for the regulation of the retention of ionic compounds in LC 314(1984)127

Toluidine
Unmodified and acylated cyclodextrin stationary phases for LC separation of aromatic compounds 301(1984)345

Tomato paste serum
LC of tomato paste serum with sample preparation by ultrafiltration 325(1985)450

Toxins
Rapid determination of the neurotoxin lolitrem B in perennial ryegrass by HPLC with fluorescence detection 321(1985)217

Fast protein LC for the isolation of *Clostridium perfringens* type A α-toxin 321(1985)235

HPLC with stop-flow UV spectral characterization of lolitrem neurotoxins from perennial ryegrass 322(1985)159

Trace analysis
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Trace analysis of amines and isocyanates using glass capillary GC and selective detection. III. Determination of aliphatic and alicyclic amines as perfluoro fatty acid amides using electron-capture and nitrogen-selective detection 303(1984)89

Derivatization at the liquid-solid interface. Application to the trace analysis of organic acids in aqueous samples 314(1984)421

Large sample volume preseparation for trace analysis in isotachophoresis 320(1985)89

Trace enrichment
Optimization of precolumn design in LC 301(1984)325

Combination of automated pre-column trace enrichment and on-line evaporation in HPLC 301(1984)335

Sample enrichment for GC-MS analysis of polynuclear aromatic hydrocarbons in water and in organic mixtures 302(1984)167

Chromatographic analysis of organic compounds in the atmosphere 302(1984)173

Trace enrichment of cytokinins using neutral polystyrene resins 321(1985)175

Trace metals
Trace metal determination by chromatography (Review) 313(1984)129

Trace organics
Identification of trace organic compounds in dimethyl sulphoxide solution using HPLC and GC-MS 324(1985)373

Tracer-pulse chromatography
GC determination of binary adsorption isotherms 302(1984)195

Trachelanthic acid
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

Tragacanth
Identification of food thickeners by monitoring of their pyrolytic products 319(1985)90

Tranexamic acid
Microdetermination of two antifibrinolytic agents (ε-aminocaproic acid and tranexamic acid) by LC and fluorimetric detection (in French) 309(1984)183

Transition metals
Application of indirect potentiometric detection with a metallic copper electrode to ion chromatography of transition metal ions 324(1985)319

Trapping
Rôle of displacement effects in trapping trace components from gases in sorbent-packed columns 324(1985)269

Tremuloidin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Trenbolone
Fast HPLC screening method for the presence of trenbolone and its major metabolite in urine of slaughter cattle 319(1985)436

Triacetin
Silver-modified mobile phase for normal-phase LC determination of prostaglandins and their 5,6-*trans* isomers in prostaglandin bulk drugs and triacetin solutions 321(1985)353

Triacetylfusarinine C
Separation of ferrichromes and other hydroxamate siderophores of fungal origin by reversed-phase chromatography 301(1984)247

Trialkylamines
Effect of trialkylamines on the response of the electron-capture detector to acridine 312(1984)49

Trialkyltin
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341

Speciation of alkyltin compounds by HPLC with a cyanopropyl-bonded silica column 315(1984)351

Triandrin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

Triazines
Parameters determining affinity partitioning of yeast enzymes using polymer-bound triazine dye ligands 303(1984)39

Isotachophoresis of cationic herbicides in waters and soils 320(1985)219

Triazolam
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Triazone
Analysis of candidate anticancer drugs by thermospray HPLC-MS 323(1985)383

Tributyl phosphate
GC analysis of the extraction solvent used in nuclear fuel reprocessing plants 325(1985)75

Tributyltin
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341

Tricarboxylic acids
Separation of tricarboxylic acid cycle acids and other related organic acids in insect haemolymph by HPLC 318(1985)112

Trichloroethanol
Determination of urochloralic acid, the glucuronic acid conjugate of trichloroethanol, by GC with electron-capture detection and its application to urine, plasma and liver 307(1984)111

Trichloroethyl alkanoates
GLC analyses. XXXVI. Separation of ethyl, 2-chloroethyl, 2,2-dichloroethyl and 2,2,2-trichloroethyl esters of aliphatic C_2-C_{20} *n*-alkanoic acids on SE-30 and OV-351 capillary columns with temperature programming 322(1985)97

2,4,4'-Trichloro2'-hydroxydiphenyl ether
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Trichothecenes
Confirmation of nivalenol and deoxynivalenol by on-line LC-MS and GC-MS. Comparison of methods 318(1985)103

Optimisation of conditions for the trimethylsilylation of trichothecene mycotoxins 319(1985)376

Triclocarban
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

Triethylammonium bicarbonate
Preparative separation of nucleotides by ion exchange HPLC using a volatile buffer system 319(1985)461

Triethylenetetramine
Separation and analysis of ethylenediamines as their *tert.* -butyldimethylsilyl derivatives by GLC and MS 314(1984)455

Trifluoroacetic acid
Analytical isotachophoresis utilizing computer simulation. II. Assessment of optimum separation conditions for urinary trifluoroacetic acid metabolized from anaesthetic halothane 305(1984)429

Trifluoroacetic anhydride
Influence of trifluoroacetic anhydride concentration during derivatization and chromatography of N(O,S)-trifluoroacetyl amino acid *n*-propyl esters 318(1985)65

4,4,4-Trifluoro-1-(pentafluoro-phenyl)butane-1,3-dione
Volatility studies of metal chelates. VII. GC behaviour of chelates derived from the aromatic β-diketones 1-phenylbutane-1,3-dione and 4,4,4-trifluoro-1-(pentafluorophenyl)butane-1,3-dione 322(1985)23

Trifluorophosphine carbonyl compounds
GC studies on group VIB organometallic complexes. The analytical and preparative separation of chromium, molybdenum and tungsten trifluorophosphine carbonyl complexes 301(1984)253

Triglycerides
Comparison of detection systems for the HPLC analysis of complex triglyceride mixtures 303(1984)386

Evaluation of the mass detector for quantitative detection of triglycerides and fatty acid methyl esters 324(1985)35

Trihalomethanes
Rapid method for the analysis of trihalomethanes in water 325(1985)221

Triiodothyronine
Use of guanidine hydrochloride in the purification by reversed-phase HPLC of thyroxinyl- and triiodothyronylpeptides derived from thyroglobulin 323(1985)395

Trilostane
Determination of trilostane and ketotrilostane in human plasma by HPLC 311(1984)434

Trimethadione
Simultaneous determination of trimethadione and its metabolite in rat and human serum by HPLC 308(1984)393

Trimethoprim
Determination of trimethoprim metabolites including conjugates in urine using HPLC with combined UV and electrochemical detection 305(1984)391

Simultaneous analysis of three antimicrobial agents in feed premixes by reversed-phase HPLC 323(1985)447

HPTLC determination of trimethoprim and sulphamethoxazole in pharmaceutical dosage forms 323(1985)456

Trimethylamine
Volatile products from acetylcholine as markers in the rapid urine test using head-space GLC 307(1984)11

Trimethylcolchicinic acid
Separation of colchicine and related hydrolysis and photodecomposition products by HPLC, using copper ion complexation 315(1984)233

3,7,11-Trimethyl-11-hydroxy-2,4-dodecadienoic acids
GC of 3,7,11-trimethyl-11-hydroxy (or -methoxy-)-2,4-dodecadienoic acids and related compounds as their methyl or trimethylsilyl esters 303(1984)370

Trimethyllysine
Determination of free trimethyllysine in plasma and tissue specimens by HPLC 306(1984)79

GC-MS characterization of N-methylated basic amino acids in human urine 309(1984)132

3,7,11-Trimethyl-11-methoxy-2,4-dodecadienoic acids
GC of 3,7,11-trimethyl-11-hydroxy (or -methoxy-)-2,4-dodecadienoic acids and related compounds as their methyl or trimethylsilyl esters 303(1984)370

Trimipramine
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Direct LC resolution of racemic drugs by means of α_1-acid glycoprotein as the chiral complexing agent in the mobile phase 316(1984)537

Triolein
Enhanced response of more highly unsaturated lipids on TLC-Iatroscan flame ionization detection after exposing the developed Chromarods to iodine vapour 319(1985)215

Triphenyltin
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341

Tripropyltin
Determination of trialkyltin, dialkyltin, and triphenyltin compounds in environmental water and sediments 315(1984)341

Trisaccharides
HPLC of trisaccharides on amine-bonded silica columns 321(1985)393

Tris(ethylenediamine)cobalt(III) iodide
Substitution-inert metal complex mobile phases in non-suppressed anion chromatography 312(1984)99

Triterpene glycosides
HPLC of oleanane saponins 315(1984)441

Tritiated compounds
Radio-GC of tritiated compounds 322(1985)332

Triton X-100
Supercritical fluid chromatography of various samples with different polarities 315(1984)39

Troleandomycin
Analysis of cortisol, methylprednisolone, and methylprednisolone hemisuccinate. Absence of effects of troleandomycin on ester hydrolysis 305(1984)271

Trypsin
Purification of chymotrypsin and trypsin by CC on agar gel particles 303(1984)283

Isolation of trypsin by column chromatography on tea particles 315(1984)478

Trypsin inhibitor
Kinetics of unfolding of proteins on hydrophobic surfaces in reversed-phase LC 317(1984)227

Tryptamine
Simultaneous determination of tryptamine and its metabolites in mouse brain by HPLC with fluorometric detection 311(1984)385

Tryptophan
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53

Determination of urinary tryptophan, 5-hydroxytryptamine and 5-hydroxyindoleacetic acid in neonatal hyperbilirubinaemic infants using reversed-phase HPLC with fluorescence detection 309(1984)145

Rapid automated analysis of biogenic amines and their metabolites using reversed-phase HPLC with electrochemical detection 309(1984)287

Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

Simultaneous determination of tryptamine and its metabolites in mouse brain by HPLC with fluorometric detection 311(1984)385

HPLC determination of hallucinogenic indoleamines with simultaneous UV photometric and voltammetric detection 320(1985)414

Tryptophol
Simultaneous determination of tryptamine and its metabolites in mouse brain by HPLC with fluorometric detection 311(1984)385

TSKgel DNA-PW
Separation of large DNA restriction fragments by high-performance gel filtration on TSKgel DNA-PW 320(1985)440

D-Tubocurarine chloride
Determination of D-tubocurarine chloride or metocurine iodide in human plasma by HPLC with UV detection 306(1984)398

Tungstate
Capillary-tube isotachophoretic separation of molybdate VI, Tungstate VI and Vanadate V ions using ion pairing equilibria 325(1985)179

Tungsten
GC studies on group VIB organometallic complexes. The analytical and preparative separation of chromium, molybdenum and tungsten trifluorophosphine carbonyl complexes 301(1984)253

Determination of molybdenum and tungsten with tiron by reversed-phase LC 320(1985)403

Turpentines
GLC study of monoterpene hydrocarbons of conifer balsams and ring opening products of pinane and carane structure 321(1985)240

Tween
Effect of surfactants on the separation of proteins by reversed-phase HPLC. I. Non-ionic surfactants (Tween) 317(1984)157

Two-dimensional polyacrylamide gel electrophoresis
Determination of protein spots separated by two-dimensional polyacrylamide gel electrophoresis 319(1985)79

Two-dimensional thin-layer chromatography

Computer simulation as an aid to optimizing continuous-development two-dimensional TLC 302(1984)135

Tyramine
Measurement of tyramine in human plasma, utilising ion-pair extraction and HPLC with amperometric detection 310(1984)11

HPLC with amperometric determination of plasma tyramine 317(1984)319

Tyrosine
Simultaneous measurement of tyrosine, tryptophan and related monoamines for determination of neurotransmitter turnover in discrete rat brain regions by LC with electrochemical detection 305(1984)27

Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Integrated scheme for the simultaneous determination of biogenic amines, precursor amino acids, and related metabolites by LC with electrochemical detection 308(1984)43

Determination of neurotransmitter systems in human cerebrospinal fluid and rat nervous tissue by HPLC with on-line data evaluation 309(1984)53

Analysis of 1- and 3-methylhistidines, aromatic and basic amino acids in rat and human urine 311(1984)267

U

Ubiquinone-10
Effect of sodium dodecyl sulphate on the extraction of ubiquinone-10 in the determination of plasma samples 310(1984)204

Ultrafiltration
Profiles of strongly polar and less polar acids obtained from human blood, plasma and serum by two-step ultrafiltration 309(1984)17

Ultraviolet detector
On-column UV absorption detector for open tubular capillary zone electrophoresis 315(1984)135

Universal liquid chromatography
Universal LC methods. V. "Pulsing" with weak eluent to resolve peaks 321(1985)33

Uracil
Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Separation of 5-fluorouracil and uracil by ion-pair reversed-phase HPLC on a column with porous polymeric packing 307(1984)464

Electrophore-labeling and alkylation of standards of nucleic acid pyrimidine bases for analysis by GC with electron-capture detection 314(1984)201

Uranyl salts
Uranyl salts as fluorescence reagents for the TLC detection of complex-forming compounds (in German) 322(1985)272

Urea
Partitioning of 1-alkanols and other compounds in Sephadex G-10 when eluted with formamide or aqueous solutions of urea, guanidinium salts and some simple electrolytes 301(1984)365

Use of a urea and guanidine-HCl-propanol solvent system to purify a growth inhibitory glycopeptide by HPLC 324(1985)173

Determination of urea and ammonia using ion-pair LC with on-line post-column derivatization in an enzymatic solid-phase reactor 325(1985)255

Urea derivatives
Separation of isocyanate prepolymer components as their urea derivatives by reversed-phase HPLC 301(1984)485

Urea linkages
α-Arylalkylamine-derived chiral stationary phases. Evaluation of urea linkages 322(1985)295

Uric acid
HPLC with polarographic and voltammetric anodic detection: simultaneous determination of allopurinol, oxipurinol and uric acid in body fluids 306(1984)205

Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Uridine monophosphate
Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Uridine phosphorylase
Pyrimidine nucleoside phosphorylase assay by automated HPLC 319(1985)351

Urocanic acid
HPLC determination of (*Z*)- and (*E*)-urocanic acid in human skin 310(1984)188

Urochloralic acid
Determination of urochloralic acid, the glucuronic acid conjugate of trichloroethanol, by GC with electron-capture detection and its application to urine, plasma and liver 307(1984)111

Urokinase
Comparison of low- and high-pressure affinity chromatography for the purification of serine and sulfhydryl esterases 317(1984)193

Uronic acids
Analysis of uronic and aldonic acids, their lactones, and related compounds by HPLC on cation-exchange resins 319(1985)159

Urushiol
Analysis of long-chain phenols in the sap of the Burmese lac tree, *Melanorrhoea usitate* by capillary GLC 318(1985)378

Usnic acid
Determination of preservatives in cosmetic products by reversed-phase HPLC. II. 325(1985)353

UV absorption detector
Simple UV absorption detector for capillary isotachophoresis 325(1985)137

UV-analyzer
The performance of the UVM-4 UV analyser in isotachophoresis 320(1985)241

V

N-Vallinyl-*n*-nonamide
Rapid quality control procedure for the determination of Scoville heat units and the detection of chillies in black pepper, via HPLC 301(1984)288

Valproic acid
Rapid analysis of valproic acid by GC 309(1984)397

Direct determination of valproate in minute whole blood samples 310(1984)208

Improved procedure for the HPLC determination of valproic acid in serum as its phenacyl ester 310(1984)450

Vanadate
Determination of vanadate(V) by conductometric anion chromatography 320(1985)450

Capillary-tube isotachophoretic separation of molybdate VI, Tungstate VI and Vanadate V ions using ion pairing equilibria 325(1985)179

Vanadium chelates
Volatility and GC of some derivatives of Schiff base chelates 318(1985)350

Van der Waals volume
Dependence of the retention of phenols upon Van der Waals volume, π-energy and hydrogen-bonding effects 302(1984)89

Vanillylmandelic acid
Analysis of vanillylmandelic acid, homovanillic acid and 5-hydroxyindoleacetic acid in human urine by HPLC and fluorometry 310(1984)379

Direct determination of urinary vanillylmandelic acid and homovanillic acid by HPLC on an anion-exchange column 311(1984)369

Vapour pressure
Evaluation of a GC method for calculating vapor pressures with organophosphorus pesticides 314(1984)37

Vasoactive intestinal peptide
Recovery of standard ^{125}I-peptides at the picogram and femtogram level by reversed-phase HPLC 312(1984)285

Veatchine
Separation of diterpenoid alkaloid mixtures using the Chromatotron 322(1985)223

Verbascoside
Micromethod for the analysis of heterosidic esters of caffeic acid (in French) 303(1984)312

Vimalin
Chemotaxonomical screening of phenolic glycosides in northern willow twigs by capillary GC 324(1985)129

γ-Vinyl-γ-aminobutyric acid
Assay of γ-vinyl-γ-aminobutyric acid (4-amino-hex-5-enoic acid) in plasma and urine by automatic amino acid analysis. Application to human pharmacokinetics 306(1984)383

HPLC resolution of enantiomers of γ-vinyl-γ-aminobutyric acid 314(1984)495

Vinyl chloride tetramer
Multi-stage chromatographic procedure for the isolation of a single oligomeric species (vinyl chloride tetramer) from poly(vinyl chloride) resin used for food packaging applications 320(1985)361

Vinylidene chloride
Comparison of a new electrochemical detector for GC analysis with the electrolytic conductivity detector 302(1984)269

Viridifloric acid
Paper electrophoresis of hydroxycarboxylic acids as their borate complexes with special reference to viridifloric, trachelanthic, lasiocarpic and heliotric acids. The importance of the "*gem*-dialkyl" effect in complex formation 314(1984)167

Viscosity
Experimental observations of the hydrodynamic behavior of solvent systems in high-speed counter-current chromatography.

III. Effects of physical properties of the solvent systems and operating temperature on the distribution of two-phase solvent systems 301(1984)405

Visnagin
Rapid separation and quantitative determination of khellin and visnagin in *Ammi visnaga* (L.) Lam. fruits by HPLC 301(1984)297

Vitamin A
HPLC of retinals, retinols (vitamin A_1) and their dehydro homologues (vitamin A_2): improvements in resolution and spectroscopic characterization of the stereoisomers 301(1984)65

Analysis of retinoids by HPLC using programmed gradient separation 305(1984)199

Gradient reversed-phase HPLC separation of naturally occurring retinoids 309(1984)299

Quantitative HPLC method for determining the isomer distribution of retinols (vitamin A_1) and 3-dehydroretinol (vitamin A_2) in fish oils 312(1984)423

Determination of vitamin A and vitamin A acetate by HPLC with fluorescence detection 317(1984)349

Vitamin B
New ion-pair reagent for the HPLC separation of B-group vitamins in pharmaceuticals 301(1984)470

Vitamin B_1
Simultaneous analysis of thiamin and riboflavin in foods by HPLC 318(1985)412

Vitamin B_2
Simultaneous analysis of thiamin and riboflavin in foods by HPLC 318(1985)412

Vitamin B_6
LC separation and quantitation of B_6 vitamers in selected rat tissues 306(1984)377

Vitamin C
Fluorometric determination of total vitamin C in whole blood by HPLC with pre-column derivatization 305(1984)53

Vitamin D
Separation of isotopically labeled vitamin D metabolites by HPLC 303(1984)229

Vitamin E
Simultaneous determination by HPLC of tocopherol isomers, α-tocopheryl quinone, and cholesterol in red blood cells and plasma 306(1984)371

Determination of α-tocopherol in human plasma by HPLC with electrochemical detection 311(1984)176

Vitamin K
Determination of (endogenous) vitamin K_1 in human plasma by reversed-phase HPLC using fluorometric detection after post-column electrochemical reduction. Comparison with UV, single and dual electrochemical detection 305(1984)61

Improved method for quantitative analysis of vitamin K_1 and vitamin K_1 2,3-epoxide in human plasma by electron-capture GLC 306(1984)333

Vitamin K_3
Fluorimetric determination of menadione sodium bisulphite (vitamin K_3) in animal feed and premixes by HPLC with post-column derivatization 301(1984)441

Volatile acids
see also Essential oils

Volatiles
GC analysis of volatiles in fluid and gas inclusions 302(1984)181

Studies on volatile metabolites of some potentially pathogenic *Bacillus* species, using automated head-space GC 305(1984)178

Chromatographic profiling of urinary volatile and organic acid metabolites of normal and diabetic C57BL/Ks mice 306(1984)23

W

Wall-jet detector
Modifications of the wall-jet electrochemical detector for LC and flow analysis 316(1984)451

Wall-jet electrochemical flow-cell
Construction of a simple wall-jet electrochemical flow-cell for HPLC 325(1985)294

Warfarin
New method for the resolution of racemic warfarin and its analogues using low-pressure LC 315(1984)378

HPLC strategies for the determination and confirmation of anticoagulant rodenticide residues in animal tissues 321(1985)255

Waxes
GC patterns of some apple surface wax constituents 325(1985)299

Whisker-walled open tubular columns
Influence of phase loading on the performance of whisker-walled open tubular columns coated with organic molten salts 324(1985)415

Whole-cell methanolysis
Whole-cell methanolysis as a rapid method for differentiation between *Actinobacillus*

actinomycetemcomitans and *Haemophilus aphrophilus* 311(1984)347

Wiener numbers
Calculation of retention indices by molecular topology. Chlorinated benzenes 319(1985)1

Window diagram optimization
Window analysis optimization of GC separations using mixed Porapaks 302(1984)119

General theory for correction for compressibility effects in binary coupled GC columns and the procedure for window diagram optimisation of relative lengths 321(1985)249

Withanolides
Reversed-phase chromatographic separation of withanolides from *Acnistus breviflorus* 315(1984)435

Wood components
Single-valve, single-pump, HPLC column-switching analysis of hydrolyzed wood components 318(1985)269

X

XAD-2
Sorption of organic compounds from urine in mutagenicity testing: choice of sorbent 310(1984)41

XAD-4
Production of mutagenic artifacts by the action of residual chlorine on XAD-4 resin 325(1985)95

XAD-7
Trace analysis of benzodiazepine drugs in blood using deactivated Amberlite XAD-7 porous polymer beads and silica capillary column GC with electron-capture detection 301(1984)137

Sorption of organic compounds from urine in mutagenicity testing: choice of sorbent 310(1984)41

Xamoterol
Measurement of xamoterol in plasma and urine by HPLC 308(1984)370

Xanthine
Ion-pair chromatography of selected nucleosides, bases and other low-molecular-weight UV absorbing compounds 307(1984)53

Simultaneous determination of allopurinol, oxipurinol, hypoxanthine and xanthine in biological fluids by HPLC 307(1984)469

Xylenes
Temperature dependence of the solution behaviour of nematic mesophases 324(1985)29

Y

Yohimbine
New method for the determination of yohimbine in biological fluids by HPLC with amperometric detection 311(1984)449

Z

Zeolites
Shape-selective sorption of monomethylalkanes by silicalite, a zeolitic form of silica 316(1984)333

Zimeldine
Reliable routine method for the determination of antidepressant drugs in plasma by HPLC 309(1984)81

Zinc iminodiacetate
Fractionation of human serum proteins by immobilized metal affinity chromatography 315(1984)167

Zingerone
HPLC separation of pungency components of ginger 319(1985)471

Zirconium(IV) oxide
Ion-exchange properties of activated carbon filled with hydrous Zr(IV) and Zr(IV)-P(V) oxides 303(1984)13

Zirconium phosphate
Ion-exchange properties of activated carbon filled with hydrous Zr(IV) and Zr(IV)-P(V) oxides 303(1984)13

Zomepirac
Simultaneous determination of zomepirac and its major metabolite zomepirac glucuronide in human plasma and urine 307(1984)371

Zone electrophoresis
Description and detection of moving sample zones in zone electrophoresis: Zone spreading due to the sample as a necessary discontinuous element 318(1985)455

Zone spreading
Description and detection of moving sample zones in zone electrophoresis: Zone spreading due to the sample as a necessary discontinuous element 318(1985)455

Completely Revised

Instrumental Liquid Chromatography

A Practical Manual on High-Performance Liquid Chromatography Methods

Second, completely revised edition

by **N.A. PARRIS,** E.I. du Pont de Nemours & Company, Biomedical Products Department, Research and Development Division, Experimental Station Laboratory
Wilmington, DE, USA

Journal of Chromatography Library 27

This extensively revised and up-to-date book is an essential tool for the HPLC user in the laboratory. It first appeared in 1976, was twice reprinted and was described in Laboratory Practice as "one of the more useful and successful texts on HPLC . . . a most readable book packed with valuable information and advice . . . strongly recommended."

Practically orientated, it is an easy-to-follow guide containing the minimum essential theoretical background. The majority of the material is based on practical experience and highlights details which may have important operational value for laboratory workers. It helps the HPLC user to select the most appropriate instrumentation, injectors, columns etc.

Applications of liquid chromatography are described with reference to the potential of the technique for qualitative, quantitative and trace analysis as well as for the preparative application. Numerous applications from the literature are tabulated and cross-referenced to sections concerned with the optimization procedures of the particular methods. The format of the original edition proved so successful that it has remained unchanged, but some 45% of the material is either new or completely revised in order to bring the column technology and applications data up-to-date.

"The style . . . is clear. The subject is placed in perspective by comparisons with other separation techniques and should provide a good reference book for all involved in practical LC," (Laboratory Practice).

"Overall the book is well written, and, because of its practical emphasis, it is highly recommended for both the aspiring and experienced chromatographer," (J. Am. Chem. Soc.).

CONTENTS: Fundamentals and Instrumentation. 1. Introduction and historical background. 2. Basic principles and terminology. 3. The chromatographic support and column. 4. Liquid chromatographic instrumentation. 5. Liquid chromatographic detection systems. 6. Modern electronic technology and its impact on LC automation. **Factors Influencing Chromatographic Selectivity.** 7. Nature of the mobile phase. 8. Liquid-solid (adsorption) chromatography. 9. Liquid-liquid (Partition) chromatography. 10. Bonded-phase chromatography. 11. Ion-exchange and ion-pair chromatography. 12. Steric exclusion chromatography. **Uses of Liquid Chromatographic Procedures.** 13. Qualitative analysis. 14. Quantitative analysis. 15. Practical aspects of trace analysis. 16. Practical aspects of preparative liquid chromatography. **Applications of Liquid Chromatography.** 17. Published LC applications. Appendices. List of abbreviations and symbols. Subject index.

1984 xiv + 432 pages. Price: US $ 83.25 / Dfl. 225.00 (including postage).
ISBN 0-444-42061-4

ELSEVIER SCIENCE PUBLISHERS

P.O. Box 211, 1000 AE Amsterdam, The Netherlands
P.O. Box 1663, Grand Central Station, New York, NY 10163, USA

7331

THEORY, PRACTICE AND APPLICATIONS of

MICROCOLUMN HIGH PERFORMANCE LIQUID CHROMATOGRAPHY

P. Kucera, *Pharmaceutical Research Products Section, Hoffmann-La Roche Inc., Nutley, NJ, USA (editor)*

Journal of Chromatography Library 28

This is the only book currently available that covers the large diverse subject of microcolumn chromatographic techniques in such a way as to satisfy, both the practical and the theoretical needs of analytical chemists and chromatographers.

The distinguished research workers and university professors who have contributed to this important work have adopted a textbook type approach to the discussion of the theoretical aspects of new microcolumn techniques. The practical coverage includes instrumentation, design, columns, detectors, injectors, connecting tubing, gradient elution and special analytical techniques, LC-MS, derivatization, etc. and applications are described using various compounds (e.g. drugs, substances of biological origin, proteins, nucleotides, industrial extracts).

The book represents a vast amount of information collected over a period of many years of intensive work and is an essential acquisition for all those who need to keep up-to-date with the latest developments in microcolumn techniques.

"This book gives us many comprehensive statements of current microcolumn HPLC and is highly recommended to chromatographers as well as analytical chemists," (TrAC).

"The vast amount of information . . . makes it essential for all who must keep abreast of latest developments in microcolumn techniques," (Food Technology in Australia).

CONTENTS: Chapter 1. Narrow-Bore and Micro-Bore Columns in Liquid Chromatography (G. Guiochon, H. Colin). 2. Design of a Microbore Column Liquid Chromatograph (P. Kucera, D.D. Dezaro). 3. Theory and Practice of High-Speed Microbore HPLC (R.A. Hartwick, D.D. Dezaro). 4. Special Analytical Techniques (P. Kucera, G. Manius). 5. Chemical Derivatization Techniques Using Microcolumns (P. Kucera, H. Umagat). 6. Applications of Microbore HPLC (P. Kucera, R.A. Hartwick). 7. Liquid Chromatography in Columns of Capillary Dimensions (M. Novotny). 8. Micro LC/MS Coupling (H. Henion). Subject Index.

1984 xvi, 302 pages. Price: US $ 61.00 / Dutch guilders 165.00 (including postage). ISBN 0-444-42290-0

ELSEVIER SCIENCE PUBLISHERS

P.O. Box 211, 1000 AE Amsterdam, The Netherlands
P.O. Box 1663, Grand Central Station, New York, NY 10163, USA

7332

PUBLICATION SCHEDULE FOR 1985

Journal of Chromatography (incorporating *Chromatographic Reviews*) and *Journal of Chromatography, Biomedical Applications*

MONTH	N 1984	D 1984	J 1985	F	M	A	M	J	J	A	S	O	N	D
Journal of Chromatography	312 314	315 316 317	318/1 318/2 319/1	319/2 319/3 320/1	320/2 321/1 321/2 322/1	322/2 322/3 323/1 323/2	324/1 324/2 325/1	325/2 326 327 328	329/1 329/2 329/3	330/1 330/2 331/1	331/2 332 333/1	333/2 CI		
Chromatographic Reviews		313				334/1				334/2			334/3	
Bibliography Section				335/1		335/2		335/3		335/4		335/5		335/6
Biomedical Applications		336/1 336/2	337/1	337/2 338/1	338/2	339/1	339/2 340 341/1	341/2	342/1	342/2	343/1	343/2	344 345/1	345/2

INFORMATION FOR AUTHORS

(Detailed *Instructions to Authors* were published in Vol. 329, No. 3, pp. 449–452. A free reprint can be obtained by application to the publisher.)

Types of Contributions. The following types of papers are published in the *Journal of Chromatography* and the section on *Biomedical Applications:* Regular research papers (Full-length papers), Short communications and Notes. Short communications are preliminary announcements of important new developments and will, whenever possible, be published with maximum speed. Notes are usually descriptions of short investigations and reflect the same quality of research as Full-length papers, but should preferably not exceed four printed pages. For review articles, see page 2 of cover under Submission of Papers.

Submission. Every paper must be accompanied by a letter from the senior author, stating that he is submitting the paper for publication in the *Journal of Chromatography*. Please do not send a letter signed by the director of the institute or the professor unless he is one of the authors.

Manuscripts. Manuscripts should be typed in double spacing on consecutively numbered pages of uniform size. The manuscript should be preceded by a sheet of manuscript paper carrying the title of the paper and the name and full postal address of the person to whom the proofs are to be sent. Authors of papers in French or German are requested to supply an English translation of the title of the paper. As a rule, papers should be divided into sections, headed by a caption (*e.g.*, Summary, Introduction, Experimental, Results, Discussion, etc.). All illustrations, photographs, tables, etc., should be on separate sheets.

Introduction. Every paper must have a concise introduction mentioning what has been done before on the topic described, and stating clearly what is new in the paper now submitted.

Summary. Full-length papers and Review articles should have a summary of 50–100 words which clearly and briefly indicates what is new, different and significant. In the case of French or German articles an additional summary in English, headed by an English translation of the title, should also be provided. (Short communications and Notes are published without a summary.)

Illustrations. The figures should be submitted in a form suitable for reproduction, drawn in Indian ink on drawing or tracing paper. Each illustration should have a legend, all the *legends* being typed (with double spacing) together on a *separate sheet.* If structures are given in the text, the original drawings should be supplied. Coloured illustrations are reproduced at the author's expense, the cost being determined by the number of pages and by the number of colours needed. The written permission of the author and publisher must be obtained for the use of any figure already published. Its source must be indicated in the legend.

References. References should be numbered in the order in which they are cited in the text, and listed in numerical sequence on a separate sheet at the end of the article. Please check a recent issue for the layout of the reference list. Abbreviations for the titles of journals should follow the system used by *Chemical Abstracts.* Articles not yet published should be given as "in press", "submitted for publication", "in preparation" or "personal communication".

Dispatch. Before sending the manuscript to the Editor please check that the envelope contains three copies of the paper complete with references, legends and figures. One of the sets of figures must be the originals suitable for direct reproduction. Please also ensure that permission to publish has been obtained from your institute.

Proofs. One set of proofs will be sent to the author to be carefully checked for printer's errors. Corrections must be restricted to instances in which the proof is at variance with the manuscript. "Extra corrections" will be inserted at the author's expense.

Reprints. Fifty reprints of Full-length papers, Short communications and Notes will be supplied free of charge. Additional reprints can be ordered by the authors. An order form containing price quotations will be sent to the authors together with the proofs of their article.

Advertisements. Advertisement rates are available from the publisher on request. The Editors of the journal accept no responsibility for the contents of the advertisements.

FIRST ANNOUNCEMENT

International Symposium on Coal Characterisation for Conversion Processes
Rolduc, The Netherlands, 28th April - 1st May 1986

Scope
The past decade has been characterized by a renewed interest in coal as an energy source and as a basic material for the chemical industry. This has induced a revival in coal research activities both from a fundamental and an applied point of view. Due to changing economics, however, the large scale introduction of new coal conversion processes has been postponed and, as a consequence, more time is available for research to gain fundamental insights into coal structure and to obtain a better understanding of its impact on conversion processes. In view of this, the initiative has been taken to organize the international "**Rolduc Symposia on Coal Science**". These are intended to fill the need for small-scale working symposia on specific subjects and with a limited number of participants (about 150), as a supplement to the large-scale Coal Science Conferences. The opportunity for in-depth discussions and maximum profit from exchange of knowledge and ideas can thereby be assured.

The first symposium of the series will be devoted to "Coal Characterisation for Conversion Processes". Its aim is to build a bridge between the techniques and methods nowadays available for coal characterisation and the need for relevant parameters to operate and design coal conversion processes.

Scientific Program
The scientific program will cover Coal Characterisation in relation to Gasification, Combustion, Pyrolysis, and Liquefaction. A full symposium day will be devoted to each of the four processes and each day's session will commence with a plenary lecture by an invited expert:
Gasification: **K.H. van Heek** (Bergbau-Forschung GmbH, Germany)
Combustion: **P.A. Roberts** (Int. Flame Research Foundation, The Netherlands)
Pyrolysis: **R. Cyprès** (Free University Brussels, Belgium)
Liquefaction: **C. Snape** (British Gas, U.K.)
Poster sessions and discussions on selected topics will be organized. There are no parallel sessions. The congress language will be English.

Proceedings
The participants will receive a hard-bound edition of the proceedings. This will be a reprint of a special issue of the journal *Fuel Processing Technology*, published by Elsevier.

Fee
A special symposium package, including registration, accommodation for four nights, all meals, and a copy of the proceedings will be available at approximately Dfl. 1050 per participant (about US $ 350).

Further Information
Further information can be obtained from
F. Kapteijn,
Secretary First International Rolduc Symposium on Coal Science,
Institute for Chemical Technology of the University of Amsterdam,
Nieuwe Achtergracht 166,
1018 WV Amsterdam,
The Netherlands.
Telephone 31 - (0) 20-522 3490/2265.
Telex FACWN 16460.